

DIET
LISTS

OF THE
PRESBYTERIAN
HOSPITAL
NEW YORK CITY


—
CARTER

DIET LISTS
OF THE
PRESBYTERIAN HOSPITAL
NEW YORK CITY

COMPILED, WITH NOTES, BY

HERBERT S. CARTER, A. M., M. D.

CONSULTING PHYSICIAN TO THE LINCOLN HOSPITAL; ASSOCIATE IN MEDICINE AT COLUMBIA UNIVERSITY, AND ASSISTANT VISITING PHYSICIAN TO THE PRESBYTERIAN HOSPITAL, NEW YORK CITY



Philadelphia and London

W. B. SAUNDERS COMPANY


1914

RM
216
C27
1913
Larg
Spec

Published May, 1913. Reprinted December, 1913

Copyright, 1913, by W. B. Saunders Company

Reprinted January, 1914



PRINTED IN AMERICA

PRESS OF
W. B. SAUNDERS COMPANY
PHILADELPHIA

PREFACE

THE Diet Lists contained in this volume have been prepared primarily for use in the Presbyterian Hospital; subsequently the comments on the different diets were added, for while the indications for the use of most of the diets is self-evident, the explanatory notes add to the completeness of the presentation and may be found useful to some readers. The Table of Standard Portions was prepared originally by Prof. Irving Fisher, of Yale University, through whose courtesy I am allowed to use the first part of it, while the actual amounts of Protein, Carbohydrate, and Fat contained in each portion was worked out from Bulletin No. 28.

The Table of Food Values has been taken from Bulletin 28, 1906, of the Department of Agriculture, Atwater and Bryant.


The author wishes to express his hearty appreciation for the assistance and advice given him in the preparation of this book by Professor Theodore C. Janeway, also for permission to use certain diet lists prepared by him; to Dr. James C. Greenway, for his courtesy in placing at the author's disposal the diet lists of the New York

Hospital; and to Dr. Herman O. Mosenthal, for some of the diet lists used in the Vanderbilt Clinic, from which great assistance was derived. The author also wishes to express his appreciation to Miss Selma Granat, for the chemical analyses of some of the diets which were made by her in the laboratory of the Presbyterian Hospital.

When diets have been quoted *in toto*, due reference is made to the source of information. The idea of the table of carbohydrate equivalents was obtained from Von Noorden's writings, but adapted to an American dietary by the author.

HERBERT S. CARTER.

NEW YORK CITY.



CONTENTS

	PAGE
INTRODUCTION.....	I
REGULAR HOUSE DIET—CONVALESCENT DIET—SOFT DIET.....	5
Regular House Diet.....	5
Convalescent Diet.....	8
Soft Diet.....	11
Milk Diet.....	11
Liquid Diet.....	11
Comments on Regular House Diet, Convalescent Diet, and Soft Diet.....	11
TYPHOID DIETS.....	14
Typhoid Fluid Diet (No. 1).....	14
Typhoid Fluid Diet (No. 2).....	15
Modified Milk Fluid Diets.....	16
Typhoid Soft Diet.....	18
Comments on Typhoid Fever Diet.....	19
SALT-POOR DIETS.....	22
Salt-poor Diet (No. 1).....	22
Salt-poor Diet (No. 2).....	23
Salt-poor Diet (No. 3).....	24
Comments on Use of Salt-poor Diets.....	24
PURIN-FREE DIETS.....	27
Liquid Purin-free Diet.....	27
Soft Purin-free Diet.....	27
Diet in Gout.....	28
Purin-free Articles of Diet.....	29
The Purin Bodies in Various Food-stuffs.....	30
Comments on Purin-free Diets.....	32

	PAGE
GASTRIC DIETS.....	34
Lenhartz Diet.....	34
Convalescent Gastric Diets (No. 1).....	48
Convalescent Gastric Diets (No. 2).....	48
Comments on Gastric Diets.....	50
GASTRIC HYPERACIDITY.....	55
DIABETIC DIETS.....	58
Standard Strict Diet.....	58
Standard Diet with Restricted Protein.....	59
Green Days.....	60
General Diabetic Diet-lists.....	61
Oatmeal Days.....	63
Comments on Diabetic Diet.....	63
ANTICONSTIPATION DIET.....	70
Comments on Anticonstipation Diet.....	70
LOW CALCIUM DIET.....	72
Comments on Low Calcium Diet.....	72
Fixed Nitrogen, Calcium, Magnesium Diet.....	73
DIARRHEA.....	74
Acute Diarrhea.....	74
Chronic Diarrhea.....	74
OBESITY DIET.....	77
Comments on Obesity Diet.....	77
GASTRIC TEST MEALS.....	79
Comments on Test Meals.....	79
SCHMIDT'S INTESTINAL TEST DIET.....	82
Comments on Schmidt's Intestinal Test Diet.....	83
TABLE OF CARBOHYDRATE EQUIVALENTS.....	84
SPECIAL DIABETIC RECIPES.....	87
Diabetic Bread and Biscuit.....	87
Special Recipes for the Use of Oatmeal.....	88
Soja Bean Meal.....	88

CONTENTS

vii

SPECIAL DIABETIC RECIPES—	PAGE
Desserts.....	90
Diabetic Milk (Wright's).....	91
Analysis of Wright's Diabetic Milk.....	92
MISCELLANEOUS RECIPES.....	93
Protein Milk (Eiweissmilch).....	95
Keller's Malt Soup.....	96
Lactose Ice-cream.....	96
Delafield's Mixture.....	97
AVERAGE WEIGHTS FOR MEN AND WOMEN.....	97
AVERAGE CHEMICAL COMPOSITION OF AMERICAN FOODS.....	98
STANDARD PORTIONS.....	111
APPENDIX.....	119
Karell Treatment for Anasarca.....	119
Comment on Karell Treatment.....	120
Gastric Diet.....	120
Von Leube Treatment for Peptic Ulcer.....	120
INDEX.....	123

DIET LISTS
OF THE
PRESBYTERIAN HOSPITAL,
NEW YORK CITY

INTRODUCTION

WITHOUT attempting to contribute anything new to the science of dietetics, for which the reader is referred to any one of the many excellent works on this subject, the author of the present little volume has in mind merely the presentation of certain diets and the principles governing their use from a purely clinical point of view. Although accuracy, approximately considered, and not exhaustiveness is aimed at, it is hoped that the completeness of the text will be sufficient for the purpose intended.

There is a constantly growing desire on the part of medical men to utilize more exactly the therapeutic effect of foods, but many are deterred by the seeming difficulty of the subject, and by the lack of the time necessary to prescribe an accurate diet list suited to one of the many varieties of disease. In addition,

institutions in which one would expect to find an economy in dietetic therapeutics, as well as accurate methods for arriving at desired results, are, for the most part, sadly lacking in the ability to give their inmates the benefit of a well-balanced ration and their board of managers an economic use of food stuffs. Hospitals and asylums of all sorts should find an easy and rational way to arrive at a satisfactory use of food products.

The needs of an individual who is laid up by a broken leg and one convalescing from an acute or wasting infectious disease are quite different, yet many hospitals have one standard diet for all such cases, nor is allowance made for the different requirements of men and women based on the caloric needs of the body per kilo of body weight.

In the diets here presented advantage has been taken of the facts developed by Professor Chittenden's experiments, that physiologic economy in nutrition can be maintained on a low level of protein, but no attempt has been made to keep to such a low standard, as this is obtained best by a gradual process of reduction in the protein, whereas patients in general are used to much larger amounts, and convalescents are in need of a certain extra quantity of protein to make up the loss due to the destructive katabolism from acute disease. A rational mean has been aimed at in the quantity of protein recommended which will allow a sufficient margin of safety and still be considerably below the

old standards of supposed protein requirement. The advantages to the individual and to the exchequer of the institution in this are obvious, when one considers the high cost of practically all forms of protein food.


In determining the number of heat units for men and women, an average weight of 70 kilos (154 pounds) has been assumed for men and 60 kilos (132 pounds) has been assumed for women as the basis of reckoning, allowing 30 to 32 calories per kilo of body weight, which is ample for an individual at rest or light work. This applies to the regular house diet, soft, gouty, and anticonstipation diets. In the convalescent diet a greater allowance is made, of approximately 35 calories per kilo of body weight. For obvious reasons the diabetic diets have a greater caloric value than the regular hospital diets, as these cases are so habitually undernourished. No hard-and-fast line has been drawn between the requirements of men and women requiring purin-free, low-calcium, or salt-poor diets, as the cases requiring these diets are never numerous in a hospital ward.

The caloric value of the primal food constituents may be stated as follows (Reubner):

	Calories.
1 gm. protein.....	4.1
1 gm. fat.....	9.3
1 gm. carbohydrate.....	4.1
1 gm. alcohol.....	7.0

The caloric needs of the body under varying conditions have been variously estimated, but the following represents the average per kilo of body weight for a man weighing 70 kilos (Von Noorden):

	Calories.
Patient in bed, 30 calories per kilo	2100
Confined to room, 32 to 35 calories per kilo	2240-2450
Light exercise, 35 to 40 calories per kilo	2450-2800
Moderate exercise, 40 to 45 calories per kilo	2800-3150
Very heavy labor, 45 to 60 calories per kilo	3150-4200



REGULAR HOUSE DIET—CONVALESCENT DIET—SOFT
DIET

REGULAR HOUSE DIET¹

EACH DAY OF THE WEEK

TOTAL quantity of milk allowed, not over 750 c.c.
(1½ pints).

Breakfast.

Coffee or tea with milk and sugar, or milk.

Bread and butter.

Two eggs to each patient in male wards.

One egg to each patient in female wards.

Cereal with milk and sugar.

Fresh fish. Hash.

Dinner.

Soup.

Meat or fish.

Potatoes (baked, boiled, or mashed) and one of the following vegetables: spinach, squash, boiled onions, beets, sweet potatoes, macaroni, tomatoes, corn.

Bread and butter.

Milk, 180 c.c. (6 oz.).

Pudding or fruit.

¹ Modified from New York Hospital diet lists.

REGULAR HOUSE DIET

Supper.

Tea or milk. Bread and butter.
 Cooked fruit (prunes, apples, or rhubarb).
 Apricots. Pears.
 Cold meat. Eggs.
 Cereal with milk and sugar. Milk-toast.

Particular Foods for the Specified Days

SUNDAY.

<i>Breakfast.</i>	<i>Dinner.</i>	<i>Supper.</i>
Wheatina.	Chicken.	Cold meat.
Eggs.	Tomatoes.	Prunes.
	Baked potatoes.	
	Fresh fruit.	

MONDAY.

<i>Breakfast.</i>	<i>Dinner.</i>	<i>Supper.</i>
Hominy.	Vegetable soup.	Eggs.
Bacon.	Roast mutton.	Apricots.
	Spinach or corn.	
	Mashed potatoes.	
	Rice pudding.	

TUESDAY.

<i>Breakfast.</i>	<i>Dinner.</i>	<i>Supper.</i>
Oatmeal.	Lamb stew.	Cereal with milk
Eggs.	Boiled potatoes.	and sugar.
	Beets.	Apples.
	Sago pudding.	

WEDNESDAY.

<i>Breakfast.</i>	<i>Dinner.</i>	<i>Supper.</i>
Wheatina.	Chicken.	Cold meat.
Fresh fish.	Baked potatoes.	Rhubarb or prunes.
	Boiled onions.	
	Baked apples.	

THURSDAY.

<i>Breakfast.</i>	<i>Dinner.</i>	<i>Supper.</i>
Hominy.	Pea soup.	Milk-toast.
Eggs.	Roast mutton.	Baked apple.
	Mashed potatoes.	
	Squash.	
	Bread pudding.	

FRIDAY.

<i>Breakfast.</i>	<i>Dinner.</i>	<i>Supper.</i>
Oatmeal.	Fish.	Eggs.
Eggs.	Boiled potatoes.	Prunes.
	Macaroni and tomatoes.	
	Rice pudding.	

SATURDAY.

<i>Breakfast.</i>	<i>Dinner.</i>	<i>Supper.</i>
Cornmeal.	Vegetable soup.	Eggs.
Hash.	Meat stew.	Apricots or pears.
	Mashed potatoes.	
	Spinach or corn.	
	Cornstarch pudding.	

Approximate Values to be Given

	Protein.	Carbohydrate.	Fat.	Total calories.
Men.	90 gm. (3 oz.).	300 gm. (10 oz.).	70 gm. (2½ oz.).	2200
Women..	80 gm. (2¾ oz.).	250 gm. (8½ oz.).	60 gm. (2 oz.).	1800

CONVALESCENT DIET¹

EACH DAY OF THE WEEK

Total quantity of milk allowed, not over 1250 c.c. (2½ pints) daily.

6 A. M.: Milk, 210 c.c. (7 oz.), if desired.

Breakfast.

Coffee or tea with milk and sugar, or milk.

1 egg, or fresh fish, or plain stew.

Cereal with milk and sugar.

Toast and butter or rolls or bread (white, graham or brown).

Fresh fish. Stew.

Dinner.

Broth or soup with barley or vegetables.

Bread and butter. Milk.

Potatoes, baked, boiled, or mashed.

Rice, macaroni, or hominy.

Beef, chicken, or fish.

Pudding, ice-cream, or

Fruit.

¹ Modified from New York Hospital diet lists.

Supper.

Tea or milk. Toast and butter or bread.
 Cooked fruit (apples, stewed or baked, prunes, rhubarb,
 apricots, or pears).

Egg.

8 P. M.: Milk, 210 C.C. (7 oz.).

Particular Foods for the Specified Days

SUNDAY.

<i>Breakfast.</i>	<i>Dinner.</i>	<i>Supper.</i>
Wheatina.	Chicken.	Egg.
1 egg.	Baked potato.	Prunes.
	Orange.	

MONDAY.

<i>Breakfast.</i>	<i>Dinner.</i>	<i>Supper.</i>
Hominy.	Roast-beef.	Egg.
Stew.	Mashed potato.	Pears or apricots.
	Rice pudding.	

TUESDAY.

<i>Breakfast.</i>	<i>Dinner.</i>	<i>Supper.</i>
Oatmeal.	Fresh fish.	Egg.
1 egg.	Boiled potato.	Apples (stewed).
	Hominy.	
	Ice-cream.	

CONVALESCENT DIET

WEDNESDAY.

<i>Breakfast.</i>	<i>Dinner.</i>	<i>Supper.</i>
Wheatina.	Chicken.	Egg.
Fresh fish.	Baked potato.	Rhubarb or prunes.
	Macaroni.	
	Tapioca pudding.	

THURSDAY.

<i>Breakfast.</i>	<i>Dinner.</i>	<i>Supper.</i>
Hominy.	Boiled beef.	Egg.
1 egg.	Mashed potato.	Baked apple.
	Rice.	
	Baked custard.	

FRIDAY.

<i>Breakfast.</i>	<i>Dinner.</i>	<i>Supper.</i>
Oatmeal.	Fresh fish.	Egg.
Egg.	Boiled potato.	Prunes.
	Macaroni.	
	Ice-cream.	

SATURDAY.

<i>Breakfast.</i>	<i>Dinner.</i>	<i>Supper.</i>
Cornmeal.	Chicken.	Egg.
Stew.	Mashed potato.	Apricots or pears.
	Hominy.	
	Cornstarch pudding.	

Approximate Values to be Given

	Protein.	Carbohydrate.	Fat.	Total Calories.
Men . . .	100 gm. (3½ oz.).	300 gm. (10 oz.).	90 gm. (3 oz.).	2500
Women..	80 gm. (2¾ oz.).	300 gm. (10 oz.).	80 gm. (2¾ oz.).	2200

SOFT DIET

Soft diet is the same as the convalescent diet, omitting meat, fish, and green vegetables.

Approximate Values to be Given

	Protein.	Carbohydrate.	Fat.	Total Calories.
Men...60 gm. (2 oz.).	350 gm. (11 $\frac{2}{3}$ oz.).	60 gm. (2 oz.).	2200	
Women..50 gm. (1 $\frac{2}{3}$ oz.).	300 gm. (10 oz.).	50 gm. (1 $\frac{2}{3}$ oz.).	1800	

MILK DIET

Milk, 3000 c.c. (3 quarts). This gives: Protein, 100 gm. (3 $\frac{1}{3}$ oz.); fat, 120 gm. (4 oz.); carbohydrate, 150 gm. (5 oz.); calories, 2160; chlorids, 5.2 gm.

LIQUID DIET

Coffee.....	165 c.c. (5 $\frac{1}{2}$ oz.)
Broth.....	675 c.c. (22 $\frac{1}{2}$ oz.)
Gruel.....	200 c.c. (6 $\frac{2}{3}$ oz.)
Tea.....	200 c.c. (6 $\frac{2}{3}$ oz.)
Milk.....	600 c.c. (20 oz.)
Sugar.....	60 gm. (2 oz.)
Egg-albumin.....	90 gm. (3 oz.)
Lemon-juice.....	60 gm. (2 oz.)

This gives: Protein, 47 gm. (1 $\frac{1}{2}$ oz.); fat, 28 gm. (1 oz.); carbohydrate, 108 gm. (3 $\frac{1}{3}$ oz.); calories, 900; chlorids, 2.6 gm.

COMMENTS ON REGULAR HOUSE DIET, CONVALESCENT DIET, AND SOFT DIET

Regular house diet, convalescent diet, and soft diet speak for themselves, and are of particular use in hospi-

tal or institutional dietaries. The variety of foods included is considerable, and should one desire to keep institutional expenses at a lower level, this can be easily done by substituting the less expensive food stuffs, keeping, however, to the same general plan. Convalescent diet should be of a higher nutritive value, more easily digestible, and contain more protein, and this change from regular house diet is indicated.

Soft diet, representing an intermediate step between fluid or milk diet and convalescent or regular diet, need not be so strictly adhered to in point of food values, as a patient's appetite can be relied upon at this stage to take a sufficient quantity.

For institutions, however, it is well to have this also standardized for economy in planning.

Fluid diet, while it contains a fair amount of protein food, is of exceedingly low caloric value, and represents what may be well to give during the first day or two of any acute illness, when one wishes to rest the organism without subjecting it to protein starvation.

Milk diet, except as modified upward in the typhoid milk diets, is of use for a time, particularly in an acute or occasionally in a chronic renal condition, its great objection being that, in order to give the necessary calories for a man, one must give large amounts of milk, 3000 to 4000 c.c. (3 to 4 quarts), which would mean that the individual must consume 100 to 130 gm. (3 to 4 oz.) of protein, an unnecessary strain to put

upon excretory organs already disabled by disease. So that, while we give not more than 3000 c.c. (3 quarts) per day for milk diet, it must be remembered that this should not be indefinitely continued.



TYPHOID DIETS

TYPHOID FLUID DIET (NO. 1)¹

- 8 A. M.: Milk and coffee, each 120 c.c. (4 oz.), 240 c.c. (8 oz.).
- 10 A. M.: Milk, hot or cold, 240 c.c. (8 oz.).
- 12 NOON: Oatmeal gruel, 120 c.c. (4 oz.), with milk, 60 c.c. (2 oz.).
- 2 P. M.: Milk, 240 c.c. (8 oz.).
- 4 P. M.: Oatmeal gruel, 120 c.c. (4 oz.), with milk, 60 c.c. (2 oz.).
- 6 P. M.: Custard with lactose (full cup).
- 8 P. M.: Hot milk, 240 c.c. (8 oz.).
- 10 P. M.: Whey, 180 c.c. (6 oz.), with one whole egg and sherry.
- 12 P. M.: Oatmeal gruel, 120 c.c. (4 oz.), milk, 60 c.c. (2 oz.).
- 2 A. M.: Milk, 240 c.c. (8 oz.).
- 4 A. M.: Broth, 240 c.c. (8 oz.), with one whole egg.
- 6 A. M.: Milk, 240 c.c. (8 oz.).

Approximate Values

Protein, 98 gm. ($3\frac{1}{3}$ oz.); fat, 52 gm. ($1\frac{2}{3}$ oz.); carbohydrates, 150 gm. (5 oz.); calories, 1900.

¹F. P. Kinnicutt.

TYPHOID FLUID DIET (NO. 2)¹*(Lower Protein)*

- 8 A. M.: Milk and coffee, each 120 c.c. (4 oz.).
 10 A. M.: Milk, hot or cold, 240 c.c. (8 oz.).
 12 NOON: Oatmeal gruel, 120 c.c. (4 oz.), with milk,
 60 c.c. (2 oz.).
 2 P. M.: Junket with cane- and milk-sugar.
 4 P. M.: Oatmeal gruel, 120 c.c. (4 oz.), with milk, 60
 c.c. (2 oz.).
 6 P. M.: Junket with cane- and milk-sugar.
 8 P. M.: Hot milk, 240 c.c. (8 oz.).
 10 P. M.: Whey, 180 c.c., with one whole egg and sherry.
 12 P. M.: Oatmeal gruel, 120 c.c. (4 oz.), with milk,
 60 c.c. (2 oz.).
 2 A. M.: Junket with cane- and milk-sugar.
 4 A. M.: Milk, 240 c.c. (8 oz.).
 6 A. M.: Milk, 240 c.c. (8 oz.).
 15 gm. ($\frac{1}{2}$ oz.) of lactose added to the four milk feedings.

Approximate Values

Protein, 71 gm. ($2\frac{1}{3}$ oz.); fat, 81 gm. ($2\frac{2}{3}$ oz.); carbo-
 hydrates, 160 gm. ($5\frac{1}{3}$ oz.); calories, 2300.

These typhoid fluid diets can be further augmented by
 the addition of the following articles:

- Lactose or cane-sugar, 30 gm. (1 oz.), 120 calories.
 Pea soup (see Special Recipes), 180 c.c. (1 cup, 6 oz.),
 192 calories (7.5 gm. protein).

¹ F. P. Kinnicutt.

Bean soup (see Special Recipes), 180 c.c. (1 cup, 6 oz.),
242 calories (12 gm. protein).

Soda cracker (one), 30 calories (.5 gm. protein).

Cream.....	100 c.c. ($3\frac{1}{3}$ oz.).	} 500 calories.
Sugar or lactose....	60 gm. (2 oz.).	
Gruel.....	150 c.c. (5 oz.).	

This adds protein, 5 gm. ($\frac{1}{8}$ oz.); fat, 20 gm. ($\frac{2}{3}$ oz.);
carbohydrates, 75 gm. ($2\frac{1}{2}$ oz.).

MODIFIED MILK FLUID DIETS¹

For 1000 calories a day: Calories.

Milk, 1000 c.c. (1 quart).....	700
Cream, 50 c.c. ($1\frac{2}{3}$ oz.).....	100
Lactose, 50 gm. ($1\frac{2}{3}$ oz.).....	200

This furnishes eight feedings, each containing:

Milk, 120 c.c. (4 oz.).....	80
Cream, 8 gm. (2 dr.).....	15
Lactose, 6 gm. ($1\frac{1}{2}$ dr.).....	24

For 1500 calories a day:

Milk, 1500 c.c. ($1\frac{1}{2}$ quarts).....	1000
Cream, 50 c.c. ($1\frac{2}{3}$ oz.).....	100
Lactose, 100 gm. ($3\frac{1}{3}$ oz.).....	400

This furnishes six feedings, each containing:

Milk, 240 c.c. (8 oz.).....	160
Cream, 8 gm. (2 dr.).....	15
Lactose, 16 gm. (4 dr.).....	64

¹ Coleman, American Journal of Medical Sciences, January, 1912.

For 2000 calories a day:

	Calories.
Milk, 1500 c.c. ($1\frac{1}{2}$ quarts).....	1000
Cream, 240 c.c. (8 oz.).....	500
Lactose, 120 gm. (4 oz.).....	500

This furnishes seven feedings, each containing:

Milk, 210 c.c. (7 oz.).....	140
Cream, 30 c.c. (1 oz.).....	60
Lactose, 18 gm. ($4\frac{1}{2}$ dr.).....	72

For 2500 calories a day:

Milk, 1500 c.c. ($1\frac{1}{2}$ quarts).....	1000
Cream, 240 c.c. (8 oz.).....	500
Lactose, 240 gm. (8 oz.).....	1000

This furnishes seven feedings, each containing:

Milk, 210 c.c. (7 oz.).....	140
Cream, 30 c.c. (1 oz.).....	60
Lactose, 36 gm. (9 dr.).....	144

For 3000 calories a day:

Milk, 1500 c.c. ($1\frac{1}{2}$ quarts).....	1000
Cream, 480 c.c. (1 pint).....	1000
Lactose, 240 c.c. (8 oz.).....	1000

This furnishes eight feedings, each containing:

Milk, 180 c.c. (6 oz.).....	120
Cream, 60 c.c. (2 oz.).....	120
Lactose, 30 gm. (1 oz.).....	120

For 3900 calories a day:

Milk, 1500 c.c. ($1\frac{1}{2}$ quarts).....	1000
Cream, 480 c.c. (1 pint).....	1000
Lactose, 480 gm. (16 oz.).....	1900

This furnishes eight feedings, each containing:		Calories
Milk, 180 c.c. (6 oz.)	120
Cream, 60 c.c. (2 oz.)	120
Lactose, 60 gm. (2 oz.)	240

TYPHOID SOFT DIET¹

Use cream, 40 per cent. (16 oz.), for each patient (Q. D.):

	Calories.
6.00 A. M. to 8.00 P. M.: Pasteurized milk, 180 c.c. (6 oz.)	120
Cream, 30 c.c. (1 oz.)	100
Each glass	220
Q. 2 H. Eight glasses	1760

Breakfast.

	Calories.
1 soft-boiled egg	60
2 tablespoonfuls hominy or farina (well cooked).	100
With milk, 90 c.c. (3 oz.)	60
With cream, 60 c.c. (2 oz.)	200
	420

Dinner.

	Calories.
Beef-juice, scraped beef, each 30 c.c. (1 oz.)	50
Oatmeal gruel, 240 c.c. (8 oz.)	100
With cream, 60 c.c. (2 oz.)	200
3 tablespoonfuls ice-cream	200
	550

¹ L. A. Conner, New York Hospital Diets.

	<i>Supper.</i>	Calories.
1	soft-boiled egg.....	60
3	tablespoonfuls wine-jelly or custard, or oatmeal gruel, 240 c.c. (8 oz.).....	100
	With cream, 60 c.c. (2 oz.).....	<u>300</u>
		460

Approximate values of this diet are protein, 101 gm. ($3\frac{1}{3}$ oz.); carbohydrate, 172 gm. ($5\frac{2}{3}$ oz.); fat, 176 gm. (6 oz.); calories, 3190.

COMMENTS ON TYPHOID FEVER DIET

Many cases of typhoid when seen for the first time, either in the hospital or private practice, have been ill for a number of days, and are very apt to have a diarrhea, which is not, strictly speaking, a part of their disease, but a complication, brought on by improper feeding. In such cases it is best, after the initial emptying of the intestinal canal by mild catharsis or enemata, to prescribe a diet which, while it is too low in heat units, still has a fair amount of protein, and has an added advantage over a strictly milk diet in that it is quite varied in composition. Such a diet is given in the typhoid liquid diets, and, while one could often advantageously give the ordinary fluid diet for this period, the special typhoid liquid diet furnishes almost twice as much protein and twice as many calories.


When the patient is ready for more food this list of

articles can be further reinforced by the addition of cream, lactose, eggs, and more gruel, or one can make use of one of the cream, milk, and lactose mixtures given in the diet lists devised by Dr. Warren Coleman. There is, however, a distinct advantage in the greater variety of foods in the first list, and it has been the experience of many that it is better borne. In early convalescence the soft typhoid diet will be found convenient, as representing a good amount of protein and a larger number of calories without irritating or coarse residue. There is apparently no question among clinicians nowadays that typhoid patients do better on a more liberal supply and variety of foods than on a monotonous and insufficient milk diet—complications are fewer (?) and convalescence is shortened. Coleman and Schaefer have ably demonstrated that the body is able, even in high fever, to digest, absorb, and properly utilize quantities of food sufficiently large to maintain nitrogenous equilibrium and body weight, their cases often actually gaining weight during the fever.

There never was a class of patients, however, who needed more individual care in the selection of the kind and quantity of food given than typhoid fever patients, and, while the high caloric feeding is a great step in advance, it must be individualized for the patient and always arrived at gradually; *i. e.*, it is not well to go from a low caloric to a high caloric diet without a gradual increase from one to the other.

Just here it might be added that when a patient's digestion fails to take care of the larger amounts of food, as shown by tympanites, flatulence, etc., the use of buttermilk or milk artificially ripened by some form of the lactic acid bacillus (see Special Recipes), if given in moderate amount wherever fresh milk is being used, will often stop the intestinal fermentation at once. It is almost as useful when a patient's stomach becomes disturbed, as shown by nausea or vomiting.

The transition from typhoid soft to convalescent diet is easily made, although fruits and green vegetables should be excluded at first.



SALT-POOR DIETS
SALT-POOR DIET NO. 1

<i>Breakfast.</i>	Gm.	Oz.
Bread.....	30	1
Sugar.....	10	$\frac{1}{3}$
Farina.....	60	2
Butter.....	30	1
Egg (1).....	40	$1\frac{1}{3}$
Coffee.....	150 c.c.	5
	320	$10\frac{2}{3}$

<i>Dinner.</i>	Gm.	Oz.
Bread.....	30	1
Butter.....	20	$\frac{2}{3}$
Sugar.....	10	$\frac{1}{3}$
Rice.....	60	2
Farina.....	100	$3\frac{1}{3}$
Tea.....	150 c.c.	5
	370	$12\frac{1}{3}$

<i>Supper.</i>	Gm.	Oz.
Egg (1).....	40	$1\frac{1}{3}$
Toast.....	15	$\frac{1}{2}$
Bread.....	30	1
Butter.....	15	$\frac{1}{2}$
Custard.....	100	$3\frac{1}{3}$
Prunes.....	60	2
Tea.....	180 c.c.	6
	440	$14\frac{2}{3}$

This contains chlorids, 1 gm.; protein, 35 gm. ($1\frac{1}{8}$ oz.); fat, 65 gm. ($2\frac{1}{8}$ oz.); carbohydrate, 140 gm. ($4\frac{2}{3}$ oz.); calories, 1300.

SALT-POOR DIET NO. 2

<i>Breakfast.</i>		
	Gm.	Oz.
Bread.....	60	2
Sugar.....	40	$1\frac{1}{3}$
Farina.....	60	2
Butter.....	35	$1\frac{1}{6}$
Egg (1).....	40	$1\frac{1}{3}$
Coffee.....	150 c.c.	5
	<hr/>	
	385	$12\frac{5}{8}$
<i>Dinner.</i>		
	Gm.	Oz.
Egg (1).....	40	$1\frac{1}{3}$
Bread.....	60	2
Butter.....	30	1
Rice.....	70	$2\frac{1}{3}$
Farina.....	100	$3\frac{1}{3}$
Tea.....	150 c.c.	5
	<hr/>	
	450	15
<i>Supper.</i>		
	Gm.	Oz.
Egg (1).....	40	$1\frac{1}{3}$
Toast.....	15	$\frac{1}{2}$
Bread.....	60	2
Butter.....	30	1
Custard.....	100	$3\frac{1}{3}$
Prunes.....	60	2
Tea.....	180 c.c.	6
	<hr/>	
	485	$15\frac{5}{8}$

This contains chlorids, 1.3 gm; protein, 50 gm. ($1\frac{2}{3}$ oz.); fat, 100 gm. ($3\frac{1}{2}$ oz.); carbohydrate, 240 gm. (8 oz.); calories, 2100.

SALT-POOR DIET NO. 3

Same as convalescent diet, without broths or soups. The fish, meat, and green vegetables must be boiled in two waters to remove most of the salt. Milk, 250 c.c. (8 oz.), only allowed.¹

COMMENTS ON USE OF SALT-POOR DIETS

In the use of salt-poor diets, or so-called salt-“free” diets (although none of them can be actually salt free), a great step in advance has been made in the dietetic treatment of edema, particularly, but not exclusively, of renal origin. We are more indebted to Widal than to any other one man for putting this therapeutic measure on a firm basis. Of course, it is particularly useful in edema associated with a greater or lesser degree of impermeability of the kidney to salt. The two requisites for the successful use of this form of treatment are, proper selection of cases and the continued and faithful use of a diet that is as nearly salt free as possible. In other words, while the edema in many cases of this kind begins to clear up at once on this diet (although it has resisted all other physical and medical measures),

¹ In all the salt-poor diets the food should be prepared absolutely without addition of salt. Salt-free butter and bread only are to be used. No extra salt allowed.

in many of them it does not, and it is not until the patients have been on this diet for a week or two or longer that apparently the kidneys, having been spared so long, are able to regain their lost power of salt excretion, and will then go ahead and rapidly empty the subcutaneous tissue of its retained salt and water.

While it is best to determine the rapidity of the salt elimination in cases in which the edema resists the dietary, so that one can be assured that one is on the right track, a clinical diagnosis can often be made when the edema is diminished promptly. While, as already stated, this treatment is particularly indicated in a special class of renal disease, it is of great assistance in many forms of edema or even serous effusions. This is due no doubt to the fact that the kidneys are spared extra labor in the excretion of the usual amount of salt taken with the normal diet, and to the fact that the body, in the presence of a reduced salt intake, seeks salt in the tissues or serous spaces in order to keep up the normal salt concentration in the urine, as an attempt at equilibrium of the inorganic constituents of the urine is in some way automatically arranged for.

With this foreword, the use of the salt-poor diets becomes evident and needs little or no further explanation.

No. 1 should be used first, then No. 2, and finally, when safe, No. 3. The length of time each should be used is a matter for individual judgment, but No. 2

should not be used, if possible, until, all edema is gone, and has been absent long enough presumably to give the kidney rest.¹

¹Determination of total chlorids in urine: Dilute 10 c.c. of urine with 90 c.c. of water, add 1 or 2 drops of 25 per cent. nitric acid. Make the mixture alkaline with 10 per cent. solution of sodium carbonate, and add a few drops of 10 per cent. potassium chromate solution as an indicator. Titrate with $\frac{1}{10}$ normal silver chlorid solution. Each cubic centimeter of the silver solution used equals .00583 gm. sodium chlorid.



PURIN-FREE DIETS

LIQUID PURIN-FREE DIET

(FOLIN)

Whole milk.....	500 c.c. (16 oz.).
Cream (18 to 22 per cent. fat)...	300 c.c. (10 oz.).
10 eggs (whites and yolks).....	450 gm. (15 oz.).
Horlick's malted milk.....	200 gm. (6 $\frac{2}{3}$ oz.).
Sugar.....	20 gm. ($\frac{2}{3}$ oz.).
NaCl.....	6 gm. ($\frac{1}{8}$ oz.).
Water.....	q. s. ad. 2000 c.c. (64 oz.).
Extra water.....	900 c.c. (30 oz.).

Yields

Protein.....	139 gm. (4 $\frac{2}{3}$ oz.)	} 2830 calories.
Fat.....	146 gm. (4 $\frac{5}{8}$ oz.)	
Carbohydrate... ..	225 gm. (7 $\frac{1}{2}$ oz.)	

SOFT PURIN-FREE DIET¹

6 A. M.: Milk, 180 c.c. (6 oz.).

8 A. M.—Breakfast:

Milk, 180 c.c. (6 oz.).

1 $\frac{1}{2}$ slices of bread and 1 pat of butter.

2 tablespoonfuls wheatina or cream of wheat with cream, 60 c.c. (2 oz.), and 2 tablespoonfuls of sugar.

1 soft-boiled egg.

¹ Vanderbilt Clinic diet lists.

12 M.—Dinner:

Milk, 180 c.c. (6 oz.).

1 soft-boiled egg, S. O. S.

Potato with cream, 30 c.c. (1 oz.), and 1 pat of butter.

Lettuce or cabbage with dressing.

1½ slices of bread with 1 pat of butter.

3 P. M.: Milk, 180 c.c. (6 oz.).

5 P. M.—Supper:

1 soft-boiled egg, S. O. S.

Milk, 180 c.c. (6 oz.).

2½ tablespoonfuls of rice with cream, 30 c.c. (1 oz.), and 1 tablespoonful of sugar.

Crackers with butter, 1 pat.

1 cube of cheese (2 inches).

1 cup of weak tea with cream, 30 c.c. (1 oz.), and 1 teaspoonful of sugar, S. O. S.

8 P. M.: Milk, 180 c.c. (6 oz.).

Protein, 80 gm. (2⅔ oz.); fat, 112 gm (3½ oz.); carbohydrate, 207 gm. (7 oz.); calories, 2300.

DIET IN GOUT¹

Breakfast:

Any fresh, cooked, or preserved fruit.

Cereal with sugar and milk or cream.

White bread, toast, muffins, biscuits, butter.

2 eggs (any style).

Milk, with slight amount of coffee, tea, or cocoa.

¹ Modified from Von Noorden.

Lunch and Dinner:

Soups.—Cream, potato, or from any vegetable given below, prepared without meat or meat extract.

Eggs, any style.

Meat and Fish.—Once a day one of the following:

Ham, boiled beef, boiled chicken, boiled mutton, pot roast, any boiled fish.

Vegetables.—Potatoes, sweet potatoes, cauliflower, lettuce, cabbage, spinach, okra, egg-plant, corn, Brussels sprouts, rice, macaroni, noodles.

Bread.—Crackers, white bread, toast, butter.

Desserts.—Fresh, cooked, or preserved fruit, all desserts (ice-cream, pudding, cake, pie, etc.) not flavored with coffee or chocolate, American, Swiss, and cream cheese.

Beverages.—Milk, cider, water, Vichy or any carbonated water. Grape-juice and raspberry syrup allowed.

Approximate Values to be Given

	Protein.	Carbohydrate.	Fat.	Total calories.
Men...	.60 gm. (2 oz.).	350 gm. (11 $\frac{3}{4}$ oz.).	60 gm. (2 oz.).	2200
Women...	.50 gm. (1 $\frac{3}{4}$ oz.).	300 gm. (10 oz.).	50 gm. (1 $\frac{3}{4}$ oz.).	1800

PURIN-FREE ARTICLES OF DIET¹

Milk.

Eggs.

Bread, white only (not graham or whole wheat).

Butter.

¹ Vanderbilt Clinic diet lists.

- Biscuits.
 Cereals (hominy, rice, farina).
 Cream.
 Sugar and syrup.
 Jam and marmalade.
 Cake.
 Cream soups.
 Vegetables:
 Potatoes (slight amount of purin).
 Cauliflower.
 Cabbage.
 Lettuce.
 Egg-plant.
 Desserts:
 Nuts.
 Cheese.
 Ice-cream and water-ices.
 Cake (any cake except with coffee or chocolate
 flavor).
 Rice, bread, farina, tapioca, or cornstarch pudding.
 Custard or cocoanut pie.

THE PURIN BODIES IN VARIOUS FOOD-STUFFS ¹

Fish:	Purin bodies, grams per kilo.
Cod.....	0.5
Plaice.....	0.7
Salmon.....	1.1
Halibut.....	1.0

¹ Hall J. Walker, *The Purin Bodies of Food Stuffs and the Rôle of Uric Acid in Health and Disease*, 2d ed. rev., London, 1903.

Meat:	Purin bodies, grams per kilo.
Beef.....	1.3-2.0
Fat.....	1.1
Mutton.....	0.96
Fat.....	—
Veal.....	1.1
Fat.....	—
Pork.....	1.2
Fat.....	0.5
Ham.....	1.1

Meat soups (varying large amounts).

Chicken.....	1.2
--------------	-----

Vegetables:

Potatoes.....	0.02
Rice.....	—
Flour (white).....	—
Bread (white).....	—
Oatmeal.....	0.53
Peas.....	0.39
Lentils.....	0.38
Beans (Haricot).....	0.63
Asparagus.....	0.21
Cabbage.....	—
Lettuce.....	—
Cauliflower.....	—
Onions.....	0.09
Tapioca.....	—

Special Foods:	Purin bodies, grams per kilo.
Milk.....	—
Butter.....	—
Eggs.....	—
Cheese (fat).....	—

Drinks:

Beer, lager.....	0.12
Ale.....	0.14
Porter.....	0.15

	Per pint (500 c.c.).
Tea.....	1.2
Cocoa.....	1.0
Chocolate.....	0.7
Coffee.....	1.7
Claret.....	—
Sherry.....	—
Port.....	—

COMMENTS ON PURIN-FREE DIETS


Since in gout the purin metabolism is faulty, the aim of its dietetic treatment should be to furnish a food containing little or no purin bodies.

In the liquid purin-free diet, as designed by Folin, we have a diet that is useful almost exclusively for scientific purposes, when one wishes to determine the amount of endogenous uric acid that is excreted. It can, however, be used to advantage during an acute attack of gout.

The soft purin-free diet can, for the most part, be used in acute gout, and is much more palatable. The amount of protein or the total calories can be reduced or increased at will to meet the requirements of large or small individuals. The list given is sufficient for a man of 70 kilos (154 pounds). Women seldom have acute gout.

The list of purin-free articles of diet is inserted to help one prescribe a variety in case the diet is continued for some time.

The diet given specifically for gout can be modified from the purin-free list as well, and, while this diet for gout is not absolutely purin free, it is of a low purin content, and boiling the meats, particularly if they are boiled in two waters, removes a great part of the purins.



GASTRIC DIETS

LENHARTZ DIET

As eggs differ in size and weight, take the total of eggs for the day of diet; beat, measure, and divide into seven feedings and put into medicine-glasses. Keep on ice and use as directed, alternating with milk.

Milk, iced, kept in bowl of cracked ice. Eggs beaten up raw and iced. Spoon kept in bowl of ice.

Patients never allowed to help themselves.

Slow feedings essential, frequent small feedings fed by spoon.

First and second days salt eggs to taste.

Third day sugar is started.

First Day

- 7 A. M.: Egg.
- 8 A. M.: Milk, 20 c.c. ($\frac{2}{3}$ oz.).
- 9 A. M.: Egg.
- 10 A. M.: Milk, 20 c.c. ($\frac{2}{3}$ oz.).
- 11 A. M.: Egg.
- 12 NOON: Milk, 15 c.c. ($\frac{1}{2}$ oz.).
- 1 P. M.: Egg.
- 2 P. M.: Milk, 15 c.c. ($\frac{1}{2}$ oz.).
- 3 P. M.: Egg.
- 4 P. M.: Milk, 15 c.c. ($\frac{1}{2}$ oz.).

5 P. M.: Egg.

6 P. M.: Milk, 15 c.c. ($\frac{1}{2}$ oz.).

7 P. M.: Egg.

Total First Day.—Eggs (raw), 2; milk, 100 c.c. ($3\frac{1}{2}$ oz.).

Second Day

7 A. M.: Egg.

8 A. M.: Milk, 35 c.c. (1 oz.).

9 A. M.: Egg.

10 A. M.: Milk, 35 c.c. (1 oz.).

11 A. M.: Egg.

12 NOON: Milk, 35 c.c. (1 oz.).

1 P. M.: Egg.

2 P. M.: Milk, 35 c.c. (1 oz.).

3 P. M.: Egg.

4 P. M.: Milk, 30 c.c. (1 oz.).

5 P. M.: Egg.

6 P. M.: Milk, 30 c.c. (1 oz.).

7 P. M.: Egg.

Total Second Day.—Eggs (raw), 3; milk, 200 c.c. ($6\frac{2}{3}$ oz.).

Third Day

7 A. M.: Egg.

Sugar, 2 gm. ($\frac{1}{2}$ dr.).

8 A. M.: Milk, 50 c.c. ($1\frac{2}{3}$ oz.).

9 A. M.: Egg.

Sugar, 3 gm. ($\frac{3}{4}$ dr.).

10 A. M.: Milk, 50 c.c. ($1\frac{2}{3}$ oz.).

11 A. M.: Egg.

Sugar, 3 gm. ($\frac{3}{4}$ dr.).

12 NOON: Milk, 50 c.c. ($1\frac{2}{3}$ oz.).

- 1 P. M.: Egg.
 Sugar, 3 gm. ($\frac{3}{4}$ dr.).
- 2 P. M.: Milk, 50 c.c. ($1\frac{2}{3}$ oz.).
- 3 P. M.: Egg.
 Sugar, 3 gm. ($\frac{3}{4}$ dr.).
- 4 P. M.: Milk, 50 c.c. ($1\frac{2}{3}$ oz.).
- 5 P. M.: Egg.
 Sugar, 3 gm. ($\frac{3}{4}$ dr.).
- 6 P. M.: Milk, 50 c.c. ($1\frac{2}{3}$ oz.).
- 7 P. M.: Egg.
 Sugar, 3 gm. ($\frac{3}{4}$ dr.).

Total Third Day.—Eggs (raw), 4; milk, 300 c.c. (10 oz.); sugar, 20 gm. (5 dr.).

Fourth Day

- 7 A. M.: Egg.
 Sugar, 2 gm. ($\frac{1}{2}$ dr.).
- 8 A. M.: Milk, 70 c.c. ($2\frac{1}{3}$ oz.).
- 9 A. M.: Egg.
 Sugar, 3 gm. ($\frac{3}{4}$ dr.).
- 10 A. M.: Milk, 70 c.c. ($2\frac{1}{3}$ oz.).
- 11 A. M.: Egg.
 Sugar, 3 gm. ($\frac{3}{4}$ dr.).
- 12 NOON: Milk, 65 c.c. (2 oz.).
- 1 P. M.: Egg.
 Sugar, 3 gm. ($\frac{3}{4}$ dr.).
- 2 P. M.: Milk, 65 c.c. (2 oz.).
- 3 P. M.: Egg.
 Sugar, 3 gm. ($\frac{3}{4}$ dr.).
- 4 P. M.: Milk, 65 c.c. (2 oz.).
- 5 P. M.: Egg.
 Sugar, 3 gm. ($\frac{3}{4}$ dr.).

6 P. M.: Milk, 65 c.c. (2 oz.).

7 P. M.: Egg.

Sugar, 3 gm. ($\frac{3}{4}$ dr.).

Total Fourth Day.—Eggs (raw), 5; milk, 400 c.c. (13 $\frac{1}{3}$ oz.); sugar, 20 gm. (5 dr.).

Fifth Day

7 A. M.: Egg.

Sugar, 4 gm. (1 dr.).

8 A. M.: Milk, 80 c.c. (2 $\frac{2}{3}$ oz.).

9 A. M.: Egg.

Sugar, 4 gm. (1 dr.).

10 A. M.: Milk, 80 c.c. (2 $\frac{2}{3}$ oz.).

11 A. M.: Egg.

Sugar, 4 gm. (1 dr.).

12 NOON: Milk, 80 c.c. (2 $\frac{2}{3}$ oz.).

1 P. M.: Egg.

Sugar, 4 $\frac{1}{2}$ gm. (1 dr.).

2 P. M.: Milk, 80 c.c. (2 $\frac{2}{3}$ oz.).

3 P. M.: Egg.

Sugar, 4 $\frac{1}{2}$ gm. (1 dr.).

4 P. M.: Milk, 80 c.c. (2 $\frac{2}{3}$ oz.).

5 P. M.: Egg.

Sugar, 4 $\frac{1}{2}$ gm. (1 dr.).

6 P. M.: Milk, 90 c.c. (3 oz.).

7 P. M.: Egg.

Sugar, 4 $\frac{1}{2}$ gm. (1 dr.).

Total Fifth Day.—Eggs (raw), 6; milk, 500 c.c. (16 $\frac{2}{3}$ oz.); sugar, 30 gm. (1 oz.).

Sixth Day

7 A. M.: Egg.

Sugar, 4 gm. (1 dr.).

- 8 A. M.: Milk, 100 c.c. ($3\frac{1}{3}$ oz.).
- 9 A. M.: Egg.
Sugar, $4\frac{1}{2}$ gm. (1 dr.).
Scraped beef, 12 gm. (3 dr.).
- 10 A. M.: Milk, 100 c.c. ($3\frac{1}{3}$ oz.).
- 11 A. M.: Egg.
Sugar, $4\frac{1}{2}$ gm. (1 dr.).
- 12 NOON: Milk, 100 c.c. ($3\frac{1}{3}$ oz.).
- 1 P. M.: Egg.
Sugar, $4\frac{1}{2}$ gm. (1 dr.).
Scraped beef, 12 gm. (3 dr.).
- 2 P. M.: Milk, 100 c.c. ($3\frac{1}{3}$ oz.).
- 3 P. M.: Egg.
Sugar, $4\frac{1}{2}$ gm. (1 dr.).
- 4 P. M.: Milk, 100 c.c. ($3\frac{1}{3}$ oz.).
- 5 P. M.: Egg.
Sugar, 4 gm. (1 dr.).
Scraped beef, 12 gm. (3 dr.).
- 6 P. M.: Milk, 100 c.c. ($3\frac{1}{3}$ oz.).
- 7 P. M.: Egg.
Sugar, $4\frac{1}{2}$ gm. (1 dr.).
- Total Sixth Day.—Eggs (raw), 7; milk, 600 c.c. (20 oz.); sugar, 30 gm. (1 oz.); scraped beef, 36 gm. (9 dr.).

Seventh Day

- 7 A. M.: 1 soft-boiled egg.
- 8 A. M.: Milk, 100 c.c. ($3\frac{1}{3}$ oz.).
- 9 A. M.: Egg.
Sugar, 13 gm. (3 dr.).
- 10 A. M.: Milk, 100 c.c. ($3\frac{1}{3}$ oz.).
Scraped beef, 23 gm. (6 dr.).
Boiled rice, 33 gm. (1 oz.).

- 11 A. M.: 1 soft-boiled egg.
 12 NOON: Milk, 125 c.c. (4 oz.).
 1 P. M.: Egg.
 Sugar, 13 gm. (3 dr.).
 2 P. M.: Milk, 125 c.c. (4 oz.).
 Scraped beef, 23 gm. (6 dr.).
 Boiled rice, 33 gm. (1 oz.).
 3 P. M.: 1 soft-boiled egg.
 4 P. M.: Milk, 125 c.c. (4 oz.).
 5 P. M.: Egg.
 Sugar, 14 gm. ($3\frac{1}{3}$ dr.).
 6 P. M.: Milk, 125 c.c. (4 oz.).
 Scraped beef, 24 gm. (6 dr.).
 Boiled rice, 34 gm. (1 oz.).
 7 P. M.: 1 soft-boiled egg.

Total Seventh Day.—Eggs (raw), 4; soft boiled, 4; milk, 700 c.c. ($23\frac{1}{3}$ oz.); sugar, 40 gm. ($1\frac{1}{3}$ oz.); scraped beef, 70 gm. ($2\frac{1}{3}$ oz.); boiled rice, 100 gm. ($3\frac{1}{3}$ oz.), served with beef-juice.

Eighth Day

Diet changes on the eighth day, requiring only 4 raw eggs, which may be divided into three feedings. The other 4 eggs are to be soft boiled and given as directed by diet.

- 7 A. M.: 1 soft-boiled egg.
 8 A. M.: Milk, 135 c.c. ($4\frac{1}{2}$ oz.).
 9 A. M.: Egg.
 Sugar, 13 gm. (3 dr.).
 10 A. M.: Milk, 133 c.c. ($4\frac{1}{2}$ oz.).
 Scraped beef, 23 gm. (6 dr.).
 Boiled rice, 33 gm. (1 oz.).

- 11 A. M.: 1 soft-boiled egg.
Zwieback, 10 gm. ($2\frac{1}{2}$ dr.).
- 12 NOON: Milk, 133 c.c. ($4\frac{1}{2}$ oz.).
- 1 P. M.: Egg.
Sugar, 13 gm. (3 dr.).
- 2 P. M.: Milk, 133 c.c. ($4\frac{1}{2}$ oz.).
Scraped beef, 23 gm. (6 dr.).
Boiled rice, 33 gm. (1 oz.).
- 3 P. M.: 1 soft-boiled egg.
- 4 P. M.: Milk, 133 c.c. ($4\frac{1}{2}$ oz.).
- 5 P. M.: Egg.
Sugar, 14 gm. ($3\frac{1}{2}$ dr.).
Zwieback, 10 gm. ($2\frac{1}{2}$ dr.).
- 6 P. M.: Milk, 133 c.c. ($4\frac{1}{2}$ oz.).
Scraped beef, 24 gm. (6 dr.).
Boiled rice, 34 gm. (1 oz.).
- 7 P. M.: 1 soft-boiled egg.

Total Eighth Day.—Eggs (raw), 4; soft boiled, 4; milk, 800 c.c. ($26\frac{2}{3}$ oz.); scraped beef, 70 gm. ($2\frac{1}{2}$ oz.); boiled rice, 100 gm. ($3\frac{1}{3}$ oz.); zwieback, 20 gm. (5 dr.); sugar, 40 gm. ($1\frac{1}{3}$ oz.).

Ninth Day

- 7 A. M.: 1 soft-boiled egg.
- 8 A. M.: Milk, 150 c.c. (5 oz.).
- 9 A. M.: Egg.
Sugar, 13 gm. (3 dr.).
- 10 A. M.: Milk, 150 c.c. (5 oz.).
Scraped beef, 23 gm. (6 dr.).
Boiled rice, 66 gm. (2 oz.).
- 11 A. M.: 1 soft-boiled egg.
Zwieback, 20 gm. (5 dr.).

12 NOON: Milk, 150 c.c. (5 oz.).

1 P. M.: Egg.

Sugar, 13 gm. (3 dr.).

2 P. M.: Milk, 150 c.c. (5 oz.).

Scraped beef, 23 gm. (6 dr.).

Boiled rice, 67 gm. (2 oz.).

3 P. M.: 1 soft-boiled egg.

Zwieback, 20 gm. (5 dr.).

4 P. M.: Milk, 150 c.c. (5 oz.).

5 P. M.: Egg.

Sugar, 14 gm. ($3\frac{1}{2}$ dr.).

6 P. M.: Milk, 150 c.c. (5 oz.).

Scraped beef, 24 gm. (6 dr.).

Boiled rice, 67 gm. (2 oz.).

7 P. M.: 1 soft-boiled egg.

Total Ninth Day.—Eggs (raw), 4; cooked, 4; milk, 900 c.c. (30 oz.); sugar, 40 gm. ($1\frac{1}{3}$ oz.); scraped beef, 70 gm. ($2\frac{1}{3}$ oz.); rice, 200 gm. ($6\frac{2}{3}$ oz.); zwieback, 40 gm. ($1\frac{1}{3}$ oz.), or toast, 20 gm. (5 dr.).

Tenth Day

7 A. M.: 1 soft-boiled egg.

8 A. M.: Milk, 166 c.c. ($5\frac{1}{2}$ oz.).

9 A. M.: Egg.

Sugar, 13 gm. (3 dr.).

10 A. M.: Milk, 168 c.c. ($5\frac{1}{2}$ oz.).

Scraped beef, 23 gm. (6 dr.).

Boiled rice, 66 gm. (2 oz.).

11 A. M.: 1 soft-boiled egg.

Zwieback, 20 gm. (5 dr.).

Butter, 4 gm. (1 dr.).

- 12 NOON: Cooked chopped chicken, 25 gm. (6 dr.).
Milk, 166 c.c. ($5\frac{1}{2}$ oz.).
- 1 P. M.: Egg.
Sugar, 13 gm. (3 dr.).
- 2 P. M.: Milk, 166 c.c. ($5\frac{1}{2}$ oz.).
Scraped beef, 23 gm. (6 dr.).
Boiled rice, 66 gm. (2 oz.).
Butter, 4 gm. (1 dr.).
- 3 P. M.: 1 soft-boiled egg.
Zwieback, 20 gm. (5 dr.).
Butter, 4 gm. (1 dr.).
- 4 P. M.: Cooked chopped chicken, 25 gm. (6 dr.).
- 5 P. M.: Egg.
Sugar, 14 gm. ($3\frac{1}{2}$ dr.).
- 6 P. M.: Milk, 166 c.c. ($5\frac{1}{2}$ oz.).
Scraped beef, 24 gm. (6 dr.).
Boiled rice, 67 gm. (2 oz.).
Butter, 4 gm. (1 dr.).
- 7 P. M.: 1 soft-boiled egg.

Total Tenth Day.—Eggs (raw), 4; cooked, 4; milk, 1000 c.c. ($33\frac{1}{3}$ oz.); sugar, 40 gm. ($1\frac{1}{3}$ oz.); scraped beef, 70 gm. ($2\frac{1}{3}$ oz.); boiled rice, 200 gm. ($6\frac{2}{3}$ oz.); zwieback, 40 gm. ($1\frac{1}{3}$ oz.), or toast, 20 gm. (5 dr.); chicken, 50 gm. ($1\frac{2}{3}$ oz.); butter, 20 gm. (5 dr.).

Eleventh Day

- 7 A. M.: 1 soft-boiled egg.
Milk, 250 c.c. ($8\frac{1}{3}$ oz.).
Zwieback, 10 gm. ($2\frac{1}{2}$ dr.).
Butter, 4 gm. (1 dr.).

8 A. M.: Egg.

Sugar, 13 gm. (3 dr.).

Scraped beef, 20 gm. (5 dr.).

Boiled rice, 75 gm. ($2\frac{1}{2}$ oz.).

Zwieback, 10 gm. ($2\frac{1}{2}$ dr.).

Butter, 6 gm. ($1\frac{1}{2}$ dr.).

11 A. M.: 1 soft-boiled egg.

Milk, 250 c.c. ($8\frac{1}{3}$ oz.).

Butter, 6 gm. ($1\frac{1}{2}$ dr.).

Zwieback, 10 gm. ($2\frac{1}{2}$ dr.).

1 P. M.: Egg.

Sugar, 13 gm. (3 dr.).

Cooked chopped chicken, 25 gm. (6 dr.).

Boiled rice, 75 gm. ($2\frac{1}{2}$ oz.).

3 P. M.: 1 soft-boiled egg.

Milk, 250 c.c. ($8\frac{1}{3}$ oz.).

Scraped beef, 20 gm. (5 dr.).

Boiled rice, 75 gm. ($2\frac{1}{2}$ oz.).

Zwieback, 10 gm. ($2\frac{1}{2}$ dr.).

Butter, 6 gm. ($1\frac{1}{2}$ dr.).

5 P. M.: Egg.

Sugar, 14 gm. ($3\frac{1}{2}$ dr.).

Cooked chopped chicken, 25 gm. (6 dr.).

Boiled rice, 75 gm. ($2\frac{1}{2}$ oz.).

Butter, 6 gm. ($1\frac{1}{2}$ dr.).

7 P. M.: 1 soft-boiled egg.

Milk, 250 c.c. ($8\frac{1}{3}$ oz.).

Zwieback, 10 gm. ($2\frac{1}{2}$ dr.).

Butter, 6 gm. ($1\frac{1}{2}$ dr.).

Scraped beef, 30 gm. (1 oz.).

Total Eleventh Day.—Eggs (raw), 4; cooked, 4; milk, 1000 c.c. ($33\frac{1}{3}$ oz.); butter, 40 gm. ($1\frac{1}{3}$ oz.); sugar,

40 gm. ($1\frac{1}{3}$ oz.); scraped beef, 70 gm. ($2\frac{1}{3}$ oz.); boiled rice, 300 gm. (30 oz.); zwieback, 60 gm. (2 oz.); chicken, 50 gm. ($1\frac{2}{3}$ oz.).

Twelfth Day.

- 7 A. M.: 1 soft-boiled egg.
Milk, 250 c.c. ($8\frac{1}{3}$ oz.).
Zwieback, 10 gm. ($2\frac{1}{2}$ dr.).
Butter, 4 gm. (1 dr.).
- 9 A. M.: Egg.
Sugar, 13 gm. (3 dr.).
Scraped beef, 35 gm. (1 oz.).
Boiled rice, 75 gm. ($2\frac{1}{2}$ oz.).
Zwieback, 10 gm. ($2\frac{1}{2}$ dr.).
Butter, 6 gm. ($1\frac{1}{2}$ dr.).
- 11 A. M.: 1 soft-boiled egg.
Milk, 250 c.c. ($8\frac{1}{3}$ oz.).
Zwieback, 20 gm. (5 dr.).
Butter, 6 gm. ($1\frac{1}{2}$ dr.).
- 1 P. M.: Egg.
Sugar, 13 gm. (3 dr.).
Cooked chopped chicken, 25 gm. (6 dr.).
Boiled rice, 75 gm. ($2\frac{1}{2}$ oz.).
Zwieback, 10 gm. ($2\frac{1}{2}$ dr.).
Butter, 6 gm. ($1\frac{1}{2}$ dr.).
- 3 P. M.: 1 soft-boiled egg.
Milk, 250 c.c. ($8\frac{1}{3}$ oz.).
Scraped beef, 35 gm. (1 oz.).
Boiled rice, 50 gm. ($1\frac{2}{3}$ oz.).
Zwieback, 10 gm. ($2\frac{1}{2}$ dr.).
Butter, 6 gm. ($1\frac{1}{2}$ dr.).

5 P. M.: Egg.

Sugar, 14 gm. ($3\frac{1}{2}$ dr.).

Cooked chopped chicken, 25 gm. (6 dr.).

Boiled rice, 75 gm. ($2\frac{1}{2}$ oz.).

Zwieback, 10 gm. ($2\frac{1}{2}$ dr.).

Butter, 6 gm. ($1\frac{1}{2}$ dr.).

7 P. M.: 1 soft-boiled egg.

Milk, 250 c.c. ($8\frac{1}{3}$ oz.).

Zwieback, 10 gm. ($2\frac{1}{2}$ dr.).

Butter, 6 gm. ($1\frac{1}{2}$ dr.).

Total Twelfth Day.—Eggs (raw), 4; cooked, 4; milk, 1000 c.c. ($33\frac{1}{3}$ oz.); sugar, 40 gm. ($1\frac{1}{3}$ oz.); scraped beef, 70 gm. ($2\frac{1}{3}$ oz.); boiled rice, 300 gm. (10 oz.); zwieback, 80 gm. ($2\frac{2}{3}$ oz.); chicken, 50 gm. ($1\frac{2}{3}$ oz.); butter, 40 gm. ($1\frac{1}{2}$ oz.).

Thirteenth Day

7 A. M.: 1 soft-boiled egg.

Milk, 142 c.c. ($4\frac{2}{3}$ oz.).

Zwieback, 10 gm. ($2\frac{1}{2}$ dr.).

Butter, 4 gm. (1 dr.).

9 A. M.: Egg.

Sugar, 13 gm. (3 dr.).

Milk, 142 c.c. ($4\frac{2}{3}$ oz.).

Scraped beef, 20 gm. (5 dr.).

Boiled rice, 75 gm. ($2\frac{1}{2}$ oz.).

Zwieback, 20 gm. (5 dr.).

Butter, 6 gm. ($1\frac{1}{2}$ dr.).

11 A. M.: 1 soft-boiled egg.

Milk, 144 c.c. (5 oz.).

Zwieback, 10 gm. ($2\frac{1}{2}$ dr.).

Butter, 6 gm. ($1\frac{1}{2}$ dr.).

1 P. M.: Egg.

Sugar, 13 gm. (3 dr.).

Milk, 142 c.c. ($4\frac{2}{3}$ oz.).

Cooked chopped chicken, 25 gm. (6 dr.).

Boiled rice, 75 gm. ($2\frac{1}{2}$ oz.).

Zwieback, 10 gm. ($2\frac{1}{2}$ dr.).

Butter, 6 gm. ($1\frac{1}{2}$ dr.).

3 P. M.: 1 soft-boiled egg.

Milk, 144 c.c. (5 oz.).

Scraped beef, 20 gm. (5 dr.).

Boiled rice, 75 gm. ($2\frac{1}{2}$ oz.).

Zwieback, 10 gm. ($2\frac{1}{2}$ dr.).

Butter, 6 gm. ($1\frac{1}{2}$ dr.).

5 P. M.: Egg.

Sugar, 14 gm. ($3\frac{1}{2}$ dr.).

Milk, 142 c.c. (5 oz.).

Cooked chopped chicken, 25 gm. (6 dr.).

Boiled rice, 75 gm. ($2\frac{1}{2}$ oz.).

Zwieback, 10 gm. ($2\frac{1}{2}$ dr.).

Butter, 6 gm. ($1\frac{1}{2}$ dr.).

7 P. M.: 1 soft-boiled egg.

Milk, 144 c.c. (5 oz.).

Zwieback, 10 gm. ($2\frac{1}{2}$ dr.).

Butter, 6 gm. ($1\frac{1}{2}$ dr.).

Total Thirteenth Day.—Eggs (raw), 4; cooked, 4; milk, 1000 c.c. ($33\frac{1}{3}$ oz.); sugar, 40 gm. ($1\frac{1}{3}$ oz.); scraped beef, 70 gm. ($2\frac{1}{3}$ oz.); boiled rice, 300 gm. (10 oz.); zwieback, 80 gm. ($2\frac{2}{3}$ oz.); chicken, 50 gm. ($1\frac{2}{3}$ oz.); butter, 40 gm. ($1\frac{1}{3}$ oz.).

Fourteenth Day

- 7 A. M.: 1 soft-boiled egg.
Minced chop.
Buttered toast.
Milk, 142 c.c. ($4\frac{2}{3}$ oz.).
- 9 A. M.: Boiled rice.
Buttered zwieback.
Custard.
Milk, 142 c.c. ($4\frac{2}{3}$ oz.).
- 11 A. M.: 1 soft-boiled egg.
Buttered zwieback.
Junket.
Milk, 144 c.c. (5 oz.).
- 1 P. M.: Minced chicken.
Boiled rice.
Buttered zwieback.
Custard.
Milk, 142 c.c. ($4\frac{2}{3}$ oz.).
- 3 P. M.: 1 soft-boiled egg.
Cooked scraped beef.
Boiled rice.
Buttered tost.
Milk, 144 c.c. (5 oz.).
- 5 P. M.: Minced chicken.
Boiled rice.
Buttered zwieback.
Custard.
Milk, 142 c.c. ($4\frac{2}{3}$ oz.).
- 7 P. M.: 1 soft-boiled egg.
Buttered toast.
Milk, 144 c.c. (5 oz.).
- Total Fourteenth Day.—Eggs (raw), 4; cooked, 4;

milk, 1000 c.c. ($33\frac{1}{3}$ oz.); sugar, 40 gm. ($1\frac{1}{3}$ oz.); scraped beef, 70 gm. ($2\frac{1}{3}$ oz.); boiled rice, 300 gm. (10 oz.); zwieback, 100 gm. ($3\frac{1}{3}$ oz.); butter, 40 gm. ($1\frac{1}{3}$ oz.) chicken, 50 gm. ($1\frac{2}{3}$ oz.). (Recapitulation of Lenhartz Diet, p. 49.)

CONVALESCENT GASTRIC DIETS, NO. 1¹

Breakfast: Boiled milk with cocoa or coffee.

Any cereal, strained, with cream.

Soft toast, buttered.

11 A. M.: Glass of milk.

Dinner: Purée of potatoes, peas, or beans with toast soaked in it, or boiled rice, with milk or cream.

Junket or custard, or jelly or tapioca pudding.

4 P. M.: Glass of milk with beaten egg in it.

Supper: 2 soft-boiled eggs.

Soft toast.

Glass of milk.

9.30 P. M.: Glass of milk, with beaten egg in it.

CONVALESCENT GASTRIC DIETS, NO. 2²

Breakfast: Boiled milk, with a little coffee or cocoa.

Any cereal, strained, with cream.

Dry toast, buttered.

1 egg.

11 A. M.: Glass of malted milk.

Lunch: 2 soft-boiled or scrambled eggs, or fresh boiled fish in place of eggs.

A little broiled bacon (eating the fat and not the lean.)

Toast and milk.

¹ T. C. Janeway.

² Ibid.

GASTRIC DIETS

RECAPITULATION OF LENHARTZ DIET

Day.	Calories, Eggs.	Milk.	Sugar.	Scraped beef.	Boiled rice.	Zwieback.	Butter.	Chicken.
1....	280 Raw 2	100 c.c. (3½ oz.)						
2....	470 Raw 3	200 c.c. (6½ oz.)						
3....	637 Raw 4	300 c.c. (10 oz.)	20 gm. (5 dr.)					
4....	777 Raw 5	400 c.c. (13½ oz.)	20 gm. (5 dr.)					
5....	966 Raw 6	500 c.c. (16½ oz.)	30 gm. (1 oz.)					
6....	1135 Raw 7	600 c.c. (20 oz.)	30 gm. (1 oz.)	36 gm. (9 dr.)				
7....	1580 Raw 4, soft 4	700 c.c. (23½ oz.)	40 gm. (1½ oz.)	70 gm. (2½ oz.)	100 gm. (3½ oz.)			
8....	1720 Raw 4, soft 4	800 c.c. (26½ oz.)	40 gm. (1½ oz.)	70 gm. (2½ oz.)	100 gm. (3½ oz.)	20 gm. (¾ oz.)		
9....	2138 Raw 4, soft 4	900 c.c. (30 oz.)	40 gm. (1½ oz.)	70 gm. (2½ oz.)	200 gm. (6½ oz.)	40 gm. (1½ oz.) of toast, 20 gm.		
10....	2478 Raw 4, soft 4	1000 c.c. (33½ oz.)	40 gm. (1½ oz.)	70 gm. (2½ oz.)	200 gm. (6½ oz.)	40 gm. (1½ oz.) of toast, 20 gm.	20 gm. (¾ oz.)	50 gm. (1½ oz.)
11....	2941 Raw 4, soft 4	1000 c.c. (33½ oz.)	40 gm. (1½ oz.)	70 gm. (2½ oz.)	300 gm. (10 oz.)	60 gm. (2 oz.) (¾ oz.)	40 gm. (1½ oz.)	50 gm. (1½ oz.)
12....	2941 Raw 4, soft 4	1000 c.c. (33½ oz.)	40 gm. (1½ oz.)	70 gm. (2½ oz.)	300 gm. (10 oz.)	80 gm. (2½ oz.)	40 gm. (1½ oz.)	50 gm. (1½ oz.)
13....	3007 Raw 4, soft 4	1000 c.c. (33½ oz.)	40 gm. (1½ oz.)	70 gm. (2½ oz.)	300 gm. (10 oz.)	80 gm. (2½ oz.)	40 gm. (1½ oz.)	50 gm. (1½ oz.)
14....	3007	Same as the thirteenth day.						

Dinner: Purée of potatoes, rice, barley, peas, asparagus, celery, or beans.

Buttered toast.

Rare beefsteak broiled, or lamb chop broiled, roast beef or lamb.

Roast or broiled chicken or minced chicken, with well-cooked rice or a well-baked potato with butter.

Junket or custard, jelly or tapioca pudding.

10.30 P. M.: Glass of milk with lime-water.

COMMENTS ON GASTRIC DIETS

Stomach.—The reputation of the Lenhartz diet in the treatment of peptic ulcer is too well established to need comment; one or two things only might be said in regard to its use. In the first place, in this country there is a deep-rooted and reasonable prejudice against the use of any hard substance in the diet of gastric ulcer, either acute or chronic, and wherever this occurs in the Lenhartz diet it would seem wise to substitute some soft substance—*e. g.*, soaked toast or bread in place of dry zwieback. Then, too, the rapidity with which this routine can be carried out varies with different persons, and it is often necessary to go more slowly than the regular schedule.

Before the introduction of this diet, it had been thought necessary to starve, or practically starve, patients with acute peptic ulcer for a week or two,

except for uncertain attempts at rectal feeding. Lenhartz went to the other extreme, and taught that one could begin at once with advantage to feed these cases, even in the face of hemorrhage, feeling that patients would do better if given food, as starvation makes it more difficult for them to further the process of repair. His idea was to furnish a highly albuminous, bland food in gradually increasing amounts, and during the feeding hours always keeping something in the stomach for the hydrochloric acid to combine with, which would otherwise keep the ulcerated surface irritated, *i. e.*, if free. A period of twelve hours' rest is given the stomach. A convenient method for using the Lenhartz diet is to have each day's diet written on a card which is hung with the patient's chart. The cards are changed every day as the ration is increased.

To many internists, however, it also seems a wise measure to give the stomach absolute rest for from forty-eight to seventy-two hours or even longer (up to one week), particularly in acute hemorrhagic cases, before beginning any mode of gastric feeding, during which time, or, at all events, for the first two or three days, normal salt solution by rectum will furnish all the fluids necessary. If it seems best to rest the stomach longer than two days, it is usually best to furnish some food by rectal alimentation. As usually done, with a thick mixture of eggs, milk, flour, etc., this form of feeding is absolutely useless, and, as proved by numerous experiments

of the author and others, almost nothing is absorbed. If, however, milk is given by rectum, which has been first sterilized and then fully peptonized for two hours or more or until it becomes watery, thin, and of a greenish-yellow color, and then scalded to kill the bacteria, which, of course, increase during the two hours or more at incubation temperature, a certain amount of protein and calories can be furnished to the organism. To be sure, this is entirely insufficient, but still enough to help maintain strength. This milk can be further reinforced by a glucose or maltose addition, although care must be taken that its presence does not increase the rectal peristalsis and result in failure to retain the enemata. The author's usual dietetic treatment of these cases is to starve for forty-eight hours, giving saline by rectum, 8 to 16 oz. every four to six hours, with the foot of the bed elevated. Then begin with rectal feeding of fully peptonized milk, 8 to 16 oz. every four to six hours, depending on the ability of the patient to hold it. Given through a rectal tube inserted fairly high, with the foot of the bed well raised and kept so for at least an hour after feedings, there is seldom any difficulty in the patient's retaining it. To each rectal feeding may be added from 1 to $1\frac{1}{2}$ oz. of glucose or maltose, first thoroughly dissolved in a little hot water. This rectal feeding is kept up for several days, or until the mouth feedings, which can begin from the third to the sixth day, are more nearly

adequate to supply the needed nourishment. After the seventh day of the Lenhartz diet, it is often best to increase it every other day instead of every day, or in some cases to work toward the use of the Convalescent Gastric Diets Nos. 1 and 2. The lower colon should be washed out once daily with saline in order to keep the mucous membrane in a clean condition. Under certain circumstances rectal feeding is impracticable; in these cases, particularly if mild, or the patient robust, it may be dispensed with.

The dietetic treatment of ambulatory cases of chronic gastric or duodenal ulceration has a very distinct place in therapeutics, for many of these cases are apparently well, except for the characteristic epigastric pain which comes on at varying times after meals, and refuse the suggestion of taking an ulcer "cure," as they will not give the necessary time. These cases are often helped surprisingly if they are put on a diet for two full weeks of milk, soft-boiled eggs (one to two minutes), and cream, given somewhat as follows:

At 8 A. M., 1 P. M., and 7 P. M.: 2 glasses of milk (and cream); 2 soft-boiled eggs.

At 11 A. M., 4 P. M., and 10 P. M.: $1\frac{1}{2}$ glasses of milk and cream.

This gives: Milk, 2040 c.c. (68 oz.); cream, 480 c.c. (16 oz.); 6 eggs.

Protein, 102 gm. ($3\frac{1}{2}$ oz.); fat, 200 gm. ($6\frac{3}{4}$ oz.); carbohydrate, 123 gm. (4 oz.); calories, 2200.

At the end of ten days, if this has been well taken, and there has been no pain for several days, the patient may be allowed well cooked farina twice a day (1 or 2 tablespoonfuls).

At the end of the two weeks other soft foods are gradually added, such as any thin cereal, soft toast, custard, finely cut white meat of boiled chicken, boiled cod or halibut, then baked potatoes, chopped steak, and anything which is soft and without acid or excess of salt.

Many cases will go for months or years after such a course of diet without a return of symptoms. These cases should never eat hard or irritating foods, such as berries with seeds, figs, hard toast, raw vegetables, etc.; acids and condiments as well should be avoided.

The *convalescent gastric diets* will be found of use after acute gastritis or following a Lenhartz ulcer cure, for it is wise to insist that for some weeks the patient shall put into the stomach only the softest foods possible.

The dietary treatment of duodenal ulcer is quite the same as that for gastric, and it is not always possible to differentiate between them clinically.¹

¹In connection with the dietary treatment of these cases of chronic ulcer of the stomach or duodenum, the use of subnitrate of bismuth in dram doses, with 2 ounces of water on early morning fasting stomach, is almost always of the greatest assistance in stopping the pain. This is often advantageously preceded by washing the stomach clean, then with silver nitrate solution, beginning 1:6000 and increasing daily until 1:2000 is used, wash again until all trace of silver disappears. This can be continued daily for seven to ten days, then every other day, twice or once a week, until all symptoms are gone.

GASTRIC HYPERACIDITY

THE dietetic treatment of a hyperacid gastric condition, of course, depends largely upon the underlying cause of the hyperacidity; thus, if it is due to ulcer of the stomach or duodenum, the proper diet for these conditions is indicated. There are cases, however, in which the hyperacidity is a symptom of indefinite and apparently of mild origin in which a modified diet is of use.

The principle upon which this diet is constructed is based upon the fact that food which is chemically, thermally, or mechanically irritating, or which stays a long time in the stomach, is almost sure to increase the hyperacidity. Hence, these things should be omitted from the diet and only bland foods used. The other underlying principle of the diet is that it should be of a high protein value, in order to offer a large amount of this element to combine with the free hydrochloric acid which is in excess in this condition.

The protein should preferably be of the least stimulating variety, hence meat is allowed only in small amount. The best kind of animal protein being that contained in eggs, milk, and boiled fish.

Diet Useful in Gastric Hyperacidity

May take:

Raw oysters.

Soups: Cream or purée (except tomato).

Fish: Fresh cod, halibut, bass, boiled with cream sauce or broiled.

Meat: Beef well done (without gristle, fat, or gravy), chicken, turkey, guinea-hen, lamb (without fat), once daily.

Vegetables: All soft-boiled green vegetables, except cabbage, cauliflower, brussel sprouts, or turnips. Baked white potato may be used moderately. Rice or macaroni.

Cheese: Any mild variety, but better without this at first.

Desserts: Cream and egg desserts of all sorts, *e. g.*, blanc mange, Bavarian cream, floating island, cup custard, junket, soft rice or bread puddings. Gelatin desserts made with little flavoring. Very little sugar used in all desserts. Fruit, none, except when constipation is a marked feature, then stewed soft fruits may be taken in good amount, but must be cooked with very little sugar and are best taken with or after a meal, never before. Cream may usually be freely used.

Bread: Toast, stale bread, roll. (Fresh butter or salt butter freshened by working it over in fresh water.)

Drinks: Weak tea, cocoa, water, and milk.

Cereals: Fine-grained varieties, well cooked.

Eggs: In all forms except fried.

Foods to be avoided:

All sour, spiced, or peppery foods, condiments, salt foods, chow-chow, etc. Sweets, fried foods.

Very hot or very cold food or drinks.

Rough hard substances, such as seeds, skins of fruit or vegetables, corn, uncooked vegetables.

Coffee, wines, liquors, beer.

Hot breads, pies, cakes, syrups, etc.



DIABETIC DIETS

STANDARD STRICT DIET¹

Breakfast: 2 eggs.

Ham, 90 gm. (3 oz.).

Coffee with 45 gm. ($1\frac{1}{2}$ oz.) cream.

Butter, 15 gm. ($\frac{1}{2}$ oz.) on the biscuit during the test period; cooked with the eggs if no biscuit or bread is taken.

Luncheon: Meat (steak or chops), 120 gm. ($\frac{1}{4}$ pound).

Green vegetables from list, 2 tablespoonfuls.

White wine, 2 claret glasses (6 oz.), or whisky or brandy, 2 tablespoonfuls (1 oz.).

Butter, 15 gm. ($\frac{1}{2}$ oz.), with the green vegetable, if no biscuit or bread is taken.

Afternoon tea with 15 gm. ($\frac{1}{2}$ oz.) of cream.

Dinner: Any clear soup.

Fish, 90 gm. (3 oz.).

Meat (beef, mutton, turkey, or chicken), 120 gm. ($\frac{1}{4}$ pound).

Green vegetables from list, 2 tablespoonfuls.

Salad with 15 gm. ($\frac{1}{2}$ oz.) of oil in the dressing.

¹ T. C. Janeway in Musser and Kelly, Practical Therapeutics.

Cream cheese, 30 gm. (1 oz.).

White wine, 2 claret glasses (6 oz.), or
whisky or brandy, 2 tablespoonfuls
(1 oz.).

Demi-tasse of coffee.

Butter, 30 gm. (1 oz.) on the fish, meat, and
green vegetables if no bread or biscuit
is taken.

Bedtime: Bouillon with 1 raw egg.

Protein, 112 gm. ($3\frac{2}{3}$ oz.); nitrogen, 18 gm. ($4\frac{1}{2}$ dr.);
fats, 160 gm. ($5\frac{1}{2}$ oz.); calories, 2200; omitting ham,
protein, 94 gm. (3 oz.); nitrogen, 15 gm. ($3\frac{3}{4}$ dr.).

For convenience in determining the carbohydrate
tolerance the following biscuits may be used, as the per-
centage of carbohydrate is practically constant.

Huntley & Palmer breakfast biscuit contains 5 gm.
carbohydrate. Uneda biscuit contains 4.6 gm. carbo-
hydrate (Mosenthal).

STANDARD DIET WITH RESTRICTED PROTEIN¹

Breakfast: 2 eggs.

Bacon, 15 gm. ($\frac{1}{2}$ oz.).

Coffee, with 45 gm. ($1\frac{1}{2}$ oz.) of cream.

Butter, 20 gm. ($\frac{2}{3}$ oz.).

Luncheon: 1 egg.

Bacon, 15 gm. ($\frac{1}{2}$ oz.).

Meat (lamb-chops, ham, or beefsteak),
60 gm. (2 oz.).

Salad, with 15 gm. ($\frac{1}{2}$ oz.) of oil in the
dressing.

¹ T. C. Janeway in Musser and Kelly, Practical Therapeutics.

White wine, 2 claret glasses, 180 c.c. (6 oz.), or whisky or brandy, 2 tablefuls, 30 c.c. (1 oz.).

Butter, 40 gm. ($1\frac{1}{3}$ oz.).

Afternoon tea with 15 gm. ($\frac{1}{2}$ oz.) of cream.

Dinner: Any clear soup.

Meat (roast pork, beef, mutton, turkey, or lamb chops), 90 gm. (3 oz.).

Vegetables from list, 2 tablespoonfuls.

Salad with 15 gm. ($\frac{1}{2}$ oz.) of oil in the dressing.

Cream cheese, 30 gm. (1 oz.).

White wine, 2 claret glasses, 180 c.c. (6 oz.), or whisky or brandy, 2 table-spoonfuls, 30 c.c. (1 oz.).

Demi-tasse of coffee.

Butter, 30 gm. (1 oz.).

Bedtime: Bouillon with 1 raw egg.

Protein, 62 gm. (2 oz.); nitrogen, 10 gm. (2 dr.); fat, 180 gm. (6 oz.); total calories equal 2500. Omitting 30 gm. of butter and $\frac{1}{2}$ oz. of bacon, calories equal 2250.

GREEN DAYS¹

Breakfast: 1 egg, boiled or poached.

Cupful of black coffee.

Dinner: Spinach, with hard-boiled egg.

Bacon, 15 gm. ($\frac{1}{2}$ oz.).

Salad with 15 gm. ($\frac{1}{2}$ oz.) of oil.

White wine, $\frac{1}{4}$ liter (4 oz.), or whisky or brandy 30 c.c. (1 oz.).

4.30 P. M.: Cup of beef-tea or chicken broth.

¹ T. C. Janeway in Musser and Kelly, Practical Therapeutics.

Supper: 1 egg, scrambled with tomato and a little butter.

Bacon, 15 gm. ($\frac{1}{2}$ oz.).

Cabbage, cauliflower, sauerkraut, string-beans, or asparagus.

White wine, $\frac{1}{8}$ liter (4 oz.), or whisky or brandy, 30 c.c. (1 oz.).

Sodium bicarbonate, 15 to 30 gm. ($\frac{1}{2}$ -1 oz.) in the twenty-four hours. Protein, 32 gm. (1 oz.); nitrogen, 5 gm. (1 dr.); carbohydrate, about 5 gm. (1 dr.); fat, 67 gm. (2 oz.); calories, 575.

GENERAL DIABETIC DIET-LIST¹

May take freely:

Soups: All meat soups and broths. May add vegetables allowed; egg or cheese.

Meats: All fresh, smoked, and cured meats (except liver), poultry and game, without sauces or gravies containing flour, pâté de foies gras.

Fish: All kinds, except oysters, clams, and scallops, cooked without bread-crumbs or meal; all dried, salted, smoked or pickled fish.

Eggs: Prepared in any way without flour.

Fats: Butter, lard, suet, olive oil, or other fats.

Cheeses: All kinds, especially cream, Swiss, English, and pineapple cheese.

Vegetables and Salads: Asparagus, beet greens, Brussels sprouts, cabbage, cauliflower, celery, chicory, cresses, cucumbers, egg-plant, endive, kohlrabi, leeks, lettuce, okra, pumpkin, radishes, rhubarb,

¹ T. C. Janeway in Musser and Kelly, Practical Therapeutics.

salsify, sauerkraut, spinach, string-beans, tomatoes, vegetable marrow.

Pickles made from the above vegetables, unsweetened; ripe olives.

Fungi: Mushrooms and truffles.

Cream: Not over 90 c.c. (3 oz.) a day.

Condiments: Salt, pepper, cayenne, paprika, curry, cinnamon, cloves, English mustard, nutmeg, caraway, capers, vinegar, and the piquant sauces in small quantities unless specially forbidden.

Desserts: Jellies made from gelatin; custards and ice-cream made with eggs and cream; all sweetened with saccharin and flavored with vanilla; coffee or brandy.

Nuts: Butternuts.

Beverages: Tea or coffee, sweetened with saccharin and with the portion of cream allowed. Whisky, brandy, rum, and other distilled liquors, up to 3 oz. a day. Light Rhine or Moselle wine, claret, or Burgundy, up to 16 oz. a day (1 pint). Mineral waters of all kinds. Lemonade in small quantity, sweetened with saccharin.

Articles Prohibited, Except as Prescribed in the Accessory Diet:

Sugars and sweets of every kind.

Pastry, puddings, preserves, cake, and ice-cream.

Bread and biscuits of all kinds, toast, crackers, and griddle cakes.

Cereals, such as rice, oatmeal, sago, hominy, tapioca, and barley.

Macaroni, potatoes, carrots, parsnips, beans, peas, beets, green corn, and turnips.

Fruit of all kinds, fresh or dried.

Soups, sauces, or gravies thickened with flour or meal, or made with milk.

Beer, ale, porter, all sweet wines, sherry or port wine, sparkling wines, cider, and liquors.

Milk, chocolate, or cocoa.

All sweet drinks and soda-water.

OATMEAL DAYS ¹

Porridge made from oatmeal, 250 gm. (8 oz.), with butter, 250 gm. (8 oz.); salt and pepper to taste.

Black coffee, light white wine, $\frac{1}{4}$ liter (8 oz.), or cognac, 60 c.c. (2 oz.).

The whites of 6 eggs may be added to the porridge if desired.

	Nitrogen.	Carbohydrate.	
	Gm.	Gm.	Calories.
Oatmeal.....	6.2	170	1025
Butter.....	0.4	...	1975
	6.6 = 42 gm. protein.		3000
Alcohol (40 gm.).....			210
6 egg whites.....	3.6		90
	10.2 = 63 gm. protein.		3300

The entire diet consists of: Protein, 63 gm. (2 oz.); nitrogen, 16.8 gm. ($\frac{1}{2}$ oz.); carbohydrate, 170 gm. ($5\frac{2}{3}$ oz.); fat, 212 gm. (7 oz.); calories, 3300.

COMMENTS ON DIABETIC DIET

Such an enormous amount has already been written on the dietetic treatment of diabetes mellitus that one hesitates about going into the details and indications.

¹ T. C. Janeway in Musser and Kelly, Practical Therapeutics.

The principles underlying the subject may, however, be stated here with propriety, and general directions given for the use of the diabetic diets and the table of carbohydrate equivalents, which will be found to be a great help in arranging a varied and, therefore, more palatable diet for patients with this disease.

Whatever may be the underlying organic lesion in a particular case, the result is the same; namely, that the power of properly metabolizing carbohydrates is impaired to a greater or less degree.

The object sought in dietetic treatment is to find the limit of the individual's carbohydrate tolerance, and then, by a slow system of gradually increasing the amount of starch or sugar, re-educate the system in its lost power to as high a degree as possible, always remembering that these cases can never go back to a free or unlimited use of carbohydrates.

The exact form which the dietetic treatment of diabetes should take must depend on the severity of the case, and, for the sake of clearness on this point, we may arbitrarily divide the cases into:

- (1) Mild, without acidosis.
- (2) Moderately severe, without acidosis or with moderate acidosis.
- (3) Severe, with marked acidosis.

In (1) and (2) the first step is to find the individual's capacity for metabolizing carbohydrates, which is done as follows:

A gradual reduction in starch and sugar in the diet is made each day, using less, until, at the end of five to seven days, the patient is taking none of either. If, at this point, the patient is still excreting sugar in the twenty-four-hour urine (and twenty-four-hour specimens are necessary to an accurate diagnosis of the condition, and to successful treatment in the early part of it at least), he should be put on the diet with restricted protein (for subsequent steps, see p. 67). If, however, the urine is sugar-free after one or two days of carbohydrate-free diet, or before that point is reached, the patient is allowed the standard diet with 100 gm. ($3\frac{1}{3}$ oz.) of white bread divided between the three meals. The second twenty-four-hour specimen is then examined, and, if no sugar is present, increase the allowance of white bread by $\frac{1}{4}$ to $\frac{1}{2}$ ounce each day until sugar just shows in the urine. The last point at which the urine fails to show sugar represents that patient's carbohydrate tolerance.

The standard diet plus one-half of the carbohydrate tolerance is the diet this patient should be put on; *e. g.*, if the point of tolerance is 60 gm. (2 oz.) of white bread, then 30 gm. (1 oz.) is all the patient should be allowed, divided between the three meals. It is just here that the table of carbohydrate equivalents is of the greatest use, as it permits the use of foods more bulky than bread, but of relatively less carbohydrate value (p. 84).

The practical application in the use of this table in

the case taken as an example, with 30 gm. (1 oz.) as the amount of carbohydrates to be given, is as follows:

Example No. 1

(If tolerance is 30 gm. white bread.)

Proto puff No. 1...	45 gm.	=	7.5 gm.	white bread.
Potato.....	22 "	=	7.5 "	" "
Oatmeal.....	40 "	=	7.5 "	" "
Beets.....	33 "	=	3.75 "	" "
Peaches.....	25 "	=	3.75 "	" "
	165 "	=	30.00 "	" "

Example No. 2

(If tolerance is 45 gm. white bread.)

Potato.....	22 gm.	=	7.5 gm.	white bread.
Oatmeal.....	40 "	=	7.5 "	" "
Corn bread.....	20 "	=	15.0 "	" "
Carrots.....	65 "	=	7.5 "	" "
Orange.....	40 "	=	7.5 "	" "
	187 "	=	45.0 "	" "

That is, we can give a varied carbohydrate diet with a bulk of 165 gm. of food worth just 30 gm. of white bread, which is our allowance, or in the second example, 187 gm. mixed carbohydrates equals 45 gm. white bread.

This is sufficient for an example, as the practical application to each case is simple and only a matter of detail and a little practice.

This means, of course, that the carbohydrates must be weighed, and at the outset it must be understood

that this is the only satisfactory way to arrive at an accurate conclusion as to what constitutes the individual's carbohydrate tolerance and the only way to use the diet successfully. Of course, later the eye becomes well trained to judge amounts, and the scales may be dispensed with. (von Noorden, however, insists that patients must always live by the scale. A difficult thing to obtain in America.)

The reason that it is necessary to allow only one-half the amount of the carbohydrate tolerance is because, while the urine becomes sugar free on the full amount of the tolerance, the blood does not always, and, in order to reduce the sugar content of the blood to normal or as nearly as may be, the lessened amount of carbohydrate only is allowed. No increase should be made in the allowance until from one to three months, depending upon the severity of the disease, unless, of course, an acidosis should develop, which would require the use of the diet as outlined for the oatmeal days. The increase must be made gradually, the object being to allow little by little more carbohydrate, but not enough to have it show in the urine.

Should the urine not become sugar free on the standard diet without carbohydrates, the standard diet, with restricted protein, should be used. If this results in a sugar-free urine, then, after a short time, the full standard diet can be again tried, for often the restricted protein will make it possible for the organism to take more pro-

tein than at first without the appearance of sugar in the urine. From this point one proceeds as in the first instance.

In the severe cases, with more or less marked acidosis, it is best to put these patients at once on the oatmeal diet, and keep them on it from two to ten days, regardless of the sugar in the urine, giving them, at the same time, large doses of bicarbonate of soda in the attempt to render the urine alkaline, a thing always to be desired and tried for in these cases. Should the acidosis grow distinctly less, but the sugar still be considerable, the patients are usually benefited by two days on the green diet, "green days" as they are called. One goes from that to the standard diet, with restricted protein, if the lessened degree of acidosis will permit, and from there to the full standard diet, if possible, after a few days. If, however, the acidosis increases in the "green days," one must return to the oatmeal diet again until that danger is past.

Oftentimes a series of diets in succession are valuable, and von Noorden's routine in a severe case is two days restricted protein diet, two days green diet, three days oatmeal diet.

Often a patient becomes sugar free on this plan even while on the oatmeal, and the organic acids drop down very soon. These seven days constitute a "set." We then go back to the diet with restricted protein. If sugar appears again, repeat the "set"—by this means

one can often get a patient so they can again take a little carbohydrate and gradually increase it, always aiming to keep below the point of actual carbohydrate tolerance.

One other indication for the use of the oatmeal days is when in the course of the disease the patient's stomach or bowels become upset, nausea, vomiting, or diarrhea ensuing.

Wright's diabetic milk fills a long felt want in the treatment of diabetes, as the loss of milk from the diet is one of the most difficult features of the restricted diet to overcome. This milk tastes surprisingly like the fresh milk from which it is prepared, and, as the analysis shows, contains an almost negligible amount of carbohydrate. (See Special Recipes.)

Note.—When it is found inconvenient to prepare this milk at home, any good milk laboratory will make it up, or a similar milk may be purchased direct from D. Whiting & Sons, 570 Rutherford Avenue, Boston, Mass.

ANTICONSTIPATION DIET¹

- 6 A. M.: Fruit, stewed or fresh.
 1½ glasses of hot water.
- Breakfast: Oatmeal, molasses, coffee, bran biscuits.
 1 egg, baked apple or prunes, butter.
- 10 A. M.: 2 glasses of water.
- Dinner: Soup, meat or fish, all vegetables, salad
 and oil, brown bread and butter, any
 dessert, fruit.
- 4 P. M.: 2 glasses of water.
- Supper: Fruit, cereal, brown bread or bran biscuits
 and butter, molasses.
- Bedtime: Dried figs, prunes or dates, 1 glass of water.

Approximate Food Value to be Given


	Protein.	Carbohydrate.	Fat.	Calories.
Men. . . .	70 gm. (2½ oz.).	300 gm. (10 oz.).	80 gm. (2¾ oz.).	2200
Women..	60 gm. (2 oz.).	250 gm. (8½ oz.).	70 gm. (2½ oz.).	1800

COMMENTS ON ANTICONSTIPATION DIET

This can be greatly modified in detail according to the patient's financial circumstances. One great underlying trouble with most people who have atonic constipation (which constitutes by far the most common type) is that the colon and rectum need stimulation by

¹ Modified from Vanderbilt Clinic diet lists.

a larger food residue than the individual normally consumes, so that in general it may be said that these cases need fruits, cooked or raw, and green vegetables in double or treble the amount the normal individual does. It is only by constant effort and encouragement that patients can be brought to do this, as they feel they are eating too much food. This prejudice can be removed when they are told of the really slight food value of these articles of diet, and that they are mainly valuable on account of their bulk.



LOW CALCIUM DIET

Boiled meat.....	250 gm.	(8 $\frac{1}{3}$ oz.)
Bread.....	100 gm.	(3 $\frac{1}{3}$ oz.)
Fish.....	100 gm.	(3 $\frac{1}{3}$ oz.)
Potatoes.....	100 gm.	(3 $\frac{1}{3}$ oz.)
Apple.....	100 gm.	(3 $\frac{1}{3}$ oz.)
Sugar.....	50 gm.	(1 $\frac{2}{3}$ oz.)
Butter.....	50 gm.	(1 $\frac{2}{3}$ oz.)
Tea (with cream).....	100 gm.	(3 $\frac{1}{3}$ oz.)
Calcium content,	0.315 gm.	

Protein, 80 gm. (1 $\frac{2}{3}$ oz.); fat, 100 gm. (3 $\frac{1}{3}$ oz.); carbohydrate, 145 gm. (5 oz.); calories, 2000.

COMMENTS ON LOW CALCIUM DIET

This was designed by Dr. C. G. L. Wolf, of the Cornell Medical School, at the suggestion of my colleague, Dr. David Bovaird, Jr., who conceived the idea that cases of chronic deforming arthritis might be greatly benefited by reducing the calcium intake, on the theory that the disease is due to a disturbed calcium metabolism. This has been used in a number of cases, both at the Presbyterian and Lincoln Hospitals, New York, and, while it has not resulted in any cures, some of the patients have been greatly relieved of their pain while on the diet. Its monotony is its worst feature, and one

can seldom get patients to continue its use for long at a time. Theoretically, tetany might seem a possible consequence of withdrawing the calcium from the diet, but in one case in which Bovaird used this diet for many months there was no evidence of this.

The diet, moreover, is of scientific value to anyone desiring to study calcium metabolism.

The fixed nitrogen, calcium, magnesium diet has been used exclusively for research work, and is an accurate analysis of the food stuffs composing the diet in these three particulars.

FIXED NITROGEN, CALCIUM, MAGNESIUM DIET

(VOGEL.)

(Twenty-four-hour Amount)

Milk.....	500 gm. (16 $\frac{2}{3}$ oz.)
Eggs.....	300 gm. (10 oz.)
Cream.....	300 gm. (10 oz.)
Rice, uncooked.....	100 gm. (3 $\frac{1}{3}$ oz.)
N in twenty-four-hour amounts.....	8.138
CaO in twenty-four-hour amounts.....	1.214
MgO in twenty-four-hour amounts.....	0.175
Calories	1900

DIARRHEA

ACUTE DIARRHEA

It seems almost unnecessary to discuss this condition from a dietetic point of view, as the principles governing the feeding of such cases seem in adults generally so well understood. After an initial purge and a period of digestive rest one can give the ordinary fluid diet without milk, consisting of broths, egg-albumin in water flavored with orange- or lemon-juice, and then gruels, followed later by thin cereals, scraped meat, and toast. Fresh milk in any form usually does not agree, and is apt to increase or at least keep up the diarrhea. One should not be afraid of a generous amount of starvation, as physiologic rest to the bowel is often all that is needed.

CHRONIC DIARRHEA

Differing from acute diarrhea (where the cause is practically always from a gross dietetic error), the causes, of which chronic diarrhea is only a symptom, are numerous. In order to treat such a condition successfully one must make an accurate diagnosis, otherwise it is time and energy thrown away. One case in point: The chronic diarrhea, usually associated with a persistent gastric achylia, is, for the most part, promptly stopped

by merely giving dilute hydrochloric acid regardless of a diet; and often no amount of dieting of any description will control this diarrhea without supplying the lack of normal acid. Hence one sees the necessity for an accurate diagnosis preliminary to any form of treatment.

There are, however, numerous cases of chronic diarrhea when, regardless of the underlying cause, there is the need of a diet which fulfils the following conditions:

- (1) Non-stimulating and non-irritating.
- (2) Easily digested, leaving little residue.
- (3) Not apt to ferment.
- (4) As astringent as possible.

In order to fulfil these conditions the articles of food may be divided into those directly useful, those to be avoided, and those possibly allowable in certain cases.

DIET IN CHRONIC DIARRHEA

May take:

Raw oysters (soft part).

Clear soups.

Fish: Any soft, white-meated, non-oily variety.

Meat: Beef, lamb, chicken, guinea-hen.

Cereals: Farina, cream of wheat, grape-nuts, shredded wheat, wheatena, flaked rice, puffed wheat or rice, well-boiled rice.

Gruels: Vermicelli.

Bread: Stale bread, toast, zwieback, toasted crackers.

Cheese: Ripe, American, Canadian, cream or cottage, pineapple.

Desserts: Gelatin desserts made with little sugar, farina pudding or almost any cereal pudding, simple cake at times (*e. g.*, sponge), blanc mange.

Drinks: Water, tea, black coffee, cocoa, claret, dry sherry, Burgundy, whisky or brandy, and water.

To be avoided:

Milk (?): Some cases can take it boiled or skimmed.

Tough or oily fish, clams, etc.

Pork, veal, ham, duck, goose.

Pickles, condiments, salads, olives, etc.

Green vegetables, salads, fruits, fresh or stewed.

Hot breads, pies, sweets of all kinds.

Sweet wines.

Articles allowable in certain cases:

Malted milk can often be used on cereals, and is better borne than fresh milk. Fresh milk boiled.

Turkey, bacon (crisp), koumiss, zoolak, baked white potato. Eggs—boiled, baked, poached, omelette, scrambled.

Baked hubbard squash, well-stewed celery, creamed spinach, small boiled onions. Boiled peas or lima beans put through a colander, without skins.

OBESITY DIET
(RICHTER.)

- Early A. M.: Cupful of tea, no milk or sugar.
Ham, 40 gm. ($1\frac{1}{3}$ oz.).
1 roll (dry).
- 10 A. M.: Fresh fruit.
- 12 A. M.: Fresh fruit.
- 2 P. M.: Clear soup, 100 gm. ($3\frac{1}{3}$ oz.); lean meat,
plenty of green vegetables without
butter or milk.
Salad.
1 or 2 glasses of lemonade without sugar.
Use saccharin if wanted.
- 4 P. M.: Cup of tea.
- 6 P. M.: Fruit.
- 8 P. M.: 75 gm. ($2\frac{2}{3}$ oz.) of lean meat, radishes,
potato (small one), sour pickles.
Take one or two steam baths per week.
Protein, 75 gm. ($2\frac{1}{2}$ oz.); fat, 50 gm. ($1\frac{2}{3}$ oz.); carbo-
hydrate, 130 gm. ($4\frac{1}{2}$ oz.); calories, 1300.

COMMENTS ON OBESITY DIET

The accumulation of fat in the subcutaneous tissues is due either to an excess of food, combined with too little work, or it is a true disease, in which case fat is deposited in spite of a small ration. In the first instance,

diet alone is necessary for reduction, combined with exercise; in the second, there must also be other methods used.

In order to reduce the body weight it is necessary to know the number of heat units which that particular individual would need if the weight were the average for the height. This is easily obtained from the tables given (page 97).

Knowing what the average weight should be, the caloric needs are reckoned at 30 to 35 calories per kilo (2.2 pounds). This is sufficient for the body at light work. When the normal caloric need is found, a diet is constructed which will supply only two-thirds, one-half, or one-fourth the required normal; always, however, keeping the protein at a good quantity to prevent unnecessary protein loss, cutting down on the fats and carbohydrates so that the caloric value is kept low. Some authors prefer to make the greatest reduction in the fats, as they are intensive fat-building bodies; others prefer the reduction principally in carbohydrates. A generous supply of water is necessary, best taken between meals, as water with meals makes it easier to take larger quantities of food; hence, water-drinking at meal time is curtailed to the lowest possible amount or even none allowed.

The sample diet given furnishes a good quantity of protein, but is low in fat, carbohydrate, and total calories.

GASTRIC TEST MEALS

EWALD: On early morning fasting stomach, give
White bread (no crust)..... 40 gm. ($1\frac{1}{3}$ oz.)
Clear tea or water..... 400 c.c. (6 oz.)
No butter, no sugar or milk.
Express one hour later.

Leube:

Clear beef soup..... 400 c.c. ($13\frac{1}{3}$ oz.)
Beefsteak..... 200 gm. ($6\frac{2}{3}$ oz.)
Bread..... 50 gm. ($1\frac{2}{3}$ oz.)
Water..... 200 c.c. ($6\frac{2}{3}$ oz.)
Express six hours later.

Riegel:

1 bowl mutton broth..... 200 c.c. ($6\frac{2}{3}$ oz.)
Beefsteak..... 200 gm. ($6\frac{2}{3}$ oz.)
Mashed potatoes..... 50 gm. ($1\frac{2}{3}$ oz.)
1 roll or piece of bread,
1 glass of water..... 200 c.c. ($6\frac{2}{3}$ oz.)
Express six hours later.

COMMENTS ON TEST MEALS

The Ewald test meal is almost too well known to need any word of explanation, but an outline of its use and a word about routine in gastric diagnosis may not be amiss.

Patients presenting themselves for diagnosis should be told to take their usual supper or dinner the night before the examination, eating with the meal a helping of spinach and a few stewed prunes or dried raisins. These are all easily recognized in the early morning fasting stomach should there be any actual retention. Nothing must be taken into the stomach after this meal until the time of examination, not even water after 10 P. M. The following morning the stomach-tube is introduced, and, by expression and the use of a Politzer bag or stiff rubber bulb, an attempt made (without the addition of water) to obtain any possible gastric contents. Whatever is obtained is examined for mucus, free hydrochloric acid, total acidity, blood, and microscopically for leukocytes, food remnants, and bacteria. After this, or even in case there is nothing obtained from the fasting stomach, the Ewald test meal is given and removed one hour later without the addition of water.¹

¹The following should be about the normal limits of gastric chemistry after the Ewald meal:

Mucus.—None, or very small amount.

Expressed.—Not over 100 to 125 c.c.

Relation of Solids to Liquids.—1 to 2 or 1 to 3.

Free Hydrochloric Acid.—15 to 30 (reckoned in cubic centimeters of $\frac{1}{10}$ normal potassium hydrate).

Combined Acids.—20 to 30.

Total Acidity.—45 to 60 (reckoned in cubic centimeters of potassium hydrate solution).

Organic Acids.—Absent.

Reaction to Dilute Lugol's Solution.—Mahogany color (blue or purple, indicating poor starch digestion; reddish or yellow, good starch digestion).

The Riegel meal is often used for the purpose of learning what sort of gastric chemistry goes on in the presence of what might be called a normal mixed meal. At times the Ewald meal will show no free acid, whereas if meat is included this may call out enough acid, if possible, to give the reaction for free acid. Still another reason for giving the Riegel meal is that the Ewald meal is rather an abnormal meal and not so apt to call out the full or normally balanced secretion that the more normal meal might.

The difference between the result of the examination of this meal and the Ewald are: There may be higher acid values, the starch should be better digested, and blood reaction be present from the meat. Of course, too, the microscopic examination shows the meat residue.

This meal or the Leube may be given to test the motor efficiency of the stomach, as there should be nothing left after six and one-half to seven hours' digestion and the stomach should be empty.

Biuret Reaction.—Normal, a purple pink.

Blood Test.—No reaction or very faint.

Microscopic Examination.—Starch grains and scattered cell detritus (no noticeable number of leukocytes, bacteria, or sarcinæ should be present).

No food from previous day should be present.

For the interpretation of the divergences from the normal finding one is referred to the regular text-books on Gastric Diseases. Any food from the previous supper must be taken as an indication of mechanical obstruction at the pylorus, or possibly an extreme degree of motor insufficiency.

SCHMIDT'S INTESTINAL TEST DIET¹

IN the morning, 0.5 liter (16 oz.) milk, or, if milk does not agree, 0.5 liter (16 oz.) cocoa, prepared from 20 gm. ($\frac{2}{3}$ oz.) cocoa powder, 10 gm. ($\frac{1}{3}$ oz.) sugar, 400 c.c. (13 oz.) water, and 100 c.c. ($3\frac{2}{3}$ oz.) milk.

In the forenoon, 0.5 liter (16 oz.) oatmeal gruel, made from 40 gm. ($1\frac{1}{3}$ oz.) oatmeal, 10 gm. ($\frac{1}{3}$ oz.) butter, 200 c.c. ($6\frac{1}{2}$ oz.) milk, 300 c.c. (10 oz.) water, 1 egg, strained.

At noon, 125 gm. (4 oz.) chopped beef (raw weight), broiled rare with 20 gm. ($\frac{2}{3}$ oz.) of butter, so that the interior will still remain raw.

To this add 250 gm. (8 oz.) potato broth, made of 190 gm. ($6\frac{1}{3}$ oz.) mashed potatoes, 100 c.c. ($3\frac{1}{3}$ oz.) milk, and 10 gm. ($\frac{1}{3}$ oz.) butter.

In the afternoon as in the morning.

In the evening as in the forenoon.

This diet consists of:

Milk.....	1.5 liters ($1\frac{1}{2}$ qt.)
Zwieback.....	100 gm. ($3\frac{1}{3}$ oz.)
Eggs.....	2
Butter.....	50 gm. ($1\frac{2}{3}$ oz.)
Beef.....	125 gm. (4 oz.)
Potatoes.....	190 gm. ($6\frac{1}{3}$ oz.)
Oatmeal (gruel).....	80 gm. ($2\frac{2}{3}$ oz.)

This contains protein 102 gm. ($3\frac{1}{3}$ oz.); fat, 111 gm. (4 oz.); carbohydrates, 191 gm. ($6\frac{1}{3}$ oz.); calories, 2234.

¹ Schmidt, The Test Diet in Intestinal Diseases.

COMMENTS ON SCHMIDT'S INTESTINAL TEST DIET

Schmidt's intestinal test diet is a convenient and uniform diet for testing the intestinal functions, and gives as a starting-point a feces of uniform composition. In Schmidt's words, "such a test diet must be one which can be taken equally well by healthy people as well as those suffering from an intestinal disorder; it must be free, but not absolutely free, from waste matter, in order that the stimulus ordinarily furnished by the ingesta should not be completely absent; it must be capable of supplying the minimum, at least, of calories required by the body (at rest), and must contain the three chief groups of food stuffs in proportionate relation to each other; finally, it must be of simple composition, easy to make, and uniformly prepared." While this is not the only diet conceivable for such a purpose, there is a great advantage in the universal use of a particular diet for testing intestinal functions, just as we use standard diets for testing the gastric functions, and it is of a definite composition and the result of its use in many cases can be correlated.

TABLE OF CARBOHYDRATE EQUIVALENTS

WHITE bread and biscuits of various sorts:

	Grams	15	30	45	60	75
Ounces	$\frac{1}{4}$	$\frac{1}{2}$	1	$1\frac{1}{2}$	2	$1\frac{1}{2}$
	"	"	"	Grams.	"	"
Potato	22	44	88	132	176	220
Hominy (cooked)	25	50	100	150	200	250
Oatmeal (cooked)	40	80	160	240	320	400
Rice (cooked)	15	30	60	90	120	150
Farina (cooked)	25	50	100	150	200	250
Shredded wheat	5	10	20	30	40	50
Indian-meal mush	27	54	108	162	216	270
Macaroni	30	60	120	180	240	300
Corn bread	10	20	40	60	80	100
Barker's gluten food, A	102	204	408	612	816	1020
Barker's gluten food, B	74	148	296	444	592	740
Barker's gluten food, C	54	108	216	224	432	540
Almond meal	65	130	260	390	520	650
Gum gluten (ground)	12	24	48	72	96	120
Soja-bean meal	50	100	200	300	400	500
Casoid flour	55	110	220	330	440	550
Pure gluten biscuit	50	100	200	300	400	500
Protopuff No. 1	45	90	180	270	360	450
Protopuff No. 2	12	24	48	72	96	120
Salvia sticks	25	50	100	150	200	250
Milk (whole)	112	224	448	672	896	1120
Cream	112	224	448	672	896	1120
Grape-fruit weighed						
with skin	187	375	750	1125	1150	1875
Rice pudding	14	28	56	84	112	140
Tapioca pudding	15	30	60	90	120	150
Beets (cooked)	65	130	260	390	520	650
Custard (baked)	30	60	120	180	240	300
Carrots	65	130	260	390	520	650

Divide by 30 to reduce grams to ounces.

	Grams.					
Corn (canned or green)	22	44	88	132	176	220
Egg plant.	90	180	360	540	720	900
Parsnips.	35	70	140	210	280	350
Green peas.	30	60	120	180	240	300
Turnips.	56	112	224	336	448	560
Baked beans.	22	44	88	132	176	220
Apples.	45	90	180	270	360	450
Bananas.	20	40	80	120	160	200
Oranges.	40	80	160	240	320	400
Peaches.	50	100	200	300	400	500
Pears.	50	100	200	300	400	500
Prunes.	24	48	96	144	192	240
Watermelon.	225	450	900			

Divide by 30 to reduce grams to ounces.

The list of food stuffs taken from Bulletin 28 of the Department of Agriculture are indispensable in reckoning the protein, fat, and carbohydrate values of foods as well as their caloric value.

The special recipes are appended and will be found useful. Those for diabetics speak for themselves, while the lactose jelly and ice-cream will be found a convenient method for exhibiting palatably a good number of calories.

Delafield's mixture is often serviceable in the feeding of gastric cases accompanied by much nausea and vomiting, beginning with dram doses every twenty minutes and gradually increasing the dose, and later lengthening the intervals.

In Eiweissmilch (protein milk) we have a preparation of German origin which is useful, at times, in conditions of intestinal disturbance associated with diarrhea. It

has been used, as has Keller's malt soup, particularly in the feeding of children, but either may be of great use in the care of adults where milk, either alone or with the usual admixtures, is not well borne.

Fisher's table of "standard portions" of 100 calories each forms the basis for the easy computation and construction of any of the diets. The addition of the actual amount of protein, fat, and carbohydrate in the individual portions simplifies the determination of the amounts of these food elements in any one of the foods listed.

Thus, if, for example, we wish to figure the actual amount of food for a day's ration in an anticonstipation diet, we take the fraction or multiples of 100 calories of the foods called for. Slight rearrangements are often necessary to keep to the approximate values set for the different diets; the following is an example of the method for using this table—*e. g.*, for anticonstipation breakfast.

	Calories.	Protein. Gm.	Fat. Gm.	Carbohydrate. Gm.
Apple (1) (early A. M.) . . .	50	0.3	0.3	11.0
Oatmeal	100	4.4	0.8	8.2
Molasses	200	1.6	...	48.0
Biscuit (johnny cake) . . .	200	6.0	3.4	35.0
Egg	100	7.9	7.2	
Apple (1)	50	0.3	0.3	11.0
Prunes	200	1.4	...	46.8
Butter	100	...	10.6	
	<u>1000</u>	<u>21.9</u>	<u>22.6</u>	<u>160.0</u>

SPECIAL DIABETIC RECIPES¹

DIABETIC BREAD AND BISCUIT

AKOLL Biscuit (Huntly and Palmer).

Carbohydrate, 2.7 per cent.; nitrogen, 7 per cent.

Each biscuit weighs 5.1 gm. and contains 0.14 gm. carbohydrate and 0.41 gm. nitrogen.

Gluten meal biscuit, made from Barker's Gluten Food A, to be had of Herman B. Barker, Somerville, Mass.

Carbohydrate not over 4 per cent.; nitrogen, 13 per cent.

Soja-bean meal biscuit from Soja-bean meal, to be had of Thomas Metcalf Company, Boston, Mass.

Protein (N. \times 6.25), 44.64 per cent.; fat, 19.43 per cent.; cane-sugar, 9.34 per cent.; starch, none.

Casoid biscuit, to be had of Thomas Leeming & Co., New York City.

Carbohydrate, 8 per cent.; nitrogen, 10 per cent.

Pure gluten biscuit and potato gluten biscuit, to be had of Battle Creek Sanitarium Food Company.

Carbohydrate, 10 per cent.; nitrogen, 12 per cent.

Protopuff No. 1, Health Food Company, New York.

Carbohydrate, 10 per cent.; nitrogen, 12 per cent.

¹ T. C. Janeway in Musser and Kelly, Practical Therapeutics.

SPECIAL RECIPES FOR THE USE OF OATMEAL

OATMEAL POPOVERS:

Into the white of 1 egg beaten lightly stir 100 gm. ($3\frac{1}{3}$ oz.) of cooked oatmeal. Mix thoroughly. Drop into heated popover pan. Bake twenty minutes. Serve hot with butter.

OATMEAL GRIDDLE CAKES:

Into the beaten white of 1 egg stir 100 gm. ($3\frac{1}{3}$ oz.) of cooked oatmeal and 5 gm. (heaping teaspoonful) melted butter. Drop into griddle pan. Turn when the oatmeal is browned on edges. Serve hot with plenty of butter and cinnamon.

OATMEAL MUFFINS:

To 130 gm. (2 half-pint cups) of oatmeal, ground as finely as possible (coffee grinder), add 1 heaping teaspoonful Royal baking powder and $\frac{1}{2}$ teaspoonful of salt. Mix well and add $1\frac{1}{3}$ cups of cold water, and at the end melted butter or lard about half the size of an egg, 28 gm. (1 oz.). Beat well for a minute, put into buttered muffin pans, and bake in a very hot oven.

SOJA-BEAN MEAL

SOJA-BEAN MEAL BISCUIT:

1 cup of cream, 2 eggs, 1 teaspoonful of baking powder, salt to taste. Soja-bean meal to make a batter, not too thick. Make into 8 cakes.

MUFFINS:

Sift together thoroughly 1 oz. of Soja-bean flour, $\frac{1}{2}$ teaspoonful baking powder, and a little salt. Stir $1\frac{1}{4}$ oz. cold water into flour, add melted butter (amount equal to size of a hickory nut), beat the white of an egg and mix in lightly. Bake in two buttered muffin pans or molds.

PANCAKE:

Sift 1 tablespoonful of Soja-bean flour with a little salt, add water gradually until a thin batter is made, beat in hard the yolk of an egg, mix in lightly the beaten white of an egg. Put all or half of the mixture in a buttered frying-pan and bake to a nice brown on both sides.

PUDDING:

Mix $2\frac{1}{2}$ oz. of water, 1 oz. Soja-bean flour, and a little salt together cold, boil in a double boiler for one-half hour adding a piece of butter the size of a hickory nut. Beat up the yolk of an egg until foamy, add to it $\frac{1}{2}$ gr. saccharin dissolved in a few drops of water, a little grated nutmeg, and a little cinnamon. After the flour mixture has cooled somewhat stir it into the beaten yolk. Add the beaten white of an egg and bake in a fairly hot oven.

GLUTEN MEAL BISCUIT:

To 1 egg add 1 heaping saltspoonful of salt and beat; then add 6 tablespoonfuls of cold water and

beat until quite thick or until it becomes in quantity from 1 to 1½ pints, and into this beat 1 tablespoonful of thin cream; add 2 heaping tablespoonfuls of dry gluten; stir this into the previous mixture; stir occasionally during one-half hour until of the consistency of thick gruel; bake thirty-five minutes in well-buttered muffin pans in hot oven.

DESSERTS

BAKED CUSTARD:

- 1 egg.
- 3 tablespoonfuls cream.
- 5 tablespoonfuls water.
- 2 or 3 saccharin tablets.
- 8 drops vanilla essence.

Beat up well. Bake in buttered dish twenty minutes, with a little nutmeg grated on top.

COFFEE ICE-CREAM:

- 3 tablespoonfuls cream.
- 3 tablespoonfuls water.
- 2 tablespoonfuls coffee with 2 or 3 saccharin tablets dissolved in it.
- 1 egg.

Mix in saucepan and beat gradually until it thickens. Then cool and freeze.

CHEESE SAVORY:

Stir together until very light 4 tablespoonfuls each of melted butter and grated cheese with 1 table-

spoonful of Barker's gluten food A, 2 tablespoonfuls cream, a flavoring of salt and cayenne, and 2 well-beaten eggs. Pour into ramikins or cases made of oiled paper, bake in a quick oven, and serve immediately.

CRANBERRIES:

Stewed and sweetened with saccharin.

DIABETIC MILK (WRIGHT'S)

Definite quantity of milk diluted with three to four volumes distilled water, to which 1 or 2:1000 glacial acetic acid has been added (6-12 c.c., dr. $1\frac{1}{2}$ -3, glacial acetic acid; 1 pint or 500 c.c. water). This precipitates all casein and fat.


Allow to settle, then strain through calico. Wash repeatedly. Redissolve curd in a 1 per cent. solution of the following mixture, enough to make the original amount of milk:

Sodium chlorid.....	11.5
Potassium chlorid.....	9.9
Monopotassium phosphate.....	13.8
Dipotassium phosphate.....	10.0
Citrate of potassium.....	5.9
Dimagnesium phosphate.....	4.0
Magnesium citrate.....	4.4
Dicalcium phosphate.....	8.0
Tricalcium phosphate.....	9.6
Calcium citrate.....	25.5
Calcium oxid.....	5.5
Sodium carbonate.....	40.0

*Analysis of Wright's Diabetic Milk*¹

Specific gravity, 1011.

Ash.....	0.200	per cent.
Fat.....	3.600	"
Protein.....	1.907	"
Carbohydrates.....	0.015	"
Total solids.....	5.722	"
NaCl.....	0.110	"

¹ Granat.

MISCELLANEOUS RECIPES

RAISED BRAN BISCUITS:

2 cups of bran.

1 cup of entire wheat flour.

Butter, salt, and baking powder.

$\frac{1}{2}$ cup of molasses, more or less to taste.

$1\frac{1}{2}$ cups of milk.

Bake for one hour in gems, in slow oven.

(Eat one to three a day, when stale.)

BRAN CRACKERS (FOORD):

Butter, 2 cups.

Buttermilk, $\frac{3}{4}$ cup.

Molasses (or honey), $\frac{3}{4}$ cup.

White flour, 1 pint.

Baking soda, 2 teaspoonfuls.

Bran, $1\frac{1}{4}$ quarts.

Bake in flat crackers, $\frac{1}{4}$ inch thick and 2 inches wide.

CREAM OF PEA SOUP:

Canned peas, $\frac{1}{3}$ cup.

Salt.

Water, $1\frac{1}{2}$ oz. (45 c.c.).

Milk, 6 oz. (180 c.c.).

Butter, 1 dram (4 gm.).

Flour, 1 dram (4 gm.).

Lactose, 1 dram (4 gm.).

Boil first three ingredients, strain, and add milk, melted butter, flour, and lactose; season. Bring to the boiling-point and strain.

30 c.c. (1 oz.) equals protein 2.6 gm. Calories, 32.

BLACK BEAN SOUP:

Black beans, 2 tablespoonfuls.

Cold water, 1 cup.

Seasoning—onion, celery salt, salt, and pepper.

Boil beans with water, strain, and add—

Milk, $\frac{2}{3}$ cup.

Butter, 2 drams (8 gm.).

Flour, 1 dram (4 gm.).

Salt and pepper, q. s.

30 c.c. (1 oz.) equals protein 1.98. Calories, 40.

LACTOSE JELLY:

Gelatin, $\frac{1}{2}$ teaspoonful

Cold water, 2 teaspoonfuls

Orange juice, $1\frac{1}{2}$ oz.

Lemon juice, 1 teaspoonful

Cane-sugar, $1\frac{1}{2}$ teaspoonfuls.

Sherry, 1 teaspoonful.

Lactose, 3 tablespoonfuls ($1\frac{1}{2}$ oz.).

Boiling water, $1\frac{1}{2}$ oz.

} 1 portion.

Lactose is boiled in water until clear. Pour over soaked gelatin to dissolve and add other parts. Strain through double gauze and chill.

May also be flavored with grape or raspberry juice.

ARTIFICIAL BUTTERMILK:

1 quart fresh milk (fat-free preferable). Sterilize at temperature of 212° F. for twenty minutes. Cool to 98° F. and put in sterile bottle. Add 1 tube (1 c.c.) of lactobacillin (The Ferment Co.). Allow to stand at temperature of 95° F. for twenty-four hours. Beat thoroughly with Dover egg-beater and chill.

For large quantity, use $\frac{1}{2}$ bottle of lactobacillin to one can of milk.

PROTEIN MILK (EIWEISSMILCH)

Heat 1 quart of whole milk to 100° F. Add 4 teaspoonfuls of essence of pepsin and stir. Let the mixture stand at 100° F. until the curd has formed. Put the mass in a linen cloth and strain off the whey from the curd. Remove the curd from the linen cloth and press it through a rather fine sieve two or three times by means of a wooden mallet or spoon. Add 1 pint of water to the curd during this process. The mixture should now look like milk, and the precipitate must be very finely divided. Add 1 pint of buttermilk to this mixture.

The composition of this food is:

Fats.....	2.5	per cent.
Sugar.....	1.5	"
Protein.....	3.0	"
Salt.....	0.5	"

1 quart of this milk contains about 370 calories.

KELLER'S MALT SOUP

2 oz. of wheat flour are mixed with 11 oz. of whole milk and then passed through a sieve. In a second vessel 3 oz. of extract of malt are mixed with 20 oz. of warm water. The two mixtures are then poured into a porcelain vessel, $2\frac{1}{2}$ dr. of 11 per cent. carbonate of potassium added, and the whole cooked with constant stirring for twenty minutes, and then brought to a momentary boil; any loss through heat is made up by the addition of boiled water.

The composition of this food is:

Fats.....	1.20	per cent.
Protein.....	2.00	“
Carbohydrates.....	12.00	“

Caloric value, 800 to the liter.

LACTOSE ICE-CREAM

Cream.....	120 c.c.	(4 oz.)
Lactose.....	30 gm.	(1 oz.)
Boiling water.....	30 c.c.	(1 oz.)
Cane-sugar.....	4 gm.	(1 dr.)
Salt.....	1 gm.	($\frac{1}{4}$ dr.)
Vanilla.....	1 gm.	($\frac{1}{4}$ dr.)
Or,		
Sherry.....	15 c.c.	($\frac{1}{2}$ oz.)

Boil lactose and water two minutes or until clear. Mix with other ingredients. Freeze. Equals 390 calories.

DELAFIELD'S MIXTURE

Feeding in nausea:

Cream.....	120 c.c. (4 oz.)
Milk.....	120 c.c. (4 oz.)
Vichy.....	120 c.c. (4 oz.)
Soda bicarbonate.....	1 $\frac{1}{3}$ gm. (20 gr.)
Cerium oxalate.....	$\frac{2}{3}$ gm. (10 gr.)

AVERAGE WEIGHTS FOR MEN AND WOMEN

AS COMPILED BY THE METROPOLITAN LIFE INSURANCE
COMPANY

<i>Men.</i>		<i>Women.</i>	
Height.	Weight.	Height.	Weight.
Ft. In.	Lbs.	Ft. In.	Lbs.
5 1.....	120	4 10.....	108
5 2.....	125	4 11.....	112
5 3.....	130	5 0.....	114
5 4.....	135	5 1.....	118
5 5.....	141	5 2.....	123
5 6.....	145	5 3.....	126
5 7.....	150	5 4.....	129
5 8.....	154	5 5.....	133
5 9.....	159	5 6.....	137
5 10.....	164	5 7.....	142
5 11.....	169	5 8.....	146
6 0.....	175	5 9.....	150
6 1.....	181	5 10.....	154
6 2.....	188	5 11.....	158

AVERAGE CHEMICAL COMPOSITION OF AMERICAN FOODS

FROM ATWATER AND BRYANT (ABSTRACT), UNITED STATES DEPARTMENT OF AGRICULTURE

(Bulletin No. 28, 1906)

Food Material.	Water.	Protein.	Fat.	Carbohy- drates.	Calories per 100 gm. ¹
A. BEEF.	Per cent.	Per cent.	Per cent.	Per cent.	
<i>Fresh:</i>					
Chuck, including shoulder...	65.0	19.2	15.4	222
Loin.....	61.3	19.0	19.1	255
Sirloin butt, as purchased...	62.5	19.7	17.7	246
Porterhouse steak.....	60.0	21.9	20.4	280
Ribs.....	57.0	17.8	24.6	302
Round.....	67.8	20.9	18.6	184
<i>Beef Organs:</i>					
Brain.....	80.6	8.8	9.3	122
Kidney.....	76.7	16.6	4.8	0.4	115
Beef liver.....	71.2	20.4	4.5	1.7	133
Sweetbreads, as purchased..	70.9	16.8	12.1	181
Tongue.....	70.8	18.9	9.2	163
<i>Cooked:</i>					
Roast, as purchased.....	48.2	22.3	28.6	357
Round steak, fat removed, as purchased.....	63.0	27.6	7.7	185
Loin steak:					
Tenderloin, broiled.....	54.8	23.5	20.4	287
<i>Canned:</i>					
Boiled beef, as purchased...	51.8	25.5	22.5	314
Corned beef.....	51.8	26.3	18.7	282
Roast beef, as purchased...	58.9	25.9	14.8	243

¹ Locke, Food Values.

CHEMICAL COMPOSITION OF AMERICAN FOODS 99

Food Material.	Water.	Protein.	Fat.	Carbohy- drates.	Calories per 100 gm.
B. VEAL.	Per cent.	Per cent.	Per cent.	Per cent.	
<i>Fresh:</i>					
Breast.....	68.2	20.3	11.0	185
Leg.....	71.7	20.7	6.7	146
Loin.....	69.5	19.9	10.0	174
Rib.....	69.8	20.2	9.4	170
Shoulder and flank, medium fat.....	65.2	19.7	14.4	215
Kidney, as purchased.....	75.8	16.9	6.4	129
C. LAMB.					
<i>Fresh:</i>					
Breast or chuck.....	56.2	19.1	23.6	298
Leg, hind.....	58.6	18.6	22.6	287
Shoulder.....	51.8	18.1	29.7	351
Forequarter.....	55.1	18.3	25.8	315
Hindquarter.....	60.9	19.6	19.1	258
<i>Cooked:</i>					
Chops, broiled.....	47.6	21.7	29.9	367
Leg, roast.....	67.1	19.7	12.7	198
D. MUTTON.					
<i>Fresh:</i>					
Chuck, lean.....	64.7	17.8	16.3	225
Leg, hind.....	63.2	18.7	17.5	239
Shoulder.....	60.2	17.5	21.8	274
Forequarter.....	52.9	15.6	30.9	352
<i>Cooked:</i>					
Mutton, leg, roast.....	50.9	25.0	22.6	313
E. PORK.					
<i>Fresh:</i>					
Ham.....	50.1	15.7	33.4	375
<i>Pickled, salted, and smoked:</i>					
Ham, smoked, boiled, as pur- chased.....	51.3	20.2	22.4	291
fried, as purchased.....	36.6	22.2	33.2	400
Bacon, smoked.....	20.2	10.5	64.8	646

100 CHEMICAL COMPOSITION OF AMERICAN FOODS

Food Material.	Water.	Protein.	Fat.	Carbohy- drates.	Calories per 100 gm.
F. SAUSAGE (as purchased).	Per cent.	Per cent.	Per cent.	Per cent.	
Bologna.....	55.2	18.2	19.7	258
Frankfort.....	57.2	19.6	18.6	1.1	258
Pork.....	39.8	13.0	44.2	1.1	468
sausage meat.....	46.2	17.4	32.5	374

G. POULTRY.

Fresh:

Chicken, broiler.....	69.7	20.7	8.3	196
young, dark meat.....	70.1	20.8	8.2	187
light meat.....	70.3	21.9	7.4	184
Duck, breast.....	73.9	22.3	2.3	151
Guinea-hen meat, not includ- ing giblets.....	68.9	23.4	6.5	191
Pheasant meat, not including giblets.....	70.0	24.7	4.6	180
Pigeon meat, not including giblets.....	63.2	22.9	12.1	243
Quail meat, not including giblets.....	66.3	25.4	7.0	208
Squabs, meat, not including giblets.....	56.6	18.5	23.8	324
Turkey, dark meat.....	57.0	21.4	20.6	316
cooked.....	53.7	39.2	4.3	265
light meat.....	63.9	25.7	9.4	235
cooked.....	58.5	34.6	4.9	240

Preserved Poultry Meat:

Potted turkey.....	56.0	17.2	22.0	306
chicken.....	56.1	19.4	20.3	306
Canned chicken soup.....	87.1	2.9	3.3	5.1	66
gumbo soup.....	91.0	2.4	0.2	4.8	35
boned chicken.....	57.6	27.7	12.8	274

H. FISH.

Fresh:

Cod, whole.....	82.6	16.5	0.4	1.2	103
Bass, black, whole.....	76.7	20.6	1.7	1.2	103
sea, whole.....	79.3	19.8	0.5	1.4	86
striped, whole.....	77.7	18.6	2.8	1.2	102
Blackfish, whole.....	79.1	18.7	1.3	1.1	89

CHEMICAL COMPOSITION OF AMERICAN FOODS 101

Food Material.	Water.	Protein.	Fat.	Carbohy- drates.	Calories per 100 gm.
	Per cent.	Per cent.	Per cent.	Per cent.	
<i>Fresh:</i>					
Bluefish, entrails removed.	78.5	19.4	1.2	1.3	90
Butterfish, whole.	70.0	18.0	11.0	1.2	176
Eels, salt water.	71.6	18.6	9.1	1.0	161
Haddock, entrails removed.	81.7	17.2	0.3	1.2	74
Halibut, steak or sections.	75.4	18.6	5.2	1.0	125
Herring, whole.	72.5	19.5	7.1	1.5	146
Mackerel, whole.	73.4	18.7	7.1	1.2	142
Perch, white, whole.	75.7	19.3	4.0	1.2	117
as purchased.	28.4	7.3	1.5	0.4	44
yellow, whole.	79.3	18.7	0.8	1.2	84
Pickrel, pike, whole.	79.8	18.7	0.5	1.1	81
Pike, gray, whole.	80.8	17.9	0.8	1.1	80
Pompano, whole.	72.8	18.8	7.5	1.0	147
Pogy (scup), whole.	75.0	18.6	5.1	1.4	123
Salmon, whole.	64.6	22.0	12.8	1.4	209
Shad, whole.	70.6	18.8	9.5	1.3	165
roe, as purchased.	71.2	20.9	3.8	1.5	133
Smelt, whole.	79.2	17.6	1.8	1.7	89
Spanish mackerel, whole.	68.1	21.5	9.4	1.5	175
Trout, brook, whole.	77.8	19.2	2.1	1.2	98
<i>Preserved and canned:</i>					
Cod, salt.	53.5	25.4	0.3	90
Herring, smoked.	34.6	36.9	15.8	299
Mackerel, salt, dressed.	43.4	17.3	26.4	316
Salmon, canned.	63.5	21.8	12.1	201
Sardines, canned.	52.3	23.0	19.7	278
<i>Shellfish, etc., Fresh:</i>					
Clams, round, removed from shell, as purchased.	80.8	10.6	1.1	5.2	75
Oysters, solids, as purchased.	88.3	6.0	1.3	3.3	51
Scallops, as purchased.	80.3	14.8	0.1	3.4	76

I. EGGS.

Hens', uncooked.	73.7	13.4	10.5	159
boiled.	73.2	13.2	12.0	169
whites.	86.2	12.3	0.2	55
yolks.	49.5	15.7	33.3	376
Egg, boiled, 1 egg (50 gm.)	36.6	6.6	6.0	169
Total, 1 egg.					83

102 CHEMICAL COMPOSITION OF AMERICAN FOODS

Food Material.	Water. Per cent.	Protein. Per cent.	Fat. Per cent.	Carbohy- drates. Per cent.	Calories per 100 gm.
J. DAIRY PRODUCTS, ETC.					
(As Purchased).					
Butter.....	11.0	1.0	85.0	795
Buttermilk.....	91.0	3.0	0.5	4.8	36
Cheese, American, pale.....	31.6	28.8	35.9	0.3	453
red.....	28.6	29.6	38.3	477
California flat.....	34.0	24.3	33.4	4.5	429
Cheddar.....	27.4	27.7	36.8	4.1	473
Cheshire.....	37.1	26.9	30.7	0.9	399
Cottage.....	72.0	20.9	1.0	4.3	112
Dutch.....	35.2	37.1	17.7	316
Full cream.....	34.2	25.9	33.7	2.4	430
Limburger.....	42.1	23.0	29.4	0.4	369
Neuchatel.....	50.0	18.7	27.4	1.5	337
Roquefort.....	29.3	22.6	29.5	1.8	375
Swiss.....	31.4	27.6	24.9	1.3	443
Cream.....	74.0	2.5	18.5	4.5	201
Koumiss.....	89.3	2.8	2.1	5.4	53
Milk, condensed, sweetened..	26.9	8.8	8.3	54.1	335
unsweetened (evapor- ated cream).....	68.2	9.6	9.3	11.2	172
skimmed.....	90.5	3.4	0.3	5.1	37
whole.....	87.0	3.3	4.0	5.0	72
whey.....	93.0	1.0	0.3	5.0	28
K. MISCELLANEOUS (As Pur- chased).					
Beef-juice.....	93.0	4.9	0.6	25
Calf's-foot jelly.....	77.6	4.3	17.4	89
Oleomargarine.....	9.5	1.2	83.0	777
VEGETABLE FOOD.					
A. FLOUR, MEALS, ETC.					
Barley meal and flour.....	11.9	10.5	2.2	72.8	362
Buckwheat flour.....	13.6	6.4	1.2	77.9	357
Cornmeal, granular.....	12.5	9.2	1.9	75.4	365
Corn Preparations:					
Cerealine.....	10.3	9.6	1.1	78.3	370
Hominy.....	11.8	8.3	0.6	79.0	364
cooked.....	79.3	2.2	0.2	17.8	84

CHEMICAL COMPOSITION OF AMERICAN FOODS 103

Food Material.	Water. Per cent.	Protein. Per cent.	Fat. Per cent.	Carbohy- drates. Per cent.	Calories per 100 gm.
Corn Preparations: Oatmeal..	7.3	16.1	7.2	67.5	410
boiled.....	84.5	2.8	0.5	11.5	63
gruel.....	91.6	1.2	0.4	6.3	34
water.....	96.0	0.7	0.1	2.9	15
Rolled oats.....	7.7	16.7	7.3	66.2	408
Rice.....	12.3	8.0	0.3	79.0	359
boiled.....	72.5	2.8	0.1	24.4	112
flaked.....	9.5	7.9	0.4	81.9	371
flour.....	8.5	8.6	6.1	68.0	370
Rye flour.....	12.9	6.8	0.9	78.7	359
meal.....	11.4	13.6	2.0	71.5	367
Wheat, entire.....	11.4	13.8	1.9	71.9	369
gluten.....	12.0	14.2	1.8	71.1	367
Graham.....	11.3	13.3	2.2	71.4	368
Prepared (self-raising).....	10.8	10.2	1.2	73.0	353
Wheat Preparations:					
Cracked and crushed.....	10.1	11.1	1.7	75.5	371
Farina.....	10.9	11.0	1.4	76.3	371
Flaked.....	8.7	13.4	1.4	74.3	373
Gems.....	10.4	10.5	2.0	76.0	374
Glutens.....	8.9	13.6	1.7	74.6	378
Macaroni.....	10.3	13.4	0.9	74.1	367
cooked.....	78.4	3.0	1.5	15.8	91
Noodles.....	10.7	11.7	1.0	75.6	367
Shredded.....	8.1	10.5	1.4	77.9	375
Spaghetti.....	10.6	12.1	0.4	76.3	366
Vermicelli.....	11.0	10.9	2.0	72.0	358

B. BREAD, CRACKERS, PASTRY,
ETC. (As Purchased).

Bread:

Brown.....	43.6	5.4	1.8	47.1	231
Corn (johnny cake).....	38.9	7.9	4.7	46.3	266
Rye.....	35.7	9.0	0.6	53.2	260
Wheat:					
Buns.....	29.0	6.3	6.5	57.3	321
Cinnamon.....	23.6	9.4	7.2	59.1	347
Currant.....	27.5	6.7	7.6	57.6	334

104 CHEMICAL COMPOSITION OF AMERICAN FOODS

Food Material.	Water.	Protein.	Fat.	Carbohy- drates.	Calories per 100 gm.
VEGETABLE FOOD.	Per cent.	Per cent.	Per cent.	Per cent.	
Bread, Wheat: Hot cross.	36.7	7.9	4.8	49.7	281
Graham.	35.7	8.9	1.8	52.1	267
Biscuit, home made.	32.9	8.7	2.6	55.3	287
soda.	22.9	9.3	13.7	52.6	381
Rolls, French.	32.0	8.5	2.5	55.7	287
Vienna.	31.7	8.5	2.2	56.5	287
White, biscuit.	35.2	8.0	1.4	54.3	269
home-made.	35.0	9.1	1.6	53.3	270
all analyses.	35.3	9.2	1.3	53.1	268
Whole wheat.	38.4	9.7	0.9	49.7	251
Zwieback.	5.8	9.8	9.9	73.5	434
Crackers:					
Soda.	5.9	9.8	9.1	73.1	424
Boston (split).	7.5	11.0	8.5	71.1	416
Egg.	5.8	12.6	14.0	66.6	454
Graham.	5.4	10.0	9.4	73.8	429
Oatmeal.	6.3	11.8	11.1	69.0	434
Oyster.	4.8	11.3	10.5	70.5	433
Pretzels.	9.6	9.7	3.9	72.8	375
Saltines.	5.6	10.6	12.7	68.5	442
Water.	6.4	11.7	5.0	75.7	405
All analyses.	6.8	10.7	8.8	71.9	420
Cake:					
Bakers'	31.4	6.3	4.6	59.9	302
Chocolate layer.	20.5	6.2	8.1	64.1	364
Drop.	16.6	7.6	14.7	60.3	316
Frosted.	18.2	5.9	9.0	64.8	374
Fruit.	17.3	5.9	10.9	64.1	388
Gingerbread.	18.8	5.8	9.0	63.5	368
Sponge.	15.3	6.3	10.7	65.9	396
Cookies, Cakes, etc.:					
Mollasses cookies.	6.2	7.2	8.7	75.7	421
Sugar cookies.	8.3	7.0	10.2	73.2	423
Ginger snaps.	6.3	6.5	8.6	76.0	418
Lady fingers.	15.0	8.8	5.0	70.6	371
Macaroons.	12.3	6.5	15.2	65.2	435
Doughnuts.	18.3	6.7	21.0	53.1	441

CHEMICAL COMPOSITION OF AMERICAN FOODS 105

Food Material.	Water.	Protein.	Fat.	Carbohy- drates.	Calories per 100 gm.
Pie:	Per cent.	Per cent.	Per cent.	Per cent.	
Apple.....	42.5	3.1	9.8	42.8	280
Cream.....	32.0	4.4	11.4	51.2	334
Custard.....	62.4	4.2	6.3	26.1	183
Lemon.....	47.4	3.6	10.1	37.4	262
Mince.....	41.3	5.8	12.3	38.1	294
Squash.....	64.2	4.4	8.4	21.7	185
Puddings:					
Rice custard.....	59.4	4.0	4.6	31.4	182
Indian meal.....	60.7	5.5	4.8	27.5	180
Tapioca.....	64.5	3.3	3.2	28.2	159
Tapioca with apple.....	70.1	0.3	0.1	29.3	122
Ice-cream.....	66.9	5.2	10.1	17.7	189

C. SUGARS, STARCHES, ETC.

(As Purchased)

Candy.....	96.0	391
Honey.....	18.2	0.4	81.2	335
Mollasses, cane.....	25.1	2.4	69.3	284
Starch, tapioca.....	11.4	0.4	0.1	88.0	364
Sugar, coffee or brown.....	95.0	389
granulated.....	100.0	410
maple.....	82.8	339
powdered.....	100.0	410

D. VEGETABLES.

Artichokes, as purchased.....	79.5	2.6	0.2	16.7	80
Asparagus, cooked, as pur- chased.....	91.1	2.1	3.3	2.2	48
Beans, butter, green.....	58.9	9.4	0.6	29.1	163
String beans, cooked.....	95.3	0.8	1.1	1.9	21
fresh, as purchased.....	83.0	2.1	0.3	6.9	40
Beets, cooked.....	88.6	2.3	0.1	7.4	41
Cabbage.....	91.5	1.6	0.3	5.6	32
Carrots, fresh.....	88.2	1.1	0.4	9.3	46
Cauliflower, as purchased.....	92.3	1.8	0.5	4.7	31
Celery.....	94.5	1.1	0.1	3.3	19
Corn, green.....	75.4	3.1	1.1	19.7	104
Cucumbers.....	95.4	0.8	0.2	3.1	18

106 CHEMICAL COMPOSITION OF AMERICAN FOODS

Food Material.	Water.	Protein.	Fat.	Carbohy- drates.	Calories per 100 gm.
VEGETABLE FOOD.	Per cent.	Per cent.	Per cent.	Per cent.	
Eggplant.....	92.9	1.2	0.3	5.1	29
Greens, beet, cooked, as pur- chased.....	89.5	2.2	3.4	3.2	54
Lentils, dried, as purchased....	8.4	25.7	1.0	59.2	357
Lettuce.....	94.7	1.2	0.3	2.9	20
Mushrooms, as purchased....	88.1	3.5	0.4	6.8	46
Okra.....	90.2	1.6	0.2	7.4	39
Onions, fresh.....	87.6	1.6	0.3	9.9	49
prepared, as purchased....	91.2	1.2	1.8	4.9	42
Parsnips.....	83.0	1.6	0.5	13.5	66
Peas, dried, as purchased.....	9.5	24.6	1.0	62.0	365
green.....	74.6	7.0	0.5	16.9	102
cooked, as purchased....	73.8	6.7	3.4	14.6	119
Potatoes, raw or fresh cooked..	78.3	2.2	0.1	18.4	85
boiled, as purchased.....	75.5	2.5	0.1	20.9	97
cooked chips, as purchased..	2.2	6.8	39.8	46.7	589
mashed and creamed, as purchased.....	75.1	2.6	3.0	17.8	111
sweet, raw or fresh.....	69.0	1.8	0.7	27.4	126
cooked and prepared, as purchased.....	51.9	3.0	2.1	42.1	204
Pumpkins.....	93.1	1.0	0.1	5.2	26
Radishes.....	91.8	1.3	0.1	5.8	30
Rhubarb.....	94.4	0.6	0.7	3.6	23
Sauerkraut, as purchased....	88.8	1.7	0.5	3.8	28
Spinach, fresh, as purchased..	92.3	2.1	0.3	3.2	24
cooked, as purchased.....	89.8	2.1	4.1	2.6	57
Squash.....	88.3	1.4	0.5	9.0	47
Tomatoes, fresh, as purchased..	94.3	0.9	0.4	3.9	23
Turnips.....	89.6	1.3	0.2	8.1	41
<i>Canned as Purchased.</i>					
Asparagus.....	94.4	1.5	0.1	2.8	19
Beans, baked.....	68.9	6.9	2.5	19.6	132
string.....	93.7	1.1	0.1	3.8	21
lima.....	79.5	4.0	0.3	14.6	79
red kidney.....	72.7	7.0	0.2	18.5	106
Brussels sprouts.....	93.7	1.5	0.1	3.4	21
Corn, green.....	76.1	2.8	1.2	19.0	103
Okra.....	94.4	0.7	0.1	3.6	19

CHEMICAL COMPOSITION OF AMERICAN FOODS 107

Food Material.	Water. Per cent.	Protein. Per cent.	Fat. Per cent.	Carbohy- drates. Per cent.	Calories per 100 gm.
Peas, green.....	85.3	3.6	0.2	9.8	56
Pumpkins.....	91.6	0.8	0.2	6.7	33
Squash.....	87.6	0.9	0.5	10.5	52
Succotash.....	75.9	3.6	1.0	18.6	103
Tomatoes.....	94.0	1.2	0.2	4.0	23

F. FRUITS, BERRIES.

Apples:					
Edible portion.....	84.6	0.4	0.5	14.2	64
As purchased (refuse, 25.0).....	63.3	0.3	0.3	10.8	49
Apricots.....	85.0	1.1	...	13.4	59
Bananas:					
Edible portion.....	75.3	1.3	0.6	22.0	101
As purchased (refuse, 35.0).....	48.9	0.8	0.4	14.3	66
Blackberries, as purchased.....	86.3	1.3	1.0	10.9	15
Cherries, as purchased.....	76.8	0.9	0.8	15.9	76
Cranberries, as purchased.....	88.9	0.4	0.6	9.9	47
Currants, as purchased.....	85.0	1.5	...	12.8	58
Figs, fresh, as purchased.....	79.1	1.5	...	18.8	84
Grapes, as purchased.....	58.0	1.0	1.2	14.4	74
Huckleberries.....	81.9	0.6	0.6	16.6	76
Lemons:					
Edible portion.....	89.3	1.0	0.7	8.5	45
As purchased (refuse, 30.0).....	62.5	0.7	0.5	5.9	32
Muskmelons:					
Edible portion.....	89.5	0.6	...	9.3	41
As purchased (refuse, 50.0).....	44.8	0.3	...	4.6	20
Nectarines.....	82.9	0.6	...	15.9	67
Oranges:					
Edible portion.....	86.9	0.8	0.2	11.6	53
As purchased (refuse, 27.0).....	63.4	0.6	0.1	8.5	37
Peaches:					
Edible portion.....	69.4	0.7	0.1	9.4	42
As purchased (refuse, 18.0).....	73.3	0.5	0.1	7.7	34
Pears:					
Edible portion.....	64.4	0.6	0.5	14.1	65
As purchased (refuse, 10.0).....	76.0	0.5	0.4	12.7	57
Pineapple.....	89.3	0.4	0.3	9.7	44

108 CHEMICAL COMPOSITION OF AMERICAN FOODS

Food Material.	Water.	Protein.	Fat.	Carbohy- drates.	Calories per 100 gm.
VEGETABLE FOOD.	Per cent.	Per cent.	Per cent.	Per cent.	
Prunes.....	79.6	0.9	...	18.9	81
Raspberries, as purchased.....	85.8	1.0	...	12.6	56
Strawberries.....	90.4	1.0	0.6	7.4	40
Watermelons:					
Edible portion.....	92.4	0.4	0.2	6.7	31
As purchased (refuse, 59.4).....	37.5	0.2	0.1	2.7	13
<i>Dried.</i>					
Apples, as purchased.....	28.1	1.6	2.2	66.1	298
Apricots, as purchased.....	29.4	4.7	1.0	62.5	284
Citron, as purchased.....	19.0	0.5	1.5	78.1	336
Currants, as purchased.....	17.2	2.4	1.7	74.2	330
Dates.....	15.4	2.1	2.8	78.4	336
Figs, as purchased.....	18.8	4.3	0.3	74.2	325
Pears.....	16.5	2.8	5.4	72.9	360
Prunes.....	22.3	2.1	...	73.3	309
Raisins.....	14.6	2.6	3.3	76.1	354
<i>Canned and Jellies.</i>					
Preserves, etc., as purchased:					
Apples, crab.....	42.4	0.3	2.4	54.4	247
sauce.....	61.1	0.2	0.8	37.2	161
Apricots.....	81.4	0.9	...	17.3	75
Blackberries.....	40.0	0.8	2.1	56.4	254
Blueberries.....	85.6	0.6	0.6	12.8	61
Cherries.....	77.2	1.1	0.1	21.1	91
Figs, stewed.....	56.5	1.2	0.3	40.9	173
Marmalade (orange peel).....	14.5	0.6	0.1	84.5	349
Peaches.....	88.1	0.7	0.1	10.8	49
Pears.....	81.1	0.3	0.3	18.0	78
Strawberries, stewed.....	74.8	0.7	...	24.0	101
Prune sauce.....	76.6	0.5	0.1	22.3	95
G. NUTS.					
Almonds.....	4.8	21.0	54.9	17.3	668
Beechnut.....	4.0	21.9	57.4	13.2	678
Brazil nuts (<i>Bertholletia ex-</i> <i>celsa</i>), edible portion.....	5.3	17.0	66.8	7.0	720
Butternuts:					
<i>Juglans cinerea</i>	4.4	27.9	61.2	3.5	698

CHEMICAL COMPOSITION OF AMERICAN FOODS 109

Food Material.	Water.	Protein.	Fat.	Carbohy- drates.	Calories per 100 gm.
	Per cent.	Per cent.	Per cent.	Per cent.	
Chestnuts:					
Edible portion.....	45.0	6.2	5.4	42.1	246
As purchased (refuse, 16.0).....	37.8	5.2	4.5	35.4	208
Cocoanuts.....	14.1	5.7	50.6	27.9	608
Cocoanut, prepared, as pur- chased.....	3.5	6.3	57.4	31.5	689
Filberts.....	3.7	15.6	65.3	13.0	725
Hickory nuts.....	3.7	15.4	67.4	11.4	737
Lichi nuts.....	17.9	2.9	0.2	77.5	332
Peanuts:					
Edible portion.....	9.2	25.8	38.6	24.4	564
As purchased (refuse, 24.5).....	6.9	19.5	29.1	18.5	427
Peanut butter, as purchased....	2.1	29.3	46.5	17.1	623
Walnuts, California.....	2.5	18.4	64.4	13.0	728

H. MISCELLANEOUS.

Chocolate.....	5.9	12.9	48.7	30.3	631
Cocoa.....	4.6	21.6	28.9	37.7	511

UNCLASSIFIED FOOD
MATERIALS.

ANIMAL AND VEGETABLE.

A. SOUPS.

Home Made (As Purchased).

Beef.....	92.9	4.4	0.4	1.1	26
Bean.....	84.3	3.2	1.4	9.4	65
Chicken.....	84.3	10.5	0.8	2.4	61
Chowder, clam.....	88.7	1.8	0.8	6.7	43
Meat stew.....	84.5	4.6	4.3	5.5	81

Canned (As Purchased).

Asparagus, cream of.....	87.4	2.5	3.2	5.5	63
Bouillon.....	96.6	2.2	0.1	0.2	11
Celery, cream of.....	88.6	2.1	2.8	5.0	55
Chicken.....	93.8	3.6	0.1	1.5	22
gumbo.....	89.2	3.8	0.9	4.7	43
Consommé.....	96.0	2.5	...	0.4	12
Corn, cream of.....	86.8	2.5	1.9	7.8	59
Julienne.....	95.9	2.7	...	0.5	13
Mock turtle.....	89.8	5.2	0.9	2.8	41

110 CHEMICAL COMPOSITION OF AMERICAN FOODS

Food Material.	Water.	Protein.	Fat.	Carbohy- drates.	Calories per 100 gm.
UNCLASSIFIED FOOD MATERIAL.	Per cent.	Per cent.	Per cent.	Per cent.	
Mulligatawny.....	89.3	3.7	0.1	5.7	40
Oxtail.....	88.8	4.0	1.3	4.3	46
Pea.....	86.9	3.6	0.7	7.6	52
cream of green.....	87.7	2.6	2.7	5.7	60
Tomato.....	90.0	1.8	1.1	5.6	41
Turtle, green.....	86.6	6.1	1.9	3.9	58
Vegetable.....	95.7	2.9	0.5	14

B. MISCELLANEOUS.

Hash.....	80.3	6.0	1.9	9.4	80
Minceat, commercial.....	27.7	6.7	1.4	60.2	288
home-made.....	54.4	4.8	6.7	32.1	214
Salad, ham.....	69.4	15.4	7.6	5.6	157
Sandwich, egg.....	41.4	9.6	12.7	34.5	299
chicken.....	48.5	12.3	5.4	32.1	232

STANDARD PORTIONS*

TABLE OF FOODS GIVING WEIGHT (IN GRAMS, OUNCES, AND ROUGH MEASURE OF A "STANDARD PORTION" OF EACH FOOD AND THE ACTUAL AMOUNT OF PROTEIN, FAT, AND CARBOHYDRATE IN EACH PORTION

Name of Food.	Portion containing 100 calories roughly described.	Weight of 100 calories.	Actual amount of protein, fat, and carbohydrate in each portion of 100 calories.†	
			Protein.	Fat.
		Gm.	Gm.	Gm.
COOKED MEATS.				
1 Beef, round, boiled (fat).....	Small serving.....	36	9.1	7.1
1 Beef, round, boiled (lean).....	Large serving.....	62	17.1	4.7
1 Beef, round, boiled (medium).....	Small serving.....	44	11.2	8.8
1 Beef, ribs, boiled.....	Small serving.....	30	5.3	7.3
2 Chicken, as purchased, canned.....	One thin slice.....	27	7.4	3.4
2 Lamb chops, boiled, edible portion, average.....	One small chop.....	27	5.8	8
2 Lamb, leg roast.....	Ordinary serving.....	50	9.8	6.3
1 Mutton, leg, boiled.....	Large serving.....	34	8.5	7.6
1 Pork, ham, boiled (fat).....	Small serving.....	20.5	4.1	4.5
1 Pork, ham, roasted (fat).....	Small serving.....	27	4.4	10.8
2 Turkey, as purchased, canned.....	Small serving.....	28	3.7	8.4
1 Veal, leg, boiled.....	Large serving.....	67.5	13.9	4.5

* Irving Fisher, Journal American Medical Association, 1907, vol. xlviii, p. 1320.

† U. S. Department Agriculture, Bulletin No. 28, 1906.

1 Experiments on Losses in Cooking Meats, Grindley, U. S. Dep't Agricult., Bull. No. 141.

2 Chemical Composition of American Food Stuffs, Atwater and Bryant, U. S. Dep't Agricult., Bull. No. 28.

Name of Food.	Portion containing 100 calories roughly described.	Weight of 100 calories.		Actual amount of protein, fat, and carbohydrate in each portion of 100 calories.		
		Gm.	Oz.	Protein.	Fat.	Carbohydrates.
UNCOOKED MEATS.						
2 Chicken, broilers, edible portion.....	Large serving.....	90	3.2	18.6	7.4	
2 Beef, loin, edible portion, average.....	Ordinary serving.....	50	1.8	9.5	9.6	
2 Beef, ribs, edible portion, average.....	Ordinary serving.....	52	1.8	9.2	13.7	
2 Beef, sirloin steak.....	Small steak.....	40	1.4	7.5	7.4	
2 Mutton, leg, hind, lean, edible portion, av.....	Ordinary serving.....	50	1.8	9.3	8.7	
2 Liver, veal, as purchased.....	Two small servings.....	79	2.8	15	39	
2 Cod, whole, edible portion.....	Two servings.....	138	4.9	15	4.1	
2 Halibut steaks, average.....	Ordinary serving.....	81	2.8	15	4.2	
2 Mackerel (Spanish).....	Ordinary serving.....	57	2.0	12.2	5.3	
2 Pork, ham, smoked, lean, edible portion, av.....	Small serving.....	36	1.3	5.9	13.9	
2 Pork, bacon, smoked, medium fat, average.....	Small serving.....	15	0.53	1.5	9.7	
DAIRY PRODUCTS.						
2 Butter, as purchased.....	Ordinary pat or ball.....	12.5	0.44	0.1	10.6	
2 Buttermilk, as purchased.....	1½ glasses.....	275	9.7	8.2	1.3	13.2
2 Cheese, American, pale, as purchased.....	1½ cubic inches.....	22	0.77	6.3	7.8	0.06
2 Cheese, cottage, as purchased.....	4 cubic inches.....	89	3.12	18.6	0.8	3.8
2 Cheese, full cream, as purchased.....	1½ cubic inches.....	23	0.82	5.9	7.7	0.55
2 Cream.....	¼ ordinary glass.....	49	1.7	1.2	9.0	2.2
2 Milk, whole, as purchased.....	Small glass.....	140	4.9	4.7	5.6	7.0
2 Milk, condensed, sweetened, as purchased.....	30	1.06	2.6	2.4	16.2

STANDARD PORTIONS

113

2 Milk, condensed, unsweetened, evaporated cream, as purchased.....	59	2.05	5.7	5.5	6.7
2 Milk, skimmed, as purchased.....	255	9.4	8.6	7.6	13.0
CEREALS.					
2 Bread, corn (johnny cake), as purchased.....	38	1.3	3	1.7	17.5
2 Bread, white, home-made, as purchased.....	38	1.3	3.4	0.6	20.2
2 Cornmeal, granular, average.....	27	0.96	2.5	0.5	20.3
2 Crackers, graham, as purchased.....	23	0.82	2.3	2.1	16.8
2 Crackers, oatmeal, as purchased.....	23	0.81	2.7	2.5	15.8
2 Hominy, cooked.....	120	4.2	2.6	0.2	21.3
2 Oatmeal, boiled, average.....	159	5.6	4.4	0.8	18.2
2 Rice, boiled, average.....	87	3.1	2.4	0.8	21.2
2 Rice, uncooked.....	28	0.98	2.2	0.8	22
2 Shredded wheat.....	27	0.94	3.5	0.4	22.5
2 Rolls, Vienna.....	35	1.2	2.9	0.77	19.7
2 Spaghetti, average.....	28	0.97	3.4	0.11	21.3
2 Macaroni, cooked, average.....	110	3.8	3.3	1.6	17.3
2 Zwieback.....	23	0.81	2.2	2.3	16.9
2 Wheat flour, white, average.....	27	0.97	2.9	0.3	20.2
VEGETABLES.					
2 Asparagus, as purchased, average.....	206	7.1	4.3	6.7	4.5
2 Beans, baked.....	75	2.6	5.2	1.9	14.7
2 Beans, string, cooked.....	480	16.6	3.8	5.3	9.1

* Chemical Composition of American Food Stuffs, Atwater and Bryant, U. S. Dep't Agricult., Bull. No. 28.

Name of Food.	Portion containing 100 calories roughly described.		Weight of 100 calories.		Actual amount of protein, fat, and carbohydrate in each portion of 100 calories.	
	Gm.	Oz.	Protein.	Fat.	Carbohydrates.	Gm.
VEGETABLES						
2 Beans, lima, canned.....	Large side dish.....	126	4.4	5.4	0.4	18.4
2 Beets, edible portion, cooked.....	Three servings.....	245	8.7	5.6	0.2	18.3
2 Cabbage, edible portion, cooked.....	Three servings.....	310	11.0	4.9	0.9	17.3
2 Carrots, edible portion, cooked.....	Two servings.....	164	5.8	1.8	0.65	15.3
2 Celery, edible portion.....		549	19	6	0.5	17.8
2 Cauliflower, as purchased.....		312	11	5.6	1.5	14.6
2 Corn, sweet, cooked.....	One side dish.....	99	3.5	2.8	1.2	19
2 Eggplant, edible portion, average.....		350	12	4.2	1	17.8
2 Lentils, cooked.....		89	3.1	22.9	0.9	52.6
2 Onions, fresh, edible portion, average.....		200	7.1	3.2	0.6	19.8
2 Onions, cooked.....	Two large servings.....	240	8.4	2.9	4.3	11.7
2 Parsnips, edible portion, average.....	One and a half servings.....	152	5.3	2.4	0.7	20.5
2 Peas, green, cooked.....	One serving.....	85	3	5.7	2.9	12.4
2 Potatoes, baked.....	One, good sized.....	86	3	1.7	0.08	15.8
2 Potatoes, boiled.....	One, large sized.....	102	3.6	2.5	0.1	21.3
2 Potatoes, mashed and creamed.....	One serving.....	89	3.1	2.3	2.6	15.8
2 Potato chips.....	One-half serving.....	17	0.6	1.1	6.7	8
2 Potatoes, sweet, cooked.....	One-half average potato.....	49	1.7	1.4	1	20.6
2 Spinach, cooked, as purchased.....	Two ordinary servings.....	174	6.1	3.6	7.13	4.5
2 Squash, edible portion, average.....		210	7.4	2.9	1	18.9

2 Succotash, canned, as purchased.....	Ordinary serving.....	100	3.5	3.6	1	18.6
2 Tomatoes, fresh, as purchased, average.....	Four average tomatoes.....	430	15	3.9	1.7	16.7
Tomatoes, canned.....	431	15.2	5.1	0.86	17.2
2 Turnips, edible portion, average.....	Two large servings.....	246	8.7	3.1	0.5	20
FRUITS (DRIED)						
2 Apples, as purchased, average.....	34	1.2	0.54	0.74	22.4
2 Apricots, as purchased, average.....	35	1.2	1.6	3	21.47
2 Figs, edible portion.....	One large one.....	31	1.1	1.3	23
2 Prunes, edible portion.....	Three large ones.....	32	1.1	0.7	23.4
2 Raisins, as purchased.....	31	1.1	0.8	1	23.6
FRUITS (FRESH OR COOKED)						
2 Apples, as purchased.....	Two apples.....	206	7.3	0.6	0.6	22
Apple sauce.....	Ordinary serving.....	111	3.9	0.2	0.8	41.3
Apricots, cooked.....	Large serving.....	131	4.6	1.1	22.6
2 Bananas, yellow, edible portion, average.....	One large.....	100	3.5	1.3	0.6	22
2 Blackberries, as purchased, average.....	170	5.9	2.2	1.7	18.5
2 Blueberries, canned, as purchased.....	165	5.8	1	1	21
Cantaloupe.....	Half ordinary serving.....	243	8.6	0.72	11
2 Cranberries, as purchased, average.....	210	7.5	0.8	1.2	20.7
2 Grapes, as purchased, average.....	136	4.8	1.3	1.6	19.5
2 Oranges, as purchased, average.....	One very large one.....	270	9.4	1.3	0.2	18.7
2 Peaches, as purchased, average.....	Three ordinary ones.....	290	10	1.4	0.3	22.3

² Chemical Composition of American Food Stuffs, Atwater and Bryant, U. S. Dep't Agricult., Bull. No. 28.

Name of Food.	Portion containing 100 calories roughly described.		Actual amount of protein, fat, and carbohydrate in each portion of 100 calories.			
	Gm.	Oz.	Protein.	Fat.	Carbohydrates.	Gm.
FRUITS (FRESH OR COOKED)						
2 Pear.....	One large pear.....	173	5.4	0.8	44.3	
Raspberries.....	178	6.2	...	22.4	
2 Strawberries, as purchased, average.....	Two servings.....	260	9.1	1.5	19.2	
CAKE, PUDDING, PASTRY DESSERTS.						
2 Cake, chocolate layer.....	Half ordinary square.....	28	0.08	2.2	12.7	
2 Sponge cake.....	Small piece.....	25	0.89	2.7	16.5	
2 Gingerbread, as purchased.....	Half ordinary square piece.....	27	0.96	2.4	18	
Custard, milk.....	Ordinary cup.....	122	4.2	7.7	32	
Custard, tapioca.....	Two-thirds ordinary cup.....	69	2.4	2.1	18.4	
2 Doughnuts, as purchased.....	Half a doughnut.....	23	0.8	4.8	12.2	
2 Pie, apple, as purchased.....	One-third ordinary piece.....	38	1.3	3.7	16.3	
2 Pie, custard, as purchased.....	One-third ordinary piece.....	55	1.9	3.4	14.3	
2 Pie, cream, as purchased.....	One-fourth ordinary piece.....	30	1.1	3.4	15.3	
2 Pie, lemon, as purchased.....	One-third ordinary piece.....	38	1.3	3.8	14.2	
Pudding, apple sago.....	81	3	0.8	23.7	
Pudding, cream, rice.....	Very small serving.....	75	2.6	3.4	23.5	
Pudding, Indian meal.....	Half ordinary serving.....	56	2	2.6	15.4	
Pudding, apple, tapioca.....	Small serving.....	79	2.8	0.07	23	
Tapioca, cooked.....	Ordinary serving.....	108	3.8	3.4	30.4	

MISCELLANEOUS.

2 Honey, as purchased.....	Four teaspoonfuls.....	30	1	0.1	24.3
2 Molasses, cane.....	35	1.2	0.8	24
2 Marmalade (orange peel).....	28	1	0.2	0.02	24
2 Sugar, granulated.....	3 teaspoonfuls (1½ lumps) ..	24	0.8	24
2 Sugar, maple or syrup.....	Four teaspoonfuls.....	35	1.2	35
2 Almonds.....	Eight.....	15	0.5	3.1	8.2	2.6
2 Chestnuts, fresh, edible portion, as purchased, average.....	40	1.4	2.4	2.1	17
2 Peanuts.....	Thirteen double.....	18	0.62	4.6	8	4.4
2 Hickory nuts.....	13	0.47	2	8.7	1.5
2 Eggs, hens', boiled.....	One large egg.....	59	2.1	7.9	7.2
2 Eggs, hens', white.....	181	6.4	22
2 Eggs, hens', yolk.....	Two yolks.....	27	0.94	4.2	9
2 Soup, beef, as purchased.....	380	13	16.7	1.5	4.1
2 Soup, bean, as purchased.....	Very large plate.....	150	5.4	4.8	2.1	14
2 Soup, cream of celery, as purchased.....	Two plates.....	180	6.3	3.7	5	9
2 Consommé, as purchased.....	830	29	20.7	3.3

² Chemical Composition of American Food Stuffs, Atwater and Bryant, U. S. Dep't Agricult., Bull. No. 28.

APPENDIX

KARELL TREATMENT FOR ANASARCA

For First Five to Seven Days

8, 12 A. M.; 4, 8 P. M.: Milk, 200 c.c.
No other fluids.

Eighth Day

Milk as above.

10 A. M.: 1 soft-boiled egg.

6 P. M.: 2 pieces of dry toast.

Ninth Day

Milk as above.

10 A. M.: 1 soft-boiled egg and 2 pieces of dry toast.

6 P. M.: 1 egg and 2 pieces of dry toast.

Tenth Day

Milk as above.

12 NOON.: Chopped meat, rice boiled in milk, vegetables.

6 P. M.: 1 soft-boiled egg.

Eleventh to Twelfth Days

Same as tenth day.

No salt is used throughout diet. Salt-free toast and butter used. Small amount of cracked ice allowed with diet. All meat can often advantageously be omitted.

¹From the New York Hospital Diet Lists.

COMMENT ON THE KARELL DIET

The usefulness of the Karell treatment in cases of anasarca is sometimes extremely satisfactory, and may succeed in eliminating the edema in certain cases where the ordinary salt-poor diets fail. The limitation of the fluid in-take, together with the low food value and minimal salt content, seem to explain its action. Besides being helpful in some nephritic cases, it is often of great use in fluid collections due to a decompensated heart.

The lessened amount of fluid taken, of course, diminishes the work of the heart, and so favors restoration of its function.

GASTRIC DIET

Von Leube Treatment for Peptic Ulcer¹

For the first three days neither food nor water is to be taken, the mouth being kept moist by washes. If the patient is very thirsty, the Murphy drip may be used. In patients who are aged, debilitated, or desiccated by vomiting and insufficient nourishment, the first period of starvation is limited to twenty-four hours. On the second day 2 to 4 ounces of Celestine Vichy is given every two hours, and on the third day alternate these doses with albumin-water, so that liquids are taken every hour.

During this time hot applications, preferably by the electric pad, are applied constantly to the epigastrium. On the fourth day peptonized milk, 2 ounces, Celestine Vichy, 2 or 3 ounces, on alternate hours are begun, and these quantities are increased every day until 8 ounces of each are taken. The hot applications are still continued and the patient kept quietly in bed.

¹As modified by G. R. Lockwood.

The bowels are now kept open by low simple enemata. If there is any acidity, alkaline powders are given, but otherwise no medicine is taken. About the tenth day are added, in order, milk-toast, junket, and fine cereal, and then one may begin one of two lines of treatment: If the bowels are constipated and the breath offensive, start with Carlsbad water, giving a glassful, hot, before the first nourishment. If necessary, one-half such dose is to be given the last thing at night. If the bowels still remain constipated, reduce the temperature of the Carlsbad to tepid, giving practically a Markbrunnen instead of Sprudel. If the bowels are not so constipated, and if pain and acidity have been permanent symptoms, some rely on nitrate of silver, giving $\frac{3}{8}$ grain three times a day for three days, $\frac{4}{8}$ grain for three days, and $\frac{5}{8}$ grain for three days, and then repeat the cycle. Diarrhea may follow such dosage.

During the third week are added, in order, spaghetti and macaroni, creamed mashed potato, fine cereal, such as hominy or cream of wheat, and creamed fresh fish, such as creamed halibut. Any vegetable that is put through a sieve, such as pea purée, or thick purée soups, without meat stock, may be given. Farinaceous desserts, such as farina, tapioca, cornstarch, blanc-mange, rice pudding, and custard, may also be given.

The patient still remains in bed and continues the use of the hot applications. The hot applications are used regularly, except in cases with recent hemorrhage.

During the fourth week the patient begins to be up and about on the convalescent gastric diets.

INDEX

- ACHYLIA, 74
Acidosis, 64
Akoll biscuit, 87
Almond meal, 84
Almonds, 108, 117
Ambulatory treatment of chronic
 peptic ulcer, 53
Anasarca, Karell treatment for,
 119, 120
Anticonstipation diet, 70
Antifat diet, 77
Apple pie, 105, 116
 sago pudding, 116
 sauce, 115
Apples, 85, 107, 108, 115
Apricots, 107, 108, 115
Arthritis deformans, 72
Artichoke, 105
Asparagus, 105, 113
 cream of, soup, 109
Atwater and Bryant, 4, 98
Average chemical composition of
 American food stuffs, 98
- BACON, 99
Baked beans, 85
 custard, 90
Bananas, 85, 107, 115
Barker's gluten flour, A, 84, 87
 B, 84
 C, 84
Barley, 102
Bass, 100
Bean soup, 117
- Beans, baked, 113
 butter, 105
 lima, 114
 string, 113
Beechnuts, 108
Beef, 98, 109, 111, 112
 soup, 117
Beef-juice, 102
Beets, 84, 105
Biscuit, home-made, 104
 soda, 104
 whole wheat, 104
Black bean soup, 94
Blackberries, 107, 108, 115
Blackfish, 100
Blueberries, 108, 115
Bluefish, 101
Bouillon, 109
Bovaird (D., Jr.), 72
Brain, 98
Bran biscuits, raised, 93
 crackers, 93
Brazil nuts, 108
Bread, 113
 brown, 103
 corn, 103
 Graham, 104
 rye, 103
 wheat, 104
Buckwheat, 102
Bulletin, 26
 Department of Agriculture, 4, 98
Butter, 102, 112
Butterfish, 101

- Buttermilk, 21, 95, 102, 112
 Butternuts, 108
- CABBAGE, 105, 114
- Cake, 116
 bakers', 104
 chocolate layer, 104
 drop, 104
 frosted, 104
 gingerbread, 104
 lady-fingers, 104
 macaroons, 104
 sponge, 104
- Calcium metabolism, 73
- Caloric needs, 4
 values, 3
- Calves'-foot jelly, 102
- Candy, 105
- Canned beef, 98
- Cantaloupe, 115
- Carbohydrate equivalents, 65, 84
 tolerance, determination of, 64
- Cardiac disease, 119, 120
- Carrots, 84, 105, 114
- Casoid flour, 84, 87
- Cauliflower, 105, 114
- Celery, 105, 114
 cream of, soup, 109
- Cerealene, 102
- Cheese, 102, 112
 savory, 90
- Cherries, 107, 108
- Chestnuts, 109, 117
- Chicken, 100, 109, 111, 112
 sandwich, 110
 soup, 100
- Chittenden, 2
- Chlorids in urine, 26
- Chocolate, 109
- Chronic diarrhea, 74
- Citron, 108
- Clam chowder, 109
- Clams, 101
- Cocoa, 109
- Cocoanuts, 109
- Cod, 112
 fish, 100
- Coleman, W., 16, 20
- Comments on anticonstipation diet,
 70
 on convalescent diet, 12
 on diabetic diets, 63
 on fluid diet, 12
 on gastric diets, 50
 on milk diet, 12
 on obesity diet, 77
 on purin-free diets, 32
 on regular diet, 11
 on salt-poor diets, 24
 on Schmidt's intestinal test diet,
 83
 on soft diet, 12
 on test meals, 79
 on typhoid fever diets, 19
- Conner, L. A., 18
- Consommé, 109
 soup, 117
- Constipation, 70
- Corn, 85, 105, 114
 bread, 84, 113
 cream of, soup, 109
- Cornmeal, 102, 113
- Crackers, 113
 Boston, 104
 egg, 104
 Graham, 104
 oatmeal, 104
 oyster, 104
 saltines, 104
 soda, 104
 water, 104
- Cranberries, 91, 107, 115
- Cream, 84, 112
 of celery soup, 117


- Cream pie, 105, 116
 Creamed potatoes, 106
 Cucumbers, 105
 Currants, 107, 108
 Custard, 116
 baked, 84
 pie, 116
- DAIRY products, 102
 Dates, 108
 Delafield's mixture, 85, 97
 Diabetes, classification of cases, 64
 severe, diet in, 68
 treatment, 64
 Diabetic bread and biscuit, 87
 diet, 58
 brandy in, 59-62
 green days, 60
 lists, 61
 oatmeal days, 63
 prohibited articles, 62
 standard, with restricted pro-
 tein, 59
 strict, 58
 whisky in, 59-62
 white wine in, 59-62
 milk, 69
 analysis, 92
 Wright's, 91
 receipts, 87
 Diarrhea, 74
 chronic, 74
 Diet, chronic diarrhea, food to
 avoid, 76
 convalescent, 8
 gastric, No. 1, 48
 No. 2, 48
 gastric, 34
 in diabetes, 58
 in diarrhea, acute, 74
 chronic, 75
 in gastric hemorrhage, 34, 51
- Diet in gastric hyperacidity, 55
 ulcer, 34, 50, 52
 in gout, 28
 in obesity, 77
 caloric needs, 78
 methods of reduction, 78
 in peptic ulcer, 34, 50
 liquid, 11
 milk, 11
 modified milk, in typhoid, 16, 17,
 18
 New York hospital, 5, 8
 purin-free, liquid, 27
 soft, 27
 regular house diet, 5
 salt-poor, 22-24
 soft, 11
 typhoid, fluid, 14-17
 soft, 18
 Doughnuts, 104, 116
 Duck, 100
 Duodenal ulcer, 54
 chronic, 53
- EDEMA, 24
 Eels, 101
 Egg sandwich, 110
 Eggplant, 85, 106, 114
 Eggs, 101, 117
 boiled, 101
 whites, 101
 yolks, 101
 Eiweissmilch, 85, 95
 Epigastric pain, 53
- FARINA, 84
 Figs, 107, 108, 115
 Filberts, 109
 Fish, 100
 Fisher, C. I., 86
 Flatulence, 21
 Flour, 113

- Folin, 27
- Food values in anticonstipation diet, 70
- in convalescent diet, 10
- in gouty diet, 29
- in regular house diet, 8
- in soft diet, 11
- in typhoid fluid diet, 14-18
- soft diet, 19
- GASTRIC diagnosis, 80
- medication in peptic ulcer, 54
- test meals, 79
- ulcer, chronic, 53
- Gingerbread, 116
- Ginger snaps, 104
- Glucose, 52
- gluten biscuit, 87
- meal biscuit, 89
- Gouty diet, 28
- Granulated sugar, 105
- Grape-fruit, 84
- Grapes, 107, 115
- Green days, 60, 68
- peas, 85
- Greens, beet, 106
- Guinea-hen, 100
- Gum gluten, ground, 84
- Gumbo soup, 100
- HADDOCK, 101
- Halibut, 101, 112
- Ham, 99, 111, 112
- salad, 110
- Hash, 110
- Herring, 101
- Hickory nuts, 109
- Hominy, 84, 102, 113
- Honey, 105, 117
- Huckleberries, 107
- Huntley and Palmer biscuit, carbohydrate content of, 59
- Hyperacidity, 55
- Hyperglycemia, 67
- ICE-CREAM, 85, 90, 96, 105
- Indian meal mush, 84
- pudding, 105, 116
- Intestinal fermentation, 21
- test diet, 82
- JANEWAY, T. C., 48, 58, 59-61, 63, 87
- KARELL treatment for anasarca, 119, 120
- Keller's malt soup, 86, 96
- Kidney, 24, 98
- Kinnicutt, F. P., 14, 15
- Koumiss, 102
- LACTOBACILLIN, 95
- Lactose, 15
- ice-cream, 96
- jelly, 85, 94
- Lamb, 99
- chops, 111
- Lemon pie, 105, 116
- Lemons, 107
- Lenhartz diet, 34-50
- recapitulation, 49
- Lentils, 106, 114
- Lettuce, 106
- Lichi nuts, 109
- Lima beans, 114
- Liver, 98, 112
- Lockwood, G. R., 120
- Low calcium diet, 72
- MACARONI, 84, 103, 113
- Mackerel, 101, 112
- Maltose, 52
- Maple-sugar, 105
- Marmalade, 108, 117

- Mashed potatoes, 106
 Meat, boiled, in gout, 33
 stew, 109
 Men, average weight, 3, 97
 Milk, 112, 113
 condensed, 102
 skimmed, 102
 whey, 102
 whole, 84, 102
 Mince pie, 105
 Mincemeat, 110
 Miscellaneous receipts, 93
 Mock turtle soup, 109
 Molasses, 105, 117
 cookies, 104
 Mosenthal, H., 59
 Mulligatawney soup, 110
 Mushrooms, 106
 Muskmelons, 107
 Mutton, 99, 111, 112
- NAUSEA, 21, 85
 Nectarines, 107
 Nitrate of silver lavage in chronic
 peptic ulcer, 54
 Nitrogen, calcium, magnesium-
 fixed diets, 73
 Noodles, 103
- OATMEAL, 63, 84, 113
 boiled, 103
 days, 63
 griddle-cakes, 88
 gruel, 103
 muffins, 88
 popovers, 88
 water, 103
 Okra, 106
 Oleomargarine, 102
 Onions, 106, 114
 Oranges, 85, 107, 115
- Oxtail soup, 110
 Oysters, 101
- PARSNIPS, 85, 106, 114
 Pea soup, 93, 110
 Peaches, 85, 107, 115
 Peanut butter, 109
 Peanuts, 109, 117
 Pears, 85, 107, 108, 116
 Peas, 107, 114
 dried, 106
 green, 106
 Peptic ulcer, 120, 121
 Peptonized milk, 52
 Perch, 101
 Pheasant, 100
 Pickerel, 101
 Pie, 116
 Pigeon, 100
 Pike, 101
 Pineapple, 107
 Pogy, 101
 Pompano, 101
 Pork, 99, 111, 112
 Potato, 84, 106, 114
 biscuit, 87
 Poultry, 100
 Powdered sugar, 105
 Pretzels, 104
 Protein milk, 85, 95
 Protopuff, Nos. 1 and 2, 84, 87
 Prunes, 85, 108, 115
 Pudding, 116
 Pumpkins, 106, 107
 Pure gluten biscuits, 84
 Purin bodies in foods, 30
 Purin-free foods, 29
- QUAIL, 100
- RADISHES, 106
 Raisins, 108, 115

- Raspberries, 108, 116
 Rectal alimentation, 51
 Reubner, 3
 Rhubarb, 106
 Rice, 84, 113
 boiled, 103
 custard, 105
 flaked, 103
 pudding, 84, 105, 116
 Richter, 77
 Ripened milk, 95
 Rolled oats, 103
 Rolls, 113
 French, 104
 Vienna, 104
 Rye flour, 103
- SALMON, 101
 Salt elimination, 25, 26
 Salt-free diet, 22-24
 Salt-poor diet, No. 1, 22
 No. 2, 23
 No. 3, 24
 Salvia sticks, 84
 Sauerkraut, 106
 Sausage, 100
 bologna, 100
 frankfort, 100
 pork, 100
 Scallops, 101
 Schaefer, 20
 Schmidt's intestinal test diet, 82
 Shad, 101
 Shredded wheat, 84, 113
 Smelt, 101
 Soda cracker, 16
 Soja-bean meal, 84, 87
 biscuit, 88
 muffins, 89
 pancakes, 89
 pudding, 89
- Soup, 117
 Spaghetti, 103, 113
 Spinach, 106, 114
 Squab, 100
 Squash, 107
 pie, 105
 Standard diabetic diets, 58, 59
 portions, 86
 Steak, 98
 Stew, meat, 109
 Strawberries, 108, 116
 String beans, 105
 Succotash, 107, 115
 Sugar cookies, 104
 granulated, 117
 Sweet potatoes, 106, 114
 Sweetbreads, 98
- TABLE of carbohydrate equivalents,
 84
 Tapioca, 116
 pudding, 84, 105
 with apple, 105
 Test meals, gastric, 79
 chemistry of, 80
 Ewald, 79, 80
 Leube, 79, 81
 Reigel, 79, 81
 Tetany, 73
 Tomato soup, 110
 Tomatoes, 107, 114
 Tongue, 98
 Trout, 101
 Turkey, 100, 111
 Turnips, 85, 115
 Turtle soup, 110
 Tympanites, 21
- ULCER, peptic, 120, 121
 Uneda biscuit, carbohydrate con-
 tent of, 59

- VANDERBILT clinic diets, 27, 29, 70
Veal, 99, 111
Vegetable soup, 110
Vermicelli, 103
Vogel, K., 73
Vomiting, 21, 85
von Leube's treatment for peptic ulcer, 120, 121
von Noorden, 4, 28
- WALKER, H. J., 30
Walnut, 109
Watermelon, 85, 108
Wheat buns, 103
 cinnamon, 103
 cracked, 103
- Wheat, currant, 103
 entire, 103
 farina, 103
 flaked, 103
 gems, 103
 gluten, 103
 Graham, 103
 shredded, 103
White bread, 113
Widal, 24
Wolf, C. G. L., 72
Women, average weight, 3, 97
Wright's diabetic milk, 91
- ZWIEBACK, 104



Books for Nurses

PUBLISHED BY

W. B. SAUNDERS COMPANY

West Washington Square Philadelphia

London: 9, Henrietta Street, Covent Garden

Sanders' Nursing A NEW WORK

Miss Sanders' new book is undoubtedly the most complete and most practical work on nursing ever published. Everything about every subject with which the nurse should be familiar is detailed in a clean cut, definite way. There is no other nursing book so full of good, practical information—information you need.

Modern Methods in Nursing. By GEORGIANA J. SANDERS, formerly Superintendent of Nurses at Massachusetts General Hospital. 12mo of 881 pages, with 227 illustrations. Cloth, \$2.50 net.

Aikens' Home Nurse's Handbook

PRACTICAL

The point about this work is this: It tells you, and *shows* you just how to do those little—but none the less important—things entirely omitted from other nursing books, or at best only incidentally treated. The chapters on "Home Treatments" and "Every-Day Care of the Baby," stand out as particularly practical. Then the "Points to be Remembered"—terse, crisp reminders—is a feature of great value.

Home Nurse's Handbook. By CHARLOTTE A. AIKENS, formerly Director of the Sibley Memorial Hospital, Washington, D.C. 12mo of 276 pages, illustrated. Cloth, \$1.50 net

Stoney's Nursing

NEW (4th) EDITION

Of this work the *American Journal of Nursing* says: "It is the fullest and most complete and may well be recommended as being of great general usefulness. The best chapter is the one on observation of symptoms which is very thorough." There are directions how to *improvise* everything.

Practical Points in Nursing. By EMILY M. A. STONEY, formerly Superintendent of the Training School for Nurses in the Carney Hospital, South Boston, Mass. 12mo, 495 pages, illustrated. Cloth, \$1.75 net.

Stoney's Materia Medica

NEW (3d) EDITION

Stoney's Materia Medica was written by a head nurse who knows just what the nurse needs. *American Medicine* says it contains "all the information in regards to drugs that a nurse should possess."

Materia Medica for Nurses. By EMILY M. A. STONEY, formerly Superintendent of the Training School for Nurses in the Carney Hospital, South Boston, Mass. 12mo volume of 300 pages. Cloth, \$1.50 net.

Stoney's Surgical Technic

NEW (3d) EDITION

The first part of the book is devoted to Bacteriology and Antiseptics; the second part to Surgical Technic, Signs of Death, Bandaging, Care of Infants, etc.

Bacteriology and Surgical Technic for Nurses. By EMILY M. A. STONEY. Revised by FREDERIC R. GRIFFITH, M. D., New York. 12mo volume of 311 pages, fully illustrated. Cloth, \$1.50 net.

Goodnow's First-Year Nursing

ILLUSTRATED

Miss Goodnow's work deals *entirely* with the *practical* side of first-year nursing work. It is the *application* of text-book knowledge. It tells the nurse *how* to do those things she is called upon to do in her first year in the training school—the *actual ward work*.

First-Year Nursing. By MINNIE GOODNOW, R. N., formerly Superintendent of the Women's Hospital, Denver. 12mo of 328 pages, illustrated. Cloth, \$1.50 net.

Aikens' Hospital Management

This is just the work for hospital superintendents, training-school principals, physicians, and all who are actively interested in hospital administration. *The Medical Record* says: "Tells in concise form exactly what a hospital should do and how it should be run, from the scrubwoman up to its financing."

Hospital Management. Arranged and edited by CHARLOTTE A. AIKENS, formerly Director of Sibley Memorial Hospital, Washington, D. C. 12mo of 488 pages, illustrated. Cloth, \$3.00 net

Aikens' Primary Studies

NEW (2d) EDITION

Trained Nurse and Hospital Review says: "It is safe to say that any pupil who has mastered even the major portion of this work would be one of the best prepared first year pupils who ever stood for examination."

Primary Studies for Nurses. By CHARLOTTE A. AIKENS, formerly Director of Sibley Memorial Hospital, Washington, D. C. 12mo of 437 pages, illustrated. Cloth, \$1.75 net

Aikens' Training-School Methods and the Head Nurse

This work not only tells how to teach, but also what should be taught the nurse and *how much*. *The Medical Record* says: "This book is original, breezy and healthy."

Hospital Training-School Methods and the Head Nurse. By CHARLOTTE A. AIKENS, formerly Director of Sibley Memorial Hospital, Washington, D. C. 12mo of 267 pages. Cloth, \$1.50 net

Aikens' Clinical Studies

NEW (2d) EDITION

This work for second and third year students is written on the same lines as the author's successful work for primary students. *Dietetic and Hygienic Gazette* says there "is a large amount of practical information in this book."

Clinical Studies for Nurses. By CHARLOTTE A. AIKENS, formerly Director of Sibley Memorial Hospital, Washington, D. C. 12mo of 569 pages, illustrated. Cloth, \$2.00 net

Bolduan and Grund's Bacteriology JUST READY

The authors have laid particular emphasis on the immediate application of bacteriology to the art of nursing. It is an *applied bacteriology* in the truest sense. A study of all the ordinary modes of transmission of infection are included.

Applied Bacteriology for Nurses. By CHARLES F. BOLDUAN, M.D., Assistant to the General Medical Officer, and MARIE GRUND, M.D., Bacteriologist, Research Laboratory, Department of Health, City of New York. 12mo of 166 pages, illustrated. Cloth, \$1.25 net.

Fiske's The Body

A NEW IDEA

Trained Nurse and Hospital Review says "it is concise, well-written and well illustrated, and should meet with favor in schools for nurses and with the graduate nurse."

Structure and Functions of the Body. By ANNETTE FISKE, A. M., Graduate of the Waltham Training School for Nurses, Massachusetts. 12mo of 227 pages, illustrated. Cloth, \$1.25 net.

Beck's Reference Handbook JUST READY NEW (3d) EDITION

This book contains all the information that a nurse requires to carry out any directions given by the physician. The *Montreal Medical Journal* says it is "cleverly systematized and shows close observation of the sickroom and hospital regime."

A Reference Handbook for Nurses. By AMANDA K. BECK, Graduate of the Illinois Training School for Nurses, Chicago, Ill. 32mo volume of 244 pages. Bound in flexible leather, \$1.25 net.

Roberts' Bacteriology & Pathology JUST READY

This new work is practical in the strictest sense. Written specially for nurses, it confines itself to information that the nurse should know. All unessential matter is excluded. The style is concise and to the point, yet clear and plain. The text is illustrated throughout.

Bacteriology and Pathology for Nurses. By JAY G. ROBERTS, Ph. G., M. D., Oskaloosa, Iowa. 12mo of 206 pages, illustrated. \$1.25 net.

DeLee's Obstetrics for Nurses

THIRD
EDITION

Dr. DeLee's book really considers two subjects—obstetrics for nurses and actual obstetric nursing. *Trained Nurse and Hospital Review* says the "book abounds with practical suggestions, and they are given with such clearness that they cannot fail to leave their impress."

Obstetrics for Nurses. By JOSEPH B. DELEE, M. D., Professor of Obstetrics at the Northwestern University Medical School, Chicago. 12mo volume of 512 pages, fully illustrated. Cloth, \$2.50 net.

Davis' Obstetric & Gynecologic Nursing

JUST READY—NEW (4th) EDITION

The Trained Nurse and Hospital Review says: "This is one of the most practical and useful books ever presented to the nursing profession." The text is illustrated.

Obstetric and Gynecologic Nursing. By EDWARD P. DAVIS, M. D., Professor of Obstetrics in the Jefferson Medical College, Philadelphia. 12mo volume of 480 pages, illustrated. Buckram, \$1.75 net.

Macfarlane's Gynecology for Nurses

ILLUSTRATED

Dr. A. M. Seabrook, Woman's Hospital of Philadelphia, says: "It is a most admirable little book, covering in a concise but attractive way the subject from the nurse's standpoint. You certainly keep up to date in all these matters, and are to be complimented upon your progress and enterprise."

A Reference Handbook of Gynecology for Nurses. By CATHARINE MACFARLANE, M. D., Gynecologist to the Woman's Hospital of Philadelphia. 32mo of 150 pages, with 70 illustrations. Flexible leather, \$1.25 net.

McKenzie's Exercise in Education and Medicine

Exercise in Education and Medicine. By R. TAIT MCKENZIE, B.A., M.D., Professor of Physical Education, and Director of the Department, University of Pennsylvania. Octavo of 406 pages, with 346 illustrations. Cloth, \$3.50 net.

Manhattan Hospital Eye, Ear, Nose, and Throat Nursing

ILLUSTRATED

This is a practical book, prepared by surgeons who know the nurse's requirements in these specialties. *New York Medical Record* says: "Every side of the question has been fully taken into consideration."

Nursing in Diseases of the Eye, Ear, Nose and Throat. By the Committee on Nurses of the Manhattan Eye, Ear, and Throat Hospital: J. EDWARD GILES, M. D., Surgeon in Eye Department; ARTHUR B. DUEL, M. D., (chairman), Surgeon in Ear Department; HARMON SMITH, M. D., Surgeon in Throat Department. Assisted by JOHN R. SHANNON, M. D., Assistant Surgeon in Eye Department; and JOHN R. PAGE, M. D., Assistant Surgeon in Ear Department. With chapters by HERBERT B. WILCOX, M. D., Attending Physician to the Hospital; and Miss EUGENIA D. AVERS, Superintendent of Nurses. 12mo of 260 pages, illustrated. Cloth, \$1.50 net.

Friedenwald and Ruhrah's Dietetics for Nurses

NEW (2d) EDITION

This work has been prepared to meet the needs of the nurse, both in training school and after graduation. *American Journal of Nursing* says it "is exactly the book for which nurses and others have long and vainly sought."

Dietetics for Nurses. By JULIUS FRIEDENWALD, M. D., Professor of Diseases of the Stomach, and JOHN RUHRAH, M. D., Professor of Diseases of Children, College of Physicians and Surgeons, Baltimore. 12mo volume of 395 pages. Cloth, \$1.50 net.

Friedenwald & Ruhrah on Diet

THIRD EDITION

Diet in Health and Disease. By JULIUS FRIEDENWALD, M. D., and JOHN RUHRAH, M. D. Octavo volume of 764 pages. Cloth, \$4.00 net.

Galbraith's Personal Hygiene and Physical Training for Women

ILLUSTRATED

Personal Hygiene and Physical Training for Women. By ANNA M. GALBRAITH, M. D., Fellow New York Academy of Medicine. 12mo of 371 pages, illustrated. Cloth, \$2.00 net.

Galbraith's Four Epochs of Woman's Life

THE NEW (2d) EDITION

The Four Epochs of Woman's Life. By ANNA M. GALBRAITH, M. D. With an Introductory Note by JOHN H. MUSSER, M. D., University of Pennsylvania. 12mo of 247 pages. Cloth, \$1.50 net.

McCombs' Diseases of Children for Nurses

NEW (2d) EDITION

Dr. McCombs' experience in lecturing to nurses has enabled him to emphasize *just those points that nurses most need to know*. *National Hospital Record* says: "We have needed a good book on children's diseases and this volume admirably fills the want." The nurse's side has been written by head nurses, very valuable being the work of Miss Jennie Manly.

Diseases of Children for Nurses. By ROBERT S. MCCOMBS, M. D., Instructor of Nurses at the Children's Hospital of Philadelphia. 12mo of 470 pages, illustrated. Cloth, \$2.00 net

Wilson's Obstetric Nursing

NEW (2d) EDITION

In Dr. Wilson's work the entire subject is covered from the beginning of pregnancy, its course, signs, labor, its actual accomplishment, the puerperium and care of the infant. *American Journal of Obstetrics* says: "Every page emphasizes the nurse's relation to the case."

A Reference Handbook of Obstetric Nursing. By W. REYNOLDS WILSON, M.D., Visiting Physician to the Philadelphia Lying-in Charity. 32mo of 355 pages, illustrated. Flexible leather, \$1.25 net

American Pocket Dictionary

NEW (7th) EDITION

The *Trained Nurse and Hospital Review* says: "We have had many occasions to refer to this dictionary, and in every instance we have found the desired information."

American Pocket Medical Dictionary. Edited by W. A. NEWMAN DORLAND, A. M., M. D., Loyola University, Chicago. Flexible leather, gold edges, \$1.00 net; with patent thumb index, \$1.25 net

Lewis' Anatomy and Physiology

SECOND EDITION

Nurses Journal of Pacific Coast says "it is not in any sense rudimentary, but comprehensive in its treatment of the subjects." The low price makes this book particularly attractive.

Anatomy and Physiology for Nurses. By LEROY LEWIS, M.D., Lecturer on Anatomy and Physiology for Nurses, Lewis Hospital, Bay City, Mich. 12mo of 375 pages, 150 illustrations. Cloth, \$1.75 net

Boyd's State Registration for Nurses

State Registration for Nurses. By LOUIE CROFT BOYD, R. N., Graduate Colorado Training School for Nurses. Price, 50 cents net.

Paul's Materia Medica

NEW (2d) EDITION

A Text-Book of Materia Medica for Nurses. By GEORGE P. PAUL, M.D., Samaritan Hospital, Troy, N. Y. 12mo of 282 pages. Cloth, \$1.50 net.

Paul's Fever Nursing

NEW (2d) EDITION

Nursing in the Acute Infectious Fevers. By GEORGE P. PAUL, M.D. 12mo of 246 pages, illustrated. Cloth, \$1.00 net.

Hoxie & Laptad's Medicine for Nurses

JUST READY—NEW (2d) EDITION, REWRITTEN

Medicine for Nurses and Housemothers. By GEORGE HOWARD HOXIE, M.D., University of Kansas; and PEARL L. LAPTAD, 12mo of 351 pages, illustrated. Cloth, \$1.50 net.

Grafstrom's Mechano-therapy

SECOND EDITION

Mechano-therapy (Massage and Medical Gymnastics). By AXEL V. GRAFSTROM, B.Sc. M.D., 12mo, 200 pages. Cloth, \$1.25 net.

Nancrede's Anatomy

NEW (7th) EDITION

Essentials of Anatomy. CHARLES B. G. DENANCREDE, M.D., University of Michigan. 12mo, 400 pages, 180 illustrations. Cloth, \$1.00 net.

Morrow's Immediate Care of Injured

Immediate Care of the Injured. By ALBERT S. MORROW, M.D., New York City Home for Aged and Infirm. Octavo of 354 pages, with 242 illustrations. Cloth, \$2.50 net. **New (2d) Edition**

Register's Fever Nursing

A Text Book on Practical Fever Nursing. By EDWARD C. REGISTER, M.D., North Carolina Medical College. Octavo of 350 pages, illustrated. Cloth, \$2.50 net.

Pyle's Personal Hygiene

NEW (5th) EDITION

A Manual of Personal Hygiene. Edited by WALTER L. PYLE, M.D., Wills Eye Hospital, Philadelphia. 12mo, 515 pages, illus. \$1.50 net.

Morris' Materia Medica

NEW (7th) EDITION

Essentials of Materia Medica, Therapeutics, and Prescription Writing. By HENRY MORRIS, M.D. Revised by W. A. BASTEDO, M.D., Columbia University, N. Y. 12mo of 300 pages, illustrated. Cloth, \$1.00 net.

Griffith's Care of the Baby

NEW (5th) EDITION

The Care of the Baby. By J. P. CROZER GRIFFITH, M.D., University of Pennsylvania. 12mo of 455 pages, illustrated. Cloth, \$1.50 net.

