
Figure 1: Mean survival of *Listeria monocytogenes* alone and in the presence of lactic acid bacteria in 0% NaCl over a 10 day period (A=4°C, B=12°C)

-LM -LM + LAB







Figure 2: Mean survival of *Listeria monocytogenes* alone and in the presence of lactic acid bacteria in 7.9 and 13.2% NaCl over a 10 day period (A=4°C, B=12°C)







Figure 3: Mean survival of *Listeria innocua* alone and in the presence of lactic acid bacteria over a 10 day period (A=4°C, B=12°C)

— LI - 0% — LI+LAB - 0% — LI - 7.9% — LI+LAB - 7.9% — LI - 13.2% — LI+LAB - 13.2%







Figure 4: Mean survival of *Listeria innocua* and *Listeria monocytogenes* in brine without exposure to lactic acid bacteria over a 10 day period (A=4°C, B=12°C)

— LI - 0% — LM - 0% — LI - 7.9% — LM - 7.9% — LM - 13.2% — LM - 13.2%






Figure 5: Mean survival of *Listeria innocua* and *Listeria monocytogenes* in brine with exposure to lactic acid bacteria over a 10 day period (A=4°C, B=12°C)

— LI - 7.9% — LM - 7.9% — LM - 13.2% — LM - 13.2%

