

## **Acknowledgements**

I wish to express my most sincere appreciation to my major professor, Dr. Rod Youngman. He provided excellent opportunities, resources, advice, encouragement, assistance, and friendship during my years as his student. We had a successful relationship together that will certainly benefit me in life.

I would also like to thank my committee members, Dr. Timothy P. Mack, Dr. Loke T. Kok, Dr. Scott M. Salom, and Dr. Paul R. Peterson, whose doors were always open to provide me with advice and guidance. Thanks also to each of them for putting a good word in on my behalf on a number of occasions.

I am deeply indebted to Mr. Curtis A. Laub for his selfless sharing of time, hard work, and friendship. We spent countless hours together in cornfields, alfalfa fields, and softball fields. I could not have asked for a better work partner or teammate.

I thank all of the professors in the entomology department at Virginia Tech who made me work hard, but treated me with respect and kindness and encouraged my success. It was a joy to be part of that department for so long.

A special thanks to those who lent a helping hand to my work, including, Alexei Sharov for his technical genius, Tree Dellinger, Peter Warren, Stephanie Stockton, Julie Smith, Dick White, Tom Stanley, Chuck Shorter, Alec Lipscomb, Bernie Cline, Rob Tanzer, Sylvia Fernandez, Eric Grissell, and Bill Day.

And last, but not least, I want to thank the special people in my life who bring joy to my being, my wonderful parents, Al and Doris Kuhar, my siblings and their kids, and my strong, beautiful wife Stacey.

## Table of Contents

Introduction .....	1
<i>Chapter 1</i>	
Alfalfa weevil literature review.....	2
<i>Chapter 2</i>	
Alfalfa weevil pest status and incidence of selected natural enemies in Virginia.....	9
<i>Chapter 3</i>	
Population dynamics and mortality of the alfalfa weevil in Virginia.....	20
<i>Chapter 4</i>	
Phenology of the alfalfa weevil in Virginia.....	46
<i>Chapter 5</i>	
Summary .....	63
References .....	65

## List of tables

2.1.	Alfalfa weevil densities in the three major alfalfa-growing regions of Virginia; 1996-98	14
2.2.	Percentage of fields exceeding the economic threshold for alfalfa weevil in Virginia, 1996-98	15
2.3.	Parasitization levels of alfalfa weevil larvae in Virginia; 1996-98	16
2.4.	Levels of <i>Z. phytonomi</i> infection in alfalfa weevil larvae in Virginia; 1996-98	17
2.5.	Economic impact of alfalfa weevil in Virginia	18
3.1.	Alfalfa weevil population levels at three locations in Virginia, 1997-98	31
3.2.	Average monthly temperatures and alfalfa weevil oviposition degree-day (ODD) accumulations at three locations in Virginia, 1997-98	32
3.3.	Comparison of predicted alfalfa weevil egg numbers according to a temperature-based oviposition function and observed egg numbers at three Virginia locations, 1997-98	33
3.4.	Alfalfa weevil population levels at three locations in Virginia, 1998-99	34
3.5.	Average monthly temperatures and alfalfa weevil oviposition degree-day (ODD) accumulations at three locations in Virginia, 1998-99	35
3.6.	Comparison of predicted alfalfa weevil egg numbers according to a temperature-based oviposition function versus observed egg numbers at three Virginia locations, 1998-99	36
3.7.	<i>Beauveria bassiana</i> infection of alfalfa weevil adults at three locations in Virginia	37
3.8.	<i>Microctonus aethiopoides</i> parasitization of alfalfa weevil adults at three locations in Virginia	38
3.9.	<i>Zoophthora phytonomi</i> infection and <i>Bathyplectes</i> spp. parasitization of alfalfa weevil larvae at three locations in Virginia	39
3.10.	Partial composite lifetables of <i>H. postica</i> at Rustburg (a), Fairfield (b), and Blacksburg, Virginia (c).	40
3.11.	Linear regression of individual submortalities ( <i>k</i> -values) against total mortality ( <i>K</i> ) of <i>H. postica</i> at three locations in Virginia, 1997-99.	41
4.1.	Occurrence of 50% of the total larval population of alfalfa weevil at three locations in Virginia	52
4.2.	Alfalfa stem height at peak infestation of alfalfa weevil larvae in three locations of Virginia	53



## List of Figures

2.1.	Virginia counties surveyed for alfalfa weevil from 1996-98.	19
3.1.	Overwintering survival of alfalfa weevil adults in Virginia. Data points represent the mean ( $\pm$ SEM) of the proportion of adult population from 3 fields. Triangles represent Rustburg, diamonds represent Fairfield, and squares represent Blacksburg. Black represents the 1997-98 season and white represents 1998-99.	42
3.2.	Fluctuations in individual mortalities ( $k$ - values) with total mortality ( $K$ ) of <i>H. postica</i> over 6 field-years at Rustburg, Virginia; $k_1$ = eggs not laid, $k_2$ = egg mortality, $k_3$ = period-one larval mortality, $k_4$ = <i>Z. phytonomi</i> , $k_5$ = larval parasitism.	43
3.3.	Fluctuations in individual mortalities ( $k$ - values) with total mortality ( $K$ ) of <i>H. postica</i> over 6 field-years at Fairfield, Virginia; $k_1$ = eggs not laid, $k_2$ = egg mortality, $k_3$ = period-one larval mortality, $k_4$ = <i>Z. phytonomi</i> , $k_5$ = larval parasitism.	44
3.4.	Fluctuations in individual mortalities ( $k$ - values) with total mortality ( $K$ ) of <i>H. postica</i> over 6 field-years at Blacksburg, Virginia; $k_1$ = eggs not laid, $k_2$ = egg mortality, $k_3$ = period-one larval mortality, $k_4$ = <i>Z. phytonomi</i> , $k_5$ = larval parasitism.	45
4.1.	Population dynamics of alfalfa weevil eggs, period-one larvae (instars 1 and 2), and period-two larvae (instars 3 and 4) at three locations in Virginia; 1997-98. Data points represent the mean $\pm$ SEM of three fields.	55
4.2.	Population dynamics of alfalfa weevil eggs, period-one larvae (instars 1 and 2), and period-two larvae (instars 3 and 4) at three locations in Virginia; 1998-99. Data points represent the mean $\pm$ SEM of three fields.	56
4.3.	Phenological area graphs of alfalfa weevil populations at three locations of Virginia; 1997-98. Each graph represents a mean of 3 fields.	57
4.4.	Phenological area graphs of alfalfa weevil populations at three locations of Virginia; 1998-99. Each graph represents a mean of 3 fields.	58
4.5.	Seasonal occurrence of alfalfa weevil larvae over time. Data represent the mean $\pm$ SEM of three fields at each of three locations in 1998 (a) and (1999 (b); Blue = Rustburg, Green = Fairfield, Pink = Blacksburg. Colored tick marks on the x-axis refer to the date of 50% occurrence of the population.	59
4.6.	Spring growth of alfalfa at three Virginia locations in 1998 and 1999. Data points represent the mean $\pm$ SEM of three fields.	60
4.7.	Phenological synchrony of the parasitoid, <i>B. anurus</i> with alfalfa weevil larval populations at three locations in Virginia, 1998. Data points represent the mean $\pm$ SEM of three fields. Hatched areas indicate the activity period of <i>B. anurus</i> adults based on a degree-day range of 177 to 260 DD base 9°C.	61
4.8.	Phenological synchrony of the parasitoid, <i>B. anurus</i> with alfalfa weevil larval	

populations at three locations in Virginia, 1999. Data points represent the mean  $\pm$  SEM of three fields. Hatched areas indicate the activity period of *B. anurus* adults based on a degree-day range of 177 to 260 DD base 9°C.

62