

Table of Figures

Chapter 1: Polypseudorotaxanes and Polyrotaxanes

Figure 1	Various types of polypseudorotaxanes and polyrotaxanes	2
Figure 2	Polyrotaxane containing a blocking group in every structural unit of the polymer chain	7
Figure 3	Synthesis of the cyclodextrin-based polyrotaxane 4 incorporating bulky groups along its polymer backbone	8
Figure 4	Synthesis of a polyrotaxane consisting of β -CD and poly(ethylene glycol)-poly(propylene glycol) triblock copolymer	10
Figure 5	Synthesis of a main-chain polypseudorotaxane based on poly(ϵ -lysine) and α -CD	11
Figure 6	Preparation of α -CD-based molecular tube	13
Figure 7	Preparation of a poly(polyrotaxane) by photoreaction	16
Figure 8	Poly(styrene/crown ether rotaxane)s	18
Figure 9	Synthesis of a main-chain polyrotaxane from a difunctional monomer incorporating a bulky group	19
Figure 10	Synthesis of a polyester main chain polypseudorotaxane	20
Figure 11	Synthesis of a side-chain polyrotaxanes based on β -CD	24
Figure 12	Preparation of a side-chain polypseudorotaxane	25
Figure 13	Preparation of a 1 D polyrotaxane based on cucurbituril	27
Figure 14	Synthesis of the first conducting polymetallorotaxanes	28
Figure 15	Mechanism for the formation of the mechanically branched and network polymers with a side chain rotaxane structure	30
Figure 16	A polypseudorotaxane with cyclobis(paraquat-phenylene) as the cyclic component	31

Figure 17	The structure of <i>p-tert</i> -butylcalix[8]arene	32
Figure 18	Polypseudorotaxanes and polyrotaxanes with new topologies	41
Figure 19	Non-covalent block and graft copolymers with pseudorotaxane structure	42

Chapter 3: Host Folding in the Formation of a Taco Complex

Figure 1	Job plot: the stoichiometry of the complex between 4a and 2 in CD ₃ COCD ₃ solution using data for H ₁ of 4a	69
Figure 2	Electrospray mass spectrum of a solution of 4a and 2 in a mixture of acetonitrile and chloroform (4:1)	70
Figure 3	X-ray crystal structures of 4a (a), 2a (b), and 4a•2a (d), and a cartoon representation of 4a•2a (c)	72
Figure 4	X-ray crystal structure of 4b	73
Figure S1	Relationship between Δ and $1/[2]_0$ for the complexation between 1b and 2 in acetone- <i>d</i> ₆ at 22°C	74

Chapter 4: First Supramolecular Poly(taco complex)

Figure 1	Job plot: the stoichiometry of the complex between 3 and 4 in CD ₃ COCD ₃ solution using data for H ₂ of 4	79
Figure 2	¹ H NMR spectra (400 MHz, CD ₃ COCD ₃ , 22 °C) of (a) 1.00 mM 2 and 1.00 mM 4 , (b) 1.00 mM 3 and 1.00 mM 4 , (c) 1.00 mM 4	79
Figure 3	Ball-stick (a) and cartoon (b) representations of 3•4	80
Figure 4	Ball-stick (a) and cartoon (b) representations of 3•4 packing structure	81

Chapter 5: Bis(*m*-phenylene)-32-Crown-10 Based Cryptands, Powerful Hosts for Paraquat Derivatives

Figure 1	X-ray structures of 1b•2a (a) and 3a•2a (b)	89
Figure 2	Job plots showing the 1:1 stoichiometries of the complexes between 3b and 2a (a), between 3c and 2a (b), between 3d and 2a (c), and between 3e and 2a (d) in CD ₃ COCD ₃	92

Figure 3	X-ray structures of 2a (a), 3b (b), and 3b•2a (c)	95
Figure 4	Two views of the X-ray structure of 3c•2a•3c	97
Figure 5	X-ray structure of 3d•2a	99
Figure 6	Ball-stick (a) and cartoon (b) representations of the 3d•2a packing structure	100
Figure 7	X-ray structures of 3e	101
Figure 8	Electrospray mass spectrum of a solution of 3b and 2a in a mixture of acetonitrile and chloroform (4:1)	103
Figure 9	Fast-atom bombardment mass spectrum of an acetone solution of 2b and 3b (1:1 molar ratio)	104
Figure 10	Electrostatic potential maps of 2a (a), 3b (b), 3c (c), 3d (d), and 3e (e) at the AM1 level as determined with the CAChe program	105

Chapter 6: A New Cryptand/Paraquat [2]Pseudorotaxane

Figure 1	Job plot showing the 1:1 stoichiometry of the complex between 1 and 2a in CD ₃ COCD ₃ solution using data for H ₁ of 1	126
Figure 2	X-ray structure of 2a , as determined by X-ray crystallography	127
Figure 3	X-ray structure of 1•2a	127
Figure 4	X-ray structure of 3•2b	128

Chapter 7: First Pseudorotaxane-Like [3]Complexes Based on Cryptands and Paraquat: Self-Assembly and Crystal Structures

Figure 1	Compounds used in this study and the solid-state structure ² of 3b•1	135
Figure 2	Job plot: the stoichiometry of the complex between 3a and 1 in CD ₃ COCD ₃ solution using data for the aromatic hydrogen atom of 3a	137
Figure 3	Partial ¹ H NMR spectra (400 MHz, CD ₃ COCD ₃ , 298K) of (a) 1.00 mM 1 , (b) 1.00 mM 2c and 1.00 mM 1 , (c) 1.00 mM 2d and 1.00 mM 1 , (d) 1.00 mM 3a and 1.00 mM 1 , (e) 1.00 mM 3b and 1.00 mM 1	138
Figure 4	Solid state structures of 3b•1•3b (a and b) and 3a•1•3a (c and d) and cartoon representations of 3b•1•3b (e) and 3a•1•3a (f)	141

Chapter 8: First Cryptand-Based [3]Pseudorotaxane: Cooperative Complexation between a Cryptand and a Bisparaquat Guest

Figure 1	Job plot: the stoichiometry of the complex between cryptand 1 and ditopic guest 2 in CD ₃ COCD ₃ solution using data for H ₁	154
Figure 2	Mole ratio plot for 2 and 3 . The solvent is acetone- <i>d</i> ₆	154
Figure 3	X-ray structure of 1 ₂ • 2	155
Figure 4	Scatchard plots for complexation of bisparaquat guest 2 with (a) (top) crown ether 3 and (b) (bottom) cryptand 1 in CD ₃ COCD ₃ at 22 °C	156

Chapter 9: Remarkably Improved Complexation of a Bisparaquat by Formation of a Pseudocryptand-Based [3]Pseudorotaxane

Figure 1	Mole ratio plot showing 1:1 stoichiometry for the complexation between 4 and 2a , 22°C	163
Figure 2	Relationship between Δ for H ₁ of 2a and 1/[4] ₀ for the complexation between 4 and 2a in CD ₃ COCD ₃ , 22°C	163
Figure 3	The influence of added TEATFA on the fraction <i>p</i> of complexed paraquat units	164
Figure 4	Two views of the X-ray structure of 4 ₂ • 2b •2H ₂ O	166

Chapter 10: Water Assisted Formation of a Pseudorotaxane and Its Dimer Based on a Supramolecular Cryptand

Figure 1	Compounds used in this study and an orthogonal view ⁶ of the cradled barbell complex 2 • 3	172
Figure 2	Partial ¹ H NMR spectra (400 MHz, 2.5:1 acetone- <i>d</i> ₆ :chloroform- <i>d</i> , 22 °C) of (a) 1.11 mM undried 3 (5.00 mM water), (b) 1.11 mM undried 3 and 2.29 mM 2 (5.50 mM water), (c) 1.11 mM undried 3 and 2.29 mM 1 (5.50 mM water), (d) 1.11 mM dried 3 and 2.29 mM 1 (1.38 mM water), (e) 2.29 mM 1 (0.50 mM water) and (f) 1.11 mM dried 3 and 2 (1.50 mM water)	173
Figure 3	(a) ORTEP diagram of 1 • 3 , (b) its cartoon representation, and (c) the cartoon representation of the offset face-to-face and edge-to-face π -stacking interactions	175

Figure 4	(a) ORTEP diagram of the dimer of 1•3 , (b) its cartoon representation, and (c) the cartoon representation of edge-to-face π -stacking interactions	176
----------	--	-----

Chapter 11: Formation of Dimers of Inclusion Cryptand/Paraquat Complexes Driven by Dipole-Dipole and Face-to-Face π -Stacking Interactions

Figure 1	Job plot showing the 1:1 stoichiometry of the complex between 1 and 3 in CD ₃ COCD ₃ solution	184
Figure 2	Partial ¹ H NMR spectra (400 MHz, 2.5:1 acetone- <i>d</i> ₆ :chloroform- <i>d</i> , 22 °C) of (a) 1.11 mM undried 3 (5.00 mM water), (b) 1.11 mM undried 3 and 2.29 mM 2 (5.50 mM water), (c) 1.11 mM undried 3 and 2.29 mM 1 (5.50 mM water), (d) 1.11 mM dried 3 and 2.29 mM 1 (1.38 mM water), (e) 2.29 mM 1 (0.50 mM water) and (f) 1.11 mM dried 3 and 2 (1.50 mM water)	185
Figure 3	X-ray structure of 1•3	186
Figure 4	Two views of the dimer structure (1•3) ₂	187

Chapter 12: An Inclusion Complex of Diquat in the Concave Cavity Provided by Two Dibenzo-24-Crown-8 Hosts

Figure 1	Job plot showing the 1:1 stoichiometry of the complex between 1 and 2 in CD ₃ COCD ₃ solution using data for H ₁ of 1	197
Figure 2	Electrospray mass spectrum of a solution of 1 and 2 in a mixture of acetonitrile and chloroform (4:1)	198
Figure 3	Two views of the solid-state structure of 2 as determined by X-ray crystallography	199
Figure 4	Stick (a) and space-filling (b) representations of the X-ray structure of 1 ₂ • 2 and stick (c) and space-filling (d) representations of a part of it showing the concave cavity provided by the two hosts	200

Chapter 13: Bis(*m*-phenylene)-32-crown-10/Monopyridinium [2]Pseudorotaxanes

Figure 1	Partial proton NMR spectra (400 MHz, 1:1 CDCl ₃ :CD ₃ COCD ₃ , 22 °C) of monopyridinium salt 2a (a, bottom), crown ether 1 (b, middle), and 30.0 mM 1 and 2a (c, top)	207
----------	--	-----

Figure 2	Job plot showing the 1:1 stoichiometry of the complex between crown ether 1 and monopyridinium 2a in 1:1 CDCl ₃ :CD ₃ COCD ₃	208
Figure 3	Two views of the X-ray structure of 1•2a	210
Figure S1	Job plot showing the 1:1 stoichiometry of the complex between crown ether 1 and monopyridinium 2b in 1:1 acetone- <i>d</i> ₆ :chloroform- <i>d</i>	212
Figure S2	Relationship between Δ and 1/[2a] ₀ for the complexation between 1 and 2a in 1:1 acetone- <i>d</i> ₆ :chloroform- <i>d</i> at 22°C	213
Figure S3	Relationship between Δ and 1/[2b] ₀ for the complexation between 1 and 2b in 1:1 acetone- <i>d</i> ₆ :chloroform- <i>d</i> at 22°C	213

Chapter 14: [2]Pseudorotaxanes Based on the Cryptand/Monopyridinium Recognition Motif

Figure 1	X-ray structures of 1a•2•1a•2H₂O (a) and 1b•2•1b (b)	220
Figure 2	Job plot showing the 1:1 stoichiometry of the complex between 1a and 3a in [D ₆]acetone	221
Figure 3	Partial proton NMR spectra (400 MHz, 1:1 CDCl ₃ :CD ₃ COCD ₃ , 22 °C) of monopyridinium salt 3a (a, bottom), cryptand 1a (b, middle), and 3.00 mM 1a and 3a (c, top)	223
Figure 4	Electrospray mass spectrum of a solution of 1a and 3a in a mixture of acetonitrile and chloroform (4:1)	226
Figure 5	Ball-stick view of one unique host-guest complex (the other is similar) in the X-ray structure of 1a•3a	228
Figure 6	Ball-stick view of the X-ray structure of 1c•3a	229
Figure 7	Ball-stick view of the X-ray structures of cryptand 1b (a) and [2]pseudorotaxane 1b•3a (b)	231
Figure 8	Ball-stick view of the X-ray structures of dimorphic forms of 1d•3a	234
Figure 9	Ball-stick view of [2]pseudorotaxane 1e•3a	236
Figure 10	Ball-stick view of [2]pseudorotaxane 1f•3a	237

Chapter 15: [3]Pseudorotaxanes Based on the Cryptand/Monopyridinium Recognition Motif

Figure 1	Compounds used in this study and the X-ray structure ³ of 1 ₂ • 4 with PF ₆ counterions removed	257
Figure 2	Partial proton NMR spectra (400 MHz, acetone- <i>d</i> ₆ , 22 °C) of bispyridinium salt 2a (a, bottom), cryptand 1 (b, middle), and 6.00 mM 1 and 3.00 mM 2a (c, top)	258
Figure 3	Mole ratio plot for 1 and 2a , indicating 2:1 stoichiometry. The solvent is acetone- <i>d</i> ₆	259
Figure 4	Scatchard plot for the complexation of cryptand host 1 with bispyridinium guest 2a in acetone- <i>d</i> ₆ at 22 °C	260
Figure 5	Ball-stick (a, top) and space-filling (b, bottom) views of the X-ray structure of 1 ₂ • 2a	263
Figure 6	Ball-stick (a, top) and space-filling (b, bottom) views of the X-ray structure of 1 ₂ • 2b	264

Chapter 16: Synthesis of a Symmetric Cylindrical Bis(crown ether) Host and Its Complexation with Paraquat

Figure 1	Two views of the X-ray structure of 1 . Two solvent molecules and hydrogens have been omitted for clarity	275
Figure 2	Partial ¹ H NMR spectra (400 MHz, acetone- <i>d</i> ₆ , 22 °C) of (a, top) cylindrical bis(crown ether) 1 , (b, middle) paraquat 2 , and (c, bottom) a solution of 0.670 mM 1 and 1.33 mM 2	276
Figure 3	Mole ratio plot for the complexation between 1 and 2 in acetone- <i>d</i> ₆ :chloroform- <i>d</i> (5:1)	276
Figure 4	Scatchard plots for the complexation of cylindrical bis(crown ether) host 1 with paraquat 2 in 5:1 CD ₃ COCD ₃ :CDCl ₃ (a, top) and CD ₃ COCD ₃ (b, bottom) at 22 °C	278
Figure 5	Electrospray mass spectrum of a solution of 1 and 2 (molar ratio 1:2) in a mixture of acetonitrile and chloroform (4:1)	279
Figure 6	Two views of the [3]pseudorotaxane 1 • 2 ₂ •(CH ₃ CN) ₃	281

Chapter 17: Slow-Exchange C₃-Symmetric Cryptand/Trispyridinium Inclusion Complexes: A New Type of Threaded Structure

Figure 1	Compounds used in this study and X-ray crystal structure of 2a	293
Figure 2	Partial proton NMR spectra (400 MHz, CD ₃ COCD ₃ , 22 °C) of cryptand 1 (a, bottom), trispyridinium 2a (b, middle), and 10.0 mM 1 and 2a (c, top)	295
Figure 3	Two views of the minimized-energy structure of 1•2a at the AM1 level with three PF ₆ counterions removed	296
Figure S1	¹ H NMR Spectrum (400 MHz, CD ₃ COCD ₃ , 22 °C) of 2a	298
Figure S2	¹ H NMR Spectrum (400 MHz, CD ₃ COCD ₃ , 22 °C) of 2b	298
Figure S3	¹ H NMR Spectrum (400 MHz, CD ₃ COCD ₃ , 22 °C) of 2c	299

Chapter 18: A Supramolecular Poly[3]pseudorotaxane from Self-Assembly of a Homoditopic Cylindrical Bis(crown ether) Host and a Bisparaquat Derivative

Figure 1	Compounds used in this study and a space-filling diagram ⁶ of the [3]pseudorotaxane 1•2₂	310
Figure 2	Partial proton NMR spectra (400 MHz, 1:1 acetonitrile- <i>d</i> ₃ :chloroform- <i>d</i> , RT) of 3 (a) and the mixture of 1 and 3 (molar ratio 1:1) at different concentrations: (b) 0.438, (c) 1.90, (d) 2.22, (e) 2.67, (f) 3.33, (g) 4.44, and (h) 6.67 mM	312
Figure 3	The partial MALDI-TOF mass spectrum of the supramolecular poly[3]pseudorotaxane 4 formed from 1 and 3	314
Figure 4	Reduced viscosity as a function of bisparaquat concentration (solutions at RT): (a) 1 and 3 (molar ratio 1:1) in 1:1 CH ₃ CN:CHCl ₃ , (b) 1 and 3 (molar ratio 1:1) in a 1:1 CH ₃ CN:CHCl ₃ solution of 0.0500 M tetrabutylammonium hexafluorophosphate, (c) 5 and 3 (molar ratio 2:1) in 1:1 CH ₃ CN:CHCl ₃	315

Chapter 19: A Supramolecular Triarm Star Polymer from a Homotritopic Tris(Crown Ether) Host and a Complementary Monotopic Paraquat-Terminated Polystyrene Guest by a Supramolecular Coupling Method

Figure 1	(a) Mole ratio plot for 3 and 2 , indicating 1:3 stoichiometry. The solvent is CDCl ₃ . [3] _o = 0.100 mM. (b) Mole ratio plot for 3 and 5 , indicating 1:3
----------	--

	stoichiometry. The solvent is 2:1 CD ₃ COCD ₃ :CDCl ₃ . [3] ₀ = 0.100 mM	325
Figure 2	Reduced viscosity as a function of concentration (solutions in chloroform at RT): (a) polymeric guest 2 , (b) monotopic host 4 and polymeric guest 2 (molar ratio 1:1), (c) ditopic host 7 and polymeric guest 2 (molar ratio 1:2), and (d) tritopic host 3 and polymeric guest 2 (molar ratio 1:3)	327
Figure 3	Scatchard plots for the complexations of homotritopic host 3 with (a) monotopic paraquat-terminated polystyrene guest 2 in CDCl ₃ and (b) model monotopic paraquat guest 5 in 2:1 CD ₃ COCD ₃ :CDCl ₃ at 22 °C	328
Figure S1	¹ H NMR Spectrum (400 MHz, CD ₃ COCD ₃ , 22 °C) of 1	331
Figure S2	¹ H NMR Spectrum (400 MHz, CDCl ₃ , 22 °C) of 2	334
Figure S3	¹ H NMR Spectrum (400 MHz, CDCl ₃ , 22 °C) of 7	335
Figure S4	¹ H NMR Spectrum (400 MHz, CDCl ₃ , 22 °C) of 3	336

Chapter 20: Formation of a Supramolecular Hyperbranched Polymer from Self-Organization of an AB₂ Monomer Containing a Crown Ether and Two Paraquat Moieties

Figure 1	Partial proton NMR spectra (400 MHz, CD ₃ CN, 22 °C) of 1 at different concentrations: (a) 0.333, (b) 1.00, (c) 10.0, (d) 20.0, (e) 30.0, (f) 40.0, (g) 50.0, (h) 60.0, (i) 70.0, (j) 80.0, (k) 90.0, and (l) 100 mM	342
Figure 2	The relationship between the chemical shift of H ₂ on 1 and the initial concentration of 1	343
Figure 3	The partial MALDI-TOF mass spectrum of the supramolecular hyperbranched polymer 4	344
Figure 4	Reduced viscosity as a function of crown concentration (solutions in acetonitrile at 22 °C): (a) 1 , (b) 2b and 5 (molar ratio 1:2)	345
Figure S1	¹ H NMR Spectrum (400 MHz, CD ₃ COCD ₃ , 22 °C) of 1	347

Chapter 21: Conclusions and Future Work

Figure 1	Dependence of <i>n</i> on log <i>K</i> _a for [H] = 1 M	357
Figure 2	The relationship between molecular weight (<i>M</i> _n) of macromolecular building block and lowest association constant to reach 95% complexation	

for a system with one end-functional group per macromolecular building
block in a neat equimolar host-guest mixture of density 1.0 g/mole

357