

Table of Tables

Chapter 5: Bis(*m*-phenylene)-32-Crown-10 Based Cryptands, Powerful Hosts for Paraquat Derivatives

Table 1	Association constants ^a and thermodynamic parameters for complexes of paraquat derivatives (2a and 2b) with crown ethers (1c and 1d) and a cryptand (3a)	88
Table 2	Association constants in acetone- <i>d</i> ₆ for complexes of paraquat 2a with crown ether 1a and different cryptands at 22 °C	93

Chapter 7: First Pseudorotaxane-Like [3]Complexes Based on Cryptands and Paraquat: Self-Assembly and Crystal Structures

Table 1	Hydrogen-bond parameters for 3b•1•3b	142
Table 2	Selected distances and angles for 3b•1•3b and 3a•1•3a	142
Table 3	Hydrogen-bond parameters for 3a•1•3a	142

Chapter 14: [2]Pseudorotaxanes Based on the Cryptand/Monopyridinium Recognition Motif

Table 1	Association constants (K_a) values (M^{-1}) in different solvents for 1a /monopyridinium complexes at 22 °C	224
Table 2	Association constants (K_a) values (M^{-1}) for cryptand/ 3a complexes at 22 °C	224
Table 3	Observed mass/charge ratios for 1:1 cryptand/Monopyridinium complexes in CH ₃ CN/CHCl ₃ (4:1) from electrospray ionization mass spectrometric characterization	225

Chapter 18: A Supramolecular Poly[3]pseudorotaxane from Self-Assembly of a Homoditopic Cylindrical Bis(crown ether) Host and a Bisparaquat Derivative

Table 1	Δ , p , and n at different initial concentrations of 1 and 3	312
---------	---	-----

Chapter 20: Formation of a Supramolecular Hyperbranched Polymer from Self-Organization of an AB₂ Monomer Containing a Crown Ether and Two Paraquat Moieties

Table 1 Calculated values of p and n at different initial concentrations of **1** based on K_a
= 380 and 100 M⁻¹

344