

Table of Contents		Page #
Abstract		ii
Acknowledgements		iv
List of Figures		ix
List of Tables		x
Chapter I	Introduction and review of literature	1
	<ul style="list-style-type: none"> • Need for new herbicide chemistries in corn • Other corn herbicides • Transgenic corn hybrids • Carotenoid biosynthesis inhibitors • RPA 201772 (or EXP 31130A) • SC 0051 (or ICIA 0051) • Mesotrione • Literature Cited 	1 3 6 8 11 15 18 21
Chapter II	Mesotrione, acetochlor, and atrazine for weed management in corn (<i>Zea mays</i>)	33
	<ul style="list-style-type: none"> • Abstract • Introduction • Methods and Materials • Results and Discussion • Acknowledgements • Literature cited 	33 34 36 39 44 45
Chapter III	Mesotrione combinations in no-till corn (<i>Zea mays</i>)	54
	<ul style="list-style-type: none"> • Abstract • Introduction • Methods and Materials • Results and Discussion • Acknowledgements • Literature Cited 	54 55 57 59 63 64
Chapter IV	Mesotrione combinations in glyphosate-resistant corn (<i>Zea</i>	70

	<i>mays</i>)	
	• Abstract	70
	• Introduction	71
	• Methods and Materials	72
	• Results and Discussion	75
	• Acknowledgements	79
	• Literature Cited	80
Chapter V	Mesotrione, imazethapyr, and imazethapyr plus imazapyr combinations in imidazolinone-resistant corn (<i>Zea mays</i>)	88
	• Abstract	88
	• Introduction	89
	• Methods and Materials	91
	• Results and Discussion	94
	• Acknowledgements	96
	• Literature Cited	97
Chapter VI	Mesotrione combinations for postemergence control of horsenettle (<i>Solanum carolinense</i>) in corn (<i>Zea mays</i>)	105
	• Abstract	105
	• Introduction	106
	• Methods and Materials	108
	• Results and Discussion	112
	• Acknowledgements	117
	• Literature Cited	118
Chapter VII	Efficacy, absorption, translocation, and metabolism of mesotrione in Canada thistle (<i>Cirsium arvense</i>)	125
	• Abstract	125
	• Introduction	126
	• Methods and Materials	129
	• Results and Discussion	134
	• Acknowledgements	139

	• Literature Cited	140
Chapter VIII	Summary	153
Vita		155

List of Figures

Chapter	Figure	Title	Page #
VII	7.1	Movement of ^{14}C mesotrione with COC plus UAN in bolting stage Canada thistle (left) and rosette stage Canada thistle (above) at 72 hours after treatment.	151
	7.2	The interrelationship between the HPPD inhibitor mesotrione and the PSII inhibitor atrazine.	152

List of Tables

Chapter	Table	Title	Page #
II	2.1	Rainfall at 7-day intervals for 28 days after preemergence applications in 1999, 2000, and 2001.	48
	2.2	Corn injury, weed control, and corn yield from the pre-package mix of mesotrione plus acetochlor in 1999 and 2000.	49
	2.3	Common ragweed control with preemergence and postemergence applications of mesotrione treatments in 1999, 2000, and 2001.	50
	2.4	Smooth pigweed and common lambsquarters control with preemergence and postemergence applications of mesotrione treatments in 1999 and 2001.	51
	2.5	Morningglory species and giant foxtail control with preemergence and postemergence applications of mesotrione treatments in 1999, 2000, and 2001.	52
	2.6	Corn injury and yields with preemergence and postemergence applications of mesotrione treatments in 1999, 2000 and 2001.	53
III	3.1	Cutleaf eveningprimrose, smooth pigweed, and common lambsquarters control with preemergence and postemergence mesotrione in no-till corn in 1999, 2000, and 2001.	67
	3.2	Common ragweed and annual grass control and corn yields with preemergence and postemergence mesotrione in no-till corn in 1999, 2000, and 2001.	68
	3.3	Evaluation of the mesotrione plus acetochlor pre-package mix applied preemergence for the control of field pansy, common ragweed, and ivyleaf morningglory and corn yields in no-till corn in 1999 and 2001.	69
IV	4.1	Glyphosate-resistant corn injury from postemergence mesotrione and mesotrione plus glyphosate treatments in 1999, 2000, and 2001.	84
	4.2	Common ragweed, common lambsquarters, and smooth pigweed control from postemergence mesotrione and mesotrione plus	84

	glyphosate combinations in 1999, 2000, and 2001.	
	4.3 Morningglory species and giant foxtail control and corn yields from postemergence mesotrione and mesotrione plus glyphosate combinations in 1999, 2000, and 2001.	86
	4.4 Injury, height, and shoot biomass reductions of glyphosate-resistant corn treated postemergence with mesotrione or mesotrione plus glyphosate in the greenhouse.	87
V	5.1 Common ragweed and common lambsquarters control from postemergence applications of mesotrione alone or in combinations with imazethapyr or imazethapyr plus imazapyr in 2000 and 2001.	101
	5.2 Control of morningglory species and giant foxtail from postemergence applications of mesotrione alone and in combinations with imazethapyr and imazethapyr plus imazapyr in 1999 and 2001.	102
	5.3 Injury and yields of corn treated postemergence with mesotrione alone or in combinations with imazethapyr or imazethapyr plus imazapyr in 1999, 2000, and 2001.	103
	5.4 Injury, height, and biomass reductions of corn treated with postemergence mesotrione alone or in combinations with imazethapyr or imazethapyr plus imazapyr in the greenhouse.	104
VI	6.1 Horsenettle control from mesotrione alone and combinations of mesotrione with primisulfuron, dicamba, and 2,4-D 5 weeks after treatment at the Kranz Farm in 1999 and Mapp Farm in 2000 and 2001.	121
	6.2 Horsenettle population counts, percent population reductions, and percent biomass reductions from mesotrione and mesotrione combinations with primisulfuron, dicamba, and 2,4-D at the Kranz Farm in 1999 and the Mapp Farm in 2000 and 2001.	122
	6.3 Horsenettle control, and horsenettle population and biomass reductions data from mesotrione and standard herbicide treatments	123

	pooled over the Mapp and Lassiter Farms in 2000 and 2001.	
	6.4 Mesotrione combinations with primisulfuron, dicamba, and 2,4-D for horsenettle control in the greenhouse.	124
VII	7.1 Mesotrione alone and in combinations with atrazine and clopyralid compared to standard herbicide treatments for POST control of Canada thistle in 2000.	144
	7.2 Postemergence mesotrione plus atrazine combinations for control and reductions of Canada thistle shoot regrowth heights and biomass in the greenhouse.	145
	7.3 Postemergence mesotrione and atrazine combinations for height and biomass reductions of Canada thistle in the greenhouse.	146
	7.4 Influence of atrazine, adjuvant, and growth stage on absorption of foliar applied ¹⁴ C mesotrione in Canada thistle.	147
	7.5 Influence of atrazine, adjuvant, and growth stage on absorption and translocation of foliar applied ¹⁴ C mesotrione in Canada thistle at 72 hours after treatment.	148
	7.6 Influence of atrazine, adjuvant, and growth stage on metabolism of foliar applied ¹⁴ C mesotrione in the treated leaf and the foliage above the treated leaf of Canada thistle at 72 hours after treatment.	149
	7.7 Influence of atrazine, adjuvant, and growth stage on metabolism of foliar applied ¹⁴ C mesotrione in foliage below the treated leaf and the roots of Canada thistle at 72 hours after treatment.	150