

LIST OF TABLES

3. EXPERIMENTAL

Table 1 Molecular Characteristics of Ethylene/1-Butene (EB), Ethylene/1-Pentene (EP) and Ethylene/1-Hexene (EH) copolymers.....	80
---	----

4. MATERIALS CHARACTERIZATION

Table 1 Peak assignments for the determination of branch content from the NMR spectra of ethylene/ α -olefin copolymers.....	101
Table 2 The integrated intensities of the significant bands in the ^{13}C NMR spectra of the ethylene copolymers. The mole percent branches and the number of branches per 1000 carbon atoms are also included	103
Table 3 Triad sequence distributions for the ethylene/1-butene copolymers.....	107
Table 4 Expected values for the random parameter, χ_R for different types of copolymers (Adopted from Ref. 12).....	108
Table 5 Branch concentrations of the ethylene/1-butene copolymers from Randall's method. Their random parameters are also included	108
Table 6 Triad sequence distributions for the ethylene/1-hexene copolymers	111
Table 7 Branch concentrations of the ethylene/1-hexene copolymers from Randall's method. Their random parameters are also included	111
Table 8 Bulk densities of ethylene/1-butene, ethylene/1-pentene and ethylene/1-hexene copolymers measured at 23°C (from L. Richardson)	113

5. DIFFERENTIAL SCANNING CALORIMETRY

Table 1 The kinetic parameters, A , B , n and k and the degrees of crystallinity at -20 and 20°C for the ethylene/ α -olefin copolymers.....	144
Table 2 Avrami exponents and associated crystal geometries for different crystallization mechanisms (Adopted from Ref. 32).....	149
Table 3 The parameters, A , B , n and k and the degrees of crystallinity at T_x for EB-912 crystallized at various temperatures (T_x) from 20 to 100°C.....	160

7. FOURIER TRANSFORM INFRARED SPECTROSCOPY

Table 1 Crystallinities for EB-912 (with ~ 5 mole % 1-butene) during (a) Heating of a quenched sample (b) Slow Cooling (c) Heating of a slowly cooled sample (A rate of 1.5°C/min was used for all three processes)	213
Table 2 Crystallinities for EB-894 (with ~ 9 mole % 1-butene) during (a) Heating of a quenched sample (b) Slow Cooling (c) Heating of a slowly cooled sample (A rate of 1.5°C/min was used for all three processes)	214
Table 3 Crystallinities for EB-883 (with ~ 13 mole % 1-butene) during (a) Heating of a quenched sample (b) Slow Cooling (c) Heating of a slowly cooled sample (A rate of 1.5°C/min was used for all three processes).....	215
Table 4 Ratios of the integrated intensity of the crystalline bands at 730 and 720 cm^{-1} in the infrared spectrum of EB-912 (~ 5 mole % 1-butene).....	225
Table 5 Ratios of the integrated intensity of the crystalline bands at 730 and 720 cm^{-1} in the infrared spectrum of EB-894 (~ 9 mole % 1-butene).....	226
Table 6 Ratios of the integrated intensity of the crystalline bands at 730 and 720 cm^{-1} in the infrared spectrum of EB-883 (~ 13 mole % 1-butene).....	227
Table 7 Relative contribution of the second crystal structure to the total crystalline fraction for the quenched ethylene/1-butene copolymers	232