

## LIST OF TABLES

<b>Table 1.1</b> Critical points of some common fluids .....	2
<b>Table 1.2</b> Research Program Outline .....	13
<b>Table 2.1</b> Polymer solutions in binary fluid mixtures .....	17
<b>Table 3.1</b> Summary of available viscometers .....	51
<b>Table 3.2</b> Literature data for polymer solution viscosity at high pressures. (Polymer solution with solvent as main component) .....	53
<b>Table 3.3</b> Literature data for polymer solution viscosity at high pressures. (Polymer solution with solute as main component) .....	54
<b>Table 5.1</b> Density ( $\rho$ ), viscosity ( $\eta$ ), excess volume ( $V^E$ ), excess viscosity ( $\Delta\eta$ ), excess Gibbs free energy ( $G_V^E$ ) of viscous flow, and Grunberg-Nissan interaction parameter ( $d_{12}$ ) of acetone + carbon dioxide mixtures. (Mixture compositions are based on mass) .....	100
<b>Table 5.2</b> Parameters for viscosity correlation with pressure (Mixture compositions are based on mass) (Equation 5.2) .....	103
<b>Table 5.3</b> Parameters for viscosity correlation with temperature (Mixture compositions are based on mass) (Equation 5.3) .....	104
<b>Table 5.4</b> Parameters for viscosity correlation with density (Mixture compositions are based on mass) (Equation 5.4) .....	105
<b>Table 6.1</b> Density and viscosity data for acetone. ....	140
<b>Table 6.2</b> Density and viscosity data for PMMA (15 K) solutions in acetone at different concentrations (c) .....	141
<b>Table 6.3</b> Density and viscosity data for PMMA (540 K) solutions in acetone at different concentrations (c) .....	143
<b>Table 6.4</b> Density and viscosity data for 5 wt % solutions of PMMA (540 K) in acetone + CO <sub>2</sub> mixtures .....	146
<b>Table 6.5</b> Activation volumes and activation energies for PMMA solutions in acetone and acetone + CO <sub>2</sub> . ....	148

<b>Table 6.6</b> Density correlation parameters for viscosity using Doolittle equation. .....	149
<b>Table 7.1.</b> Density and viscosity data for 5 wt % PCL solution in acetone. ....	181
<b>Table 7.2</b> Density and viscosity data for 5 wt % PCL (14 K) solutions in acetone + CO <sub>2</sub> at different CO <sub>2</sub> concentrations (c) .....	182
<b>Table 7.3</b> Activation volume, activation energy and close-packed volume for 5 wt % PCL solutions in acetone and acetone + CO <sub>2</sub> mixtures .....	185
<b>Table 8.1</b> Liquid-liquid (L-L) demixing pressures (P) at different temperatures (T) for poly ( $\epsilon$ -caprolactone) ( <i>M<sub>w</sub></i> = 14 K and 65 K) solutions in acetone + carbon dioxide mixtures. .....	215
<b>Table 8.2</b> Density data for 5 wt % PCL ( <i>M<sub>w</sub></i> = 14 K) solution in acetone solutions at high pressures .....	217
<b>Table 8.3</b> Density data for the 5 wt % PCL ( <i>M<sub>w</sub></i> = 14 K) solutions in acetone + CO <sub>2</sub> mixtures containing 5, 10, 20 wt % CO <sub>2</sub> at high pressures .....	218
<b>Table 8.4</b> Density data for the 1 wt % PCL ( <i>M<sub>w</sub></i> = 14 K) solutions in acetone (59 wt %) + CO <sub>2</sub> (40 wt %) mixture at high pressures .....	219
<b>Table 8.5</b> Density data for the 5 wt % PCL ( <i>M<sub>w</sub></i> = 14 K) solutions in acetone (55 wt %) + CO <sub>2</sub> (40 wt %) mixture at high pressures .....	220
<b>Table 8.6</b> Density data for the 5 wt % PCL ( <i>M<sub>w</sub></i> = 14 K) solutions in acetone + CO <sub>2</sub> mixture containing 50 and 60 wt % CO <sub>2</sub> at high pressures .....	221
<b>Table 8.7</b> Density data for the 10 wt % PCL ( <i>M<sub>w</sub></i> = 14 K) solutions in acetone (50 wt %) + CO <sub>2</sub> (40 wt %) mixture at high pressures .....	222
<b>Table 8.8</b> Density data for the 20 wt % PCL ( <i>M<sub>w</sub></i> = 14 K) solutions in acetone (40 wt %) + CO <sub>2</sub> (40 wt %) mixture at high pressures .....	223
<b>Table 8.9</b> Density data for the 10 wt % PCL ( <i>M<sub>w</sub></i> = 65 K) solutions in acetone (50 wt %) + CO <sub>2</sub> (40 wt %) mixture at high pressures .....	224
<b>Table 9.1</b> Demixing pressures at different temperatures for 10 wt % PCL ( <i>M<sub>w</sub></i> = 14,300), PMMA ( <i>M<sub>w</sub></i> = 15,000 or 540,000), and their blend solutions in acetone + carbon dioxide mixtures. ....	258
<b>Table 9.2</b> Melting and crystallization temperatures, and heat of melting and crystallization for PCL ( <i>M<sub>w</sub></i> = 14 K) and its blends with PMMA ( <i>M<sub>w</sub></i> = 15 K or 540 K) .....	261

<b>Table 10.1</b> High-pressure homogeneous homopolymerization of methyl methacrylate in acetone at 343 K with 0.04 wt % AIBN as initiator. ....	301
<b>Table 10.2</b> Changing rates of densities and viscosities in both the early and late stages of the polymerizations and the transition time between the two stages at different pressures .....	302
<b>Table 10.3</b> Apparent overall rate constants for MMA polymerizations in acetone at different pressures s estimated from change in density or viscosity .....	303
<b>Table 11.1</b> High-pressure heterogeneous Homopolymerizations of MDO to produce PCL in CO <sub>2</sub> .....	341
<b>Table 11.2</b> Assignments for the <sup>1</sup> H and <sup>13</sup> C NMR spectrums for poly (ε-caprolactone) from MDO polymerization. ....	342
<b>Table 11.3</b> High-pressure heterogeneous copolymerizations of 2-methylene-1,3-dioxepane (MDO) with methyl methacrylate (MMA), styrene (ST), and acrylonitrile (AN) in CO <sub>2</sub> . .....	343
<b>Table 11.4</b> Assignments for the <sup>1</sup> H and <sup>13</sup> C NMR spectrums for PMMA sample. .....	344
<b>Table 11.5</b> Assignments for the <sup>1</sup> H and <sup>13</sup> C NMR spectrums for poly (MDO-co-MMA) sample. ....	345
<b>Table 10.6</b> Estimation of the T <sub>g</sub> of PCL from MDO polymerization .....	346
<b>Table 10.7</b> Estimation of the MDO fraction in copolymers .....	346