

**Table 3.** Recovery of *Cylindrocladium parasiticum* from speckled peanut seed of VA 98R after pods were dried indoors using forced air with no heat (air) or outdoors using forced air with supplemental heat at three locations within drying trailers (top, middle, bottom) in 2000.

Trial and drying condition	Temperature (°C)			Seed assay <sup>a</sup> (% positive)
	Max.	Min.	Avg.	
<b><i>Trial 1</i></b>				
Air.....	26.6	20.9	24.9	98
Heat-top.....	32.5	24.4	27.9	98
Heat-middle.....	33.5	23.8	29.0	100
Heat-bottom.....	35.4	23.8	30.0	96
<b><i>Trial 2</i></b>				
Air.....	22.5	17.8	20.0	98
Heat-top.....	24.8	9.3	14.6	96
Heat-middle.....	28.3	8.5	15.5	98
Heat-bottom.....	30.0	3.3	12.5	98
<b><i>Trial 3</i></b>				
Air.....	22.0	11.3	19.6	96
Heat-top.....	19.6	7.0	14.0	92
Heat-middle.....	16.0	6.4	14.2	94
Heat-bottom.....	21.9	4.5	13.7	74
<b><i>Trial 4</i></b>				
Air.....	27.0	24.0	25.5	98
Heat-top.....	27.8	22.0	26.2	100
Heat-middle.....	-	-	-	100
Heat-bottom.....	33.5	25.5	28.0	96
<b><i>Trial 5</i></b>				
Air.....	27.5	20.3	23.0	100
Heat-top.....	26.8	12.0	18.0	100
Heat-middle.....	30.8	12.5	20.2	100
Heat-bottom.....	32.3	11.5	20.3	100
<b><i>Trial 6</i></b>				
Air.....	26.5	22.6	22.6	94
Heat-top.....	26.5	17.5	20.0	96
Heat-middle.....	27.0	17.0	19.3	94
Heat-bottom.....	29.5	14.5	19.3	98

<sup>a</sup> Data are recovery percentages of *C. parasiticum* from 50 seed. Seeds were assayed on a selective medium.