

## List of Figures

|   | Page |
|---|------|
| Figure 1.1 Distribution of natural products as drugs  | 3    |
| Figure 1.2 World market for drugs from plant sources  | 7    |
| Figure 1.3 <i>Podophyllum peltatum</i>  | 9    |
| Figure 1.4 <i>Catharanthus roseus</i>   | 10   |
| Figure 1.5 <i>Taxus brevifolia</i>  | 11   |
| Figure 1.6 96 well microtiter plate for A2780 bioassay  | 17   |
| Figure 2.1 <i>Strophanthus boivinii</i>   | 25   |
| Figure 2.2 Basic skeleton of cardenolide glycosides   | 28   |
| Figure 2.3 Two classes of cardiac glycosides  | 29   |
| Figure 2.4 Cardenolide glycosides isolated by Mr. Adou from <i>S.boivinii</i>                     | 32   |
| Figure 2.5 <sup>1</sup> H NMR spectrum of compound <b>1</b> in pyridine- <i>d</i> <sub>5</sub>    | 34   |
| Figure 2.6 Key HMBC correlations of fragment <b>1</b>   | 35   |
| Figure 2.7 Key HMBC correlations of A/B and B/C rings of compound <b>1</b>                        | 36   |
| Figure 2.8 Key HMBC correlations for connecting sugar moiety to the aglycone of compound <b>1</b> | 36   |
| Figure 2.9 Key HMBC correlations for attachment of sugar II to sugar I of compound <b>1</b>       | 36   |
| Figure 2.10 Key ROESY correlations of aglycone portion of compound <b>1</b>                       | 37   |

|  |    |
|--|----|
| Figure 2.11 ROESY correlations of the D ring and the lactone unit of compound <b>1</b>   | 38 |
| Figure 2.12 Key ROESY correlations of sugar I and the aglycone of compound <b>1</b>  | 38 |
| Figure 2.13 Final structure and key HMBC and ROESY correlations of compound <b>1</b>   | 39 |
| Figure 2.14 $^1\text{H}$ NMR spectrum of compound <b>2</b> in pyridine- $d_5$  | 41 |
| Figure 2.15 Key HMBC correlations of the aglycone of compound <b>2</b>   | 43 |
| Figure 2.16 Key ROESY correlations of the aglycone of compound <b>2</b>  | 43 |
| Figure 2.17 Key ROESY correlations of the sugar portion of compound <b>2</b>   | 44 |
| Figure 2.18 HMBC connection of the sugar to the aglycone of compound <b>2</b>  | 44 |
| Figure 2.19 Final structure and key HMBC and ROESY correlations of compound <b>2</b>   | 45 |
| Figure 2.20 Key HMBC correlations of the two sugar moieties of compound <b>3</b>   | 47 |
| Figure 2.21 Key ROESY correlations of the two sugar moieties of compound <b>3</b>  | 48 |
| Figure 2.22 Final structure and key HMBC and ROESY correlations of compound <b>3</b>   | 49 |
| Figure 2.23 $^1\text{H}$ NMR spectrum of compound <b>4</b> in methanol- $d_4$  | 51 |
| Figure 2.24 Digitoxigenin-3-O- $\beta$ -D-glucopyranosyl-(1 $\rightarrow$ 4)- $\alpha$ -L-acofriopyranoside (compound <b>4</b> ) | 52 |
| Figure 3.1 $^1\text{H}$ NMR spectrum of compound <b>1</b> in methanol- $d_4$   | 68 |

|  |    |
|--|----|
| Figure 3.2 Key HMBC correlations of A/B ring fusion of fragment 1                              | 69 |
| Figure 3.3 Key HMBC correlations of B/C ring fusion of fragment 2                              | 70 |
| Figure 3.4 Key HMBC correlations of C/D ring fusion of fragment 3                              | 70 |
| Figure 3.5 HMBC correlations for attachment of the alkyl side chain at C-17                    | 70 |
| Figure 3.6 Key ROESY correlations of compound <b>1</b>   | 71 |
| Figure 3.7 Final structure and key HMBC correlations of compound <b>1</b>                      | 72 |
| Figure 3.8 Key ROESY correlations of compound <b>1</b>   | 72 |
| Figure 3.9 <sup>1</sup> H NMR spectrum of compound <b>2</b> in pyridine- <i>d</i> <sub>5</sub> | 74 |
| Figure 3.10 Key HMBC correlations of A/B ring fusion of fragment 1                             | 75 |
| Figure 3.11 Final structure and key HMBC and ROESY correlations of compound <b>2</b>           | 77 |