

List of Figures

Figure		Page
1	Historical corn yields 1890-1999 (USDA NASS).	12
2	1998 Double Cropped-corn – Whitethorne gray leaf spot (<i>Cercospora zea-maydis</i>) disease progress curve.	22
3	Area under the disease progress curve (AUDPC) response for fungicide in 1998.	22
4	Surface plot of the number of plants exhibiting virus symptoms of MDM and MCD in each plot of the 1999 double-cropped corn experiment at Whitethorne.	25
5	Reference map for latitude and longitude in Virginia.	31
6	Day length as a function of latitude.	39
7	Day length as a function of latitude in the middle-Atlantic U.S.	40
8	Primary soil component name for Virginia from the USGS STATSGO database.	43
9	Surface soil texture for Virginia from the USGS STATSGO database.	43
10	Soil drainage classification for Virginia from the USGS STATSGO database.	44
11	30-year mean annual frost free days for Virginia.	44
12	30-year mean annual frost free days minus 50 for Virginia.	45
13	30-year mean accumulated degree days from April to October (base 50°F) for Virginia.	45
14	30-year mean accumulated degree days from June to November (base 50°F) for Virginia.	46
15	30-year mean first frost date for Virginia.	46
16	30-year mean last frost date in Virginia.	47
17	Percent slope calculated from the National Elevation Dataset (USGS).	47

18	Aspect calculated from the National Elevation Dataset (USGS).	48
19	National Land Cover Dataset (c. 1992) (USGS).	48
20	Relative suitability classification for annual degree-day (base 50°F) accumulation used for full season corn calculations.	49
21	Relative suitability classification for annual degree-day (base 50°F) accumulation, used for double cropped corn calculations.	50
22	Relative suitability classification for the mean number of frost free days used for full season corn calculations.	50
23	Relative suitability classification for the mean number of frost free days minus 50, used for double cropped corn calculations.	51
24	June 30-year mean Thornthwaite potential evapotranspiration (cm) in Virginia.	51
25	July 30-year mean Thornthwaite potential evapotranspiration (cm) in Virginia.	52
26	August 30-year mean Thornthwaite potential evapotranspiration (cm) in Virginia.	52
27	June 30-year mean Thornthwaite potential evapotranspiration minus precipitation (cm) in Virginia.	53
28	July 30-year mean Thornthwaite potential evapotranspiration minus precipitation (cm) in Virginia.	53
29	August 30-year mean Thornthwaite potential evapotranspiration minus precipitation in Virginia.	54
30	Thornthwaite potential evapotranspiration minus precipitation (cm) during full season corn silking months in Virginia.	54
31	Relative suitability classification for full season corn PET-P for silking months only.	55
32	Thornthwaite potential evapotranspiration minus precipitation (cm) during double cropped corn silking months in Virginia.	55
33	Relative suitability classification for combined double cropped corn PET-P silking months only.	56

34	The difference in PET-P expressed as double cropped corn silking months minus full season corn silking months.	57
35	Relative suitability classification for soil surface texture used in the analysis.	58
36	Relative suitability classification for soil drainage used in the analysis.	58
37	Combined relative suitability classification for soil surface texture and drainage used in the analysis.	59
38	Relative suitability classification for percent slope as calculated from NED.	59
39	Relative suitability classification for aspect as calculated from NED.	60
40	Relative suitability classification for combined full season corn components.	60
41	Relative suitability classification for combined double cropped corn components.	61