

Personal Acknowledges

I would like to express my gratitude to my research advisor, Dr. Larry Taylor, for his patience and guidance throughout my research since 2005. My special thanks to Dr. Medhi Ashraf-Khorassani for his time, encouragement, and advice whenever I have needed it. I am also grateful to my committee members, Dr. David Kingston and Dr. Sungsool Wi, for their suggestions and understanding, and for taking the time to serve as my committee members.

I am grateful to my parents, my brother and sister for their constant love and support. I would like to extend my deepest thanks to my wife, Josephine Chan for her love, encouragement, and inspiration.

I would also like to thank all my friends in Blacksburg who have helped in making me feel at home here.

Acknowledgement to Funding

The Author would like to thank Dr. J. David Pinkston for critically reviewing the manuscript. We thank Jeff Caldwell from Princeton Chromatography Inc. for donating the 4-ethylpyridine column; Mettler Toledo AutoChem for donating the SFC instrumentation and Isco Inc. for donating the SFE instrumentation.

TABLE OF CONTENTS

	<u>Page</u>
Abstract	ii
Personal Acknowledgments	iii
Acknowledgement to Funding	iv
Table of Contents	v
List of Figures	vii
List of Tables	x
I. Introduction	1
II. Previous Chromatographic Methods for Analysis of Phospholipids	7
III. Feasibility of Phospholipids Separation via Packed Column Supercritical Fluid Chromatography with Mass Spectrometric and Light Scattering Detection	
3.1 Introduction	14
3.2 Experimental	
3.2.1 Chemicals	19
3.2.2 SFC-ELSD	19
3.2.3 SFC-MS Instrumentation	20
3.3 Results and Discussion	
3.3.1 Elution with Primary Mobile Phase Modifier	21
3.3.2 Elution with Basic Additive and Primary Modifier	27
3.3.3 Elution with Acidic Additive and Primary Modifier	31
3.3.4 Elution with Ionic Additive and Primary Modifier	33
3.3.5 Effect of Number of Theoretical Plates	36
3.4 Conclusions	38
IV. The Isolation and Separation of Phospholipids from Soybean Lecithin by Supercritical Fluid Technologies	
4.1 Introduction	39
4.2 Supercritical Fluid Extraction	44
4.3 Experimental	
4.3.1 SFE	46
4.4 Results and Discussion	
4.4.1 Deoiling of crude soybean lecithin	47
4.4.2 Pressure effect on SFE of PC from de-oiled lecithin	48
4.4.3 Temperature effect on SFE of PC from de-oiled lecithin	51

4.4.4 SFE fractionation of PC from de-oiled lecithin	51
4.4.5 Internal adsorbent effect on SFE extraction of de-oiled lecithin	57
4.4.6 Identification of the other polar lipids	59
4.5 Conclusion	65
References	66

LIST OF FIGURES

<u>Figure</u>	<u>Description</u>	<u>Page</u>
1	Classification of Lipids	4
2	Structure of selected lipids and their primary hydrolysis products	5
3	Basic Phospholipids Structure	6
4	Phase diagram of carbon dioxide	11
5	Proposed mechanism for modification of bare silica phase by ionic additive	16
6	Molecular structures of selected phospholipids: phosphatidylcholine (PC) (M.M.=757 Da), phosphatidyl-ethanolamine (PE) (M.M.=715 Da), phosphatidylinositol (M.M.=833 Da), phosphatidylserine (PS) (M.M.=758 Da)	18
7	SFC-ELSD traces for the separation of phospholipid mixture with pressurized CO ₂ -modified with pure methanol on various stationary phases. See Experimental Section for chromatographic conditions.	22
8	Mass spectra of the two PC peak components eluting on the 4-ethylpyridine column with only methanol as the modifier. A = major peak; B = shoulder eluting in front of the main peak.	24
9	Mass spectra of the two PE peak components eluting on the 4-ethylpyridine column with only methanol as the modifier. A = main peak; B = shoulder eluting in front of the main peak	25
10	SFC-ELSD of phospholipid mixture with modified CO ₂ on the four stationary phases. The modifier consisted of methanol with 5.0 mM isopropylamine. Chromatographic conditions: flow rate 2 mL/min, methanol/additive concentration raised from 15% to 55% at 4% per minute and held for 5 minutes at 55%.	28
11	Mass spectrum of PI peak components eluting on the diol column. Methanol is the primary modifier. Isopropyl amine (5.0 mM in methanol) was used as the additive.	29
12	Mass spectrum of PS peak component eluting on the diol column. Methanol is the primary modifier. Isopropyl amine (5.0 mM in methanol) was used as the additive.	30

13	SFC-ELSD of phospholipid mixture with modified CO ₂ on the four stationary phases. The modifier consisted of methanol with 5 mM trifluoroacetic acid. Chromatographic conditions: flow rate 2 mL/min, methanol/additive concentration raised from 15% to 55% at 4% per minute and held for 5 minutes at 55%.	32
14	SFC-ELSD of phospholipid mixture with modified CO ₂ on the four stationary phases. The modifier consisted of methanol with 5 mM ammonium acetate. Chromatographic conditions: flow rate 2 mL/min, methanol/additive concentration raised from 15% to 55% at 4% per minute and held for 5 minutes at 55%.	34
15	Effect of number of theoretical plates on separation of PC	37
16	Molecular structures of phospholipids isolated from soybean lecithin via supercritical fluid fractionation	41
17	Proposed molecular structures of glycolipids isolated from soybean lecithin via supercritical fluid fractionation	42
18	SFC/ELSD trace of soybean lecithin extract with pure CO ₂ . Pressure: 360 atm, Temperature: 40°C, mass of soybean lecithin: 110 mg	49
19	SFC/ELSD trace of soybean lecithin extracts at different pressure using 5% methanol/ CO ₂ at 40°C.	50
20	SFC/ ELSD trace of de-oiled soybean lecithin extracts at different modifier percentage without silica adsorbent	54
21	SFC/ ELSD trace of de-oiled soybean lecithin extracts at different modifier percentage without silica adsorbent (cont')	55
22	Packed extraction vessel with soybean lecithin and silica gel	56
23	SFC/ ELSD trace of soybean lecithin extracts at different modifier percentage with silica as the internal adsorbent	58
24	Mass spectrum of polar lipid I(retention time = 5.1min) eluting on the diol column with 8% methanol as the modifier (Fraction 2 from Figure 23)	60
25	SFC/UV trace of the second peak (retention time = 5.8 min) in Fraction 2 from Figure 23 (<i>m/z</i> range: 480-800)	61

26	Mass spectra of polar lipid II eluting on the diol column in Fraction # 2 from Figure 23. A = front peak; B = shoulder peak.	62
27	Mass spectrum of the polar lipid III (retention time = 8.9 min) eluting on the diol column with fraction #4 (from Figure 23) as the modifier.	64

TABLE OF CONTENTS

<u>Figure</u>	<u>Description</u>	<u>Page</u>
1	Physical properties of gas, supercritical fluid and liquid	12
2	Main components of polar lipids from soybean lecithin	40
3	Optimized dynamic SFE conditions for removal of PC from soybean lecithin	52