

## List of Figures

<b>Figure I-1.</b> The evolution of <i>supramolecular chemistry</i> .	2
<b>Figure I-2.</b> Comparison of biological and material science supramolecular systems.	4
<b>Figure I-3.</b> Crystal structure of valinomycin binding a potassium ion.	5
<b>Figure I-4.</b> Self-assembly of helicates.	8
<b>Figure I-5.</b> A rotaxane, <b>I-9</b> , and pseudorotaxane, <b>I-10</b> .	10
<b>Figure I-6.</b> Various polyrotaxanes and polypseudorotaxanes.	12
<b>Figure I-7.</b> ‘Directed’ or ‘chemical conversion’ approach for rotaxane synthesis.	13
<b>Figure I-8.</b> <i>Statistical</i> and <i>thermodynamic</i> control approaches to rotaxane synthesis.	14
<b>Figure I-9.</b> Self-assembling approaches to rotaxanes.	15
<b>Figure I-10.</b> Example of a rotaxane synthesis using metal complexation.	18
<b>Figure II-1.</b> Comparison of binding constants for 18-crown-6 (left) and benzo-18-crown-6 (right).	28
<b>Figure II-2.</b> Comparison of the conformations of 18-crown-6 in hydrophilic and lipophilic medium.	29
<b>Figure II-3.</b> Pedersen’s synthetic methods for crown ether syntheses.	31
<b>Figure II-4.</b> Example of template effect for the synthesis of 18-crown-6.	32
<b>Figure II-5.</b> Synthetic method for synthesis of phenylene-based crown ethers.	33
<b>Figure II-6.</b> Synthesis of bis( <i>meta</i> -phenylene)-26-crown-8, <b>II-15</b> .	34
<b>Figure II-7.</b> Synthesis of various mono-substituted <b>DB24C8s</b> .	37
<b>Figure II-8.</b> Synthesis of <b>II-17</b> by <b>Method A</b> .	38
<b>Figure II-9.</b> Synthesis of succinimide ester of 1,3-phenylene-16-crown-5, <b>II-29</b> .	39
<b>Figure II-10.</b> Synthesis of mono-substituted <b>BMP32C10s</b> .	40
<b>Figure II-11.</b> Synthesis of diacid bis- <i>meta</i> -phenylene-20-crown-6 ( <b>II-37</b> ).	41
<b>Figure II-12.</b> Attempted synthesis of bis(5-carbomethoxy-1,3-phenylene)-bis(1,4-phenylene)-30-crown-8 ( <b>II-45</b> ).	42
<b>Figure III-1.</b> First reported cryptand.	62
<b>Figure III-2.</b> Simmons' "in-out" bicyclic amines with the "out" conformation on the left and "in" on the right.	63
<b>Figure III-3.</b> Synthesis of bis-1,3,5-phenylene cryptands.	66
<b>Figure III-4.</b> Synthesis of 5-benzyloxyresorcinol ( <b>III-8</b> ) via <b>Method B</b> .	69
<b>Figure III-5.</b> First DSC of bis-1,3,5-phenylene-[5.5.5]-cryptand, <b>III-5</b> .	71
<b>Figure III-6.</b> Second DSC of bis-1,3,5-phenylene-[5.5.5]-cryptand, <b>III-5</b> .	72
<b>Figure IV-1.</b> Synthesis of dibenzyl- ( <b>IV-1</b> ) and di- <i>n</i> -butyl- ( <b>IV-2</b> ) ammonium hexafluorophosphates.	81
<b>Figure IV-2.</b> Synthesis of diphenethylammonium hexafluorophosphate ( <b>IV-8</b> ).	83
<b>Figure IV-3.</b> Synthesis of diammonium salts.	84
<b>Figure IV-4.</b> Synthesis of partially deuterated diammonium salts.	84
<b>Figure V-1.</b> Comparison of uncomplexed and complexed 18-crown-6 ( <b>V-2</b> , <b>18C6</b> ).	96
<b>Figure V-2.</b> Cation binding by crown ethers with various ring sizes.	99
<b>Figure V-3.</b> Cation binding of bibracchial lariat ethers.	101

<b>Figure V-4.</b> Complex between <b>18C6</b> and $R-NH_3^+$ .	102
<b>Figure V-5.</b> Binding constants for crown-type hosts with <i>t</i> -butyl ammonium thiocyanate.	103
<b>Figure V-6.</b> Cartoon depicting "logical progression" from the simple [1.1]pseudorotaxane	107
<b>Figure V-7.</b> First rotaxane formed between a secondary ammonium ion and <b>DB24C8</b> .	108
<b>Figure V-8.</b> Rotaxanes from secondary ammonium ions and crown ethers produced by Stoddart <i>et al.</i>	108
<b>Figure V-9.</b> Crystal structure of a hydrogen-bonded polypseudorotaxane from the dibenzylammonium ion <b>V-5</b> and <b>DB24C8</b> .	109
<b>Figure V-10.</b> Two different exchange rates, a) slow- and b) fast-exchange, relative to NMR time scale for complexation.	111
<b>Figure V-11.</b> Mole ratio plot for complex formed between the bicyclic molecule <b>V-6</b> and <i>p</i> -nitrophenol.	112
<b>Figure V-12.</b> Plot of $[S]_o/[R]_o$ versus $\delta$ for various $K_a$ values	113
<b>Figure V-13.</b> Plot of $[RS]$ versus $[S]$ at various $K_a$ values. $[R]_o = 10\text{mM}$	114
<b>Figure V-14.</b> Stacked $^1\text{H}$ NMR plots (400 MHz, $\text{CD}_3\text{CN}$ , ambient $T$ ) for a) <b>BMP26C8</b> , b) <b>BMP26C8</b> + <b>V-9•PF<sub>6</sub></b> (1:1 stoichiometry, 0.010 M), and c) <b>V-9•PF<sub>6</sub></b> .	124
<b>Figure V-15.</b> Stacked $^1\text{H}$ NMR plots (400 MHz, $\text{CD}_3\text{CN}$ , ambient $T$ ) for a) <b>BMP26C8</b> , b) <b>BMP26C8</b> + <b>V-10•PF<sub>6</sub></b> (1:1 stoichiometry, 0.010 M), and c) <b>V-10•PF<sub>6</sub></b> .	125
<b>Figure V-16.</b> Positive ion ESI-MS for 1:1 mixture of <b>BMP26C8</b> and <b>V-8•PF<sub>6</sub></b> .	126
<b>Figure V-17.</b> Positive ion ESI-MS for 1:1 mixture of <b>BMP26C8</b> and <b>V-9•PF<sub>6</sub></b> .	127
<b>Figure V-18.</b> Positive ion ESI-MS for 1:1 mixture of <b>BMP26C8</b> and <b>V-10•PF<sub>6</sub></b> .	129
<b>Figure V-19.</b> X-ray structure of 26-membered macrocycle <b>BMP26C8</b> : a) ORTEP drawing b) space-filling representation.	131
<b>Figure V-20.</b> Stacked $^1\text{H}$ NMR plots (400 MHz, $\text{CD}_3\text{CN}$ , ambient $T$ ) for a) <b>BMP32C10</b> , b) <b>BMP32C10</b> + <b>V-9•PF<sub>6</sub></b> (1:1 stoichiometry, 0.010 M), and c) <b>V-9•PF<sub>6</sub></b> .	133
<b>Figure V-21.</b> Low Resolution FAB-MS for <b>BMP32C10</b> + <b>V-9•PF<sub>6</sub></b> .	134
<b>Figure V-22.</b> Side-on and top views of the X-ray structure of the [2.1]pseudorotaxane for <b>BMP32C10:(V-9)<sub>2</sub>•2(PF<sub>6</sub>)</b>	135
<b>Figure V-23.</b> Mole ratio plot for <b>BMP32C10</b> + <b>V-9•PF<sub>6</sub></b> in acetonitrile- $d_3$ ( $[V-9•PF_6] = 0.010\text{ M}$ ).	137
<b>Figure V-24.</b> Scatchard plot for <b>BMP32C10</b> + <b>V-9•PF<sub>6</sub></b> .	138
<b>Figure V-25.</b> Stacked $^1\text{H}$ NMR plots (400 MHz, acetone- $d_6$ , ambient $T$ ) for a) <b>BMP32C10</b> , b) <b>BMP32C10</b> + <b>V-8•PF<sub>6</sub></b> (1:1 stoichiometry, 0.010 M), and c) <b>V-8•PF<sub>6</sub></b> .	143
<b>Figure V-26.</b> LRFAB-MS for <b>BMP32C10</b> + <b>(V-8)<sub>2</sub>•2(PF<sub>6</sub>)</b> .	144
<b>Figure V-27.</b> Side-on and top views of the X-ray structure of the pseudorotaxane pseudorotaxane <b>BMP32C10:(V-8)<sub>2</sub>•2(PF<sub>6</sub>)</b>	145
<b>Figure V-28.</b> Stacked $^1\text{H}$ NMR (400 MHz, acetone- $d_6$ ) spectra of a) <b>BMP32C10</b> (0.010 M), b) 1:1 solution of <b>BMP32C10</b> and <b>V-14•2(PF<sub>6</sub>)</b> (0.010 M), and c) <b>V-14•2(PF<sub>6</sub>)</b> (0.010 M).	148

<b>Figure V-29.</b> Mole ratio plot for complex between <b>BMP32C10</b> and <b>V-14•2(PF<sub>6</sub>)</b> in acetone-d <sub>6</sub> .	149
<b>Figure V-30.</b> Crystal structure of "cradled barbell" complex between <b>BMP32C10</b> and <b>V-14•2(PF<sub>6</sub>)</b> .	152
<b>Figure V-31.</b> Packing diagram illustrating the $\pi$ -stacking between the aromatic rings of <b>BMP32C10</b> (blue) and <b>V-14•2(PF<sub>6</sub>)</b> (red).	153
<b>Figure V-32.</b> Stacked <sup>1</sup> H NMR plots (400 MHz, CD <sub>3</sub> CN, ambient <i>T</i> ) for a) <b>DB24C8</b> + <b>V-8•PF<sub>6</sub></b> (1:1 stoichiometry, 0.010 M), b) <b>V-8•PF<sub>6</sub></b> , and c) <b>DB24C8</b> . Subscripts c = "complexed" in c).	156
<b>Figure V-33.</b> Mole ratio plot for complex between <b>DB24C8</b> and <b>V-10•PF<sub>6</sub></b> in acetonitrile-d <sub>3</sub> .	158
<b>Figure V-34.</b> Scatchard plot for <b>DB24C8</b> + <b>V-10•PF<sub>6</sub></b> .	159
<b>Figure V-35.</b> Equilibrium equations for the complexation of secondary diammonium salts the <i>p</i> -substituted diammonium salt <b>V-13•2(PF<sub>6</sub>)</b> and the <i>m</i> -substituted diammonium salt <b>V-14•2(PF<sub>6</sub>)</b> with <b>DB24C8</b> .	163
<b>Figure V-36.</b> Stacked <sup>1</sup> H NMR (400 MHz, CDCl <sub>3</sub> :CD <sub>3</sub> CN, 1:1, v:v) spectra of a) <b>DB24C8</b> (0.010 M), b) <b>DB24C8</b> (0.100 M) + <b>V-13•2PF<sub>6</sub></b> (0.010 M) c) <b>DB24C8</b> (0.010 M) + <b>V-13•2PF<sub>6</sub></b> (0.010 M), and d) <b>V-13•2PF<sub>6</sub></b> (0.010 M).	164
<b>Figure V-37.</b> Stacked <sup>1</sup> H NMR (400 MHz, CDCl <sub>3</sub> :CD <sub>3</sub> CN, 1:1, v:v) spectra of a) <b>DB24C8</b> (0.010 M) + <b>V-15•2PF<sub>6</sub></b> (0.010 M) and b) <b>V-23</b> (0.010 M) + <b>V-13•2PF<sub>6</sub></b> (0.010 M).	165
<b>Figure V-38.</b> Stacked <sup>1</sup> H NMR (400 MHz, CDCl <sub>3</sub> :CD <sub>3</sub> CN, 1:1, v:v) spectra of a) <b>DB24C8</b> (0.010 M), b) <b>DB24C8</b> (0.100 M) + <b>V-14•2PF<sub>6</sub></b> (0.010 M) c) <b>DB24C8</b> (0.010 M) + <b>V-14•2PF<sub>6</sub></b> (0.010 M), and d) <b>V-14•2PF<sub>6</sub></b> (0.010 M).	166
<b>Figure V-39.</b> Stacked <sup>1</sup> H NMR (400 MHz, CDCl <sub>3</sub> :CD <sub>3</sub> CN, 1:1, v:v) spectra of a) <b>DB24C8</b> (0.010 M) + <b>V-16•2PF<sub>6</sub></b> (0.010 M) and b) <b>DB24C8</b> (0.010 M) + <b>V-14•2PF<sub>6</sub></b> (0.010 M).	167
<b>Figure V-40.</b> ESI-MS of <b>30C10</b> + <b>V-8•PF<sub>6</sub></b> .	171
<b>Figure V-41.</b> ESI-MS of <b>30C10</b> + <b>V-9•PF<sub>6</sub></b> .	171
<b>Figure V-42.</b> Mole ratio plot for complex between <b>30C10</b> and <b>V-10•PF<sub>6</sub></b> in acetone-d <sub>6</sub> .	173
<b>Figure V-43.</b> Stacked <sup>1</sup> H NMR (400 MHz, CDCl <sub>3</sub> :CD <sub>3</sub> CN, 3:1, v:v) spectra of a) <b>32-Cryptand</b> (0.010 M), b) 1:1 solution of <b>32-Cryptand</b> and <b>V-9•PF<sub>6</sub></b> (0.010 M), and c) <b>V-9•PF<sub>6</sub></b> (0.010 M).	176
<b>Figure V-44.</b> Mole ratio plot for complex between <b>32-Cryptand</b> and <b>V-9•PF<sub>6</sub></b> in acetonitrile-d <sub>3</sub> .	177
<b>Figure V-45.</b> Scatchard plot for <b>32-Cryptand</b> + <b>V-9•PF<sub>6</sub></b> .	178
<b>Figure V-46.</b> Plot of [BMP32C10]:[V-9•PF <sub>6</sub> ] versus $\Delta\delta$ for protons of <b>BMP32C10</b> .	179
<b>Figure V-47.</b> Plot of <b>32-Cryptand</b> : <b>V-9•PF<sub>6</sub></b> versus $\Delta\delta$ for protons of <b>32-Cryptand</b> .	180
<b>Figure VI-1.</b> Flory's "2,3,2,3,1,0 molecule." A hypothetical "dendrimer."	200
<b>Figure VI-2.</b> <i>Divergent</i> and <i>convergent</i> dendrimer approaches.	203
<b>Figure VI-3.</b> The synthesis of the poly(propylene imine) dendrimers	

(reactions <b>A</b> and <b>B</b> ) and alternative, unwanted reactions <b>C</b> and <b>D</b> as discussed in the text.	206
<b>Figure VI-4.</b> Synthesis of <i>N</i> -methyl- <i>N,N'</i> -bis[ <i>N</i> -(5-carboxy-1,3-phenylene-16-crown-5)propylamino]amine ( <b>VI-8</b> ) and <i>N</i> -methyl- <i>N,N'</i> -bis[ <i>N</i> -(5-carboxy-dibenzo-24-crown-8)propylamino]amine ( <b>VI-9</b> ).	209
<b>Figure VI-5.</b> <sup>1</sup> H NMR (300 MHz, CDCl <sub>3</sub> , ambient T) of <i>N</i> -methyl- <i>N,N'</i> -bis[ <i>N</i> -(5-carboxy-1,3-phenylene-16-crown-5)propylamino]amine ( <b>VI-8</b> ).	210
<b>Figure VI-6.</b> <sup>13</sup> C NMR (75 MHz, CDCl <sub>3</sub> , ambient T) of <i>N</i> -methyl- <i>N,N'</i> -bis[ <i>N</i> -(5-carboxy-1,3-phenylene-16-crown-5)propylamino]amine ( <b>VI-8</b> ).	210
<b>Figure VI-7.</b> 2D COSY NMR (300 MHz, CDCl <sub>3</sub> , ambient T) of <i>N</i> -methyl- <i>N,N'</i> -bis[ <i>N</i> -(5-carboxy-1,3-phenylene-16-crown-5)propylamino]amine ( <b>VI-8</b> ).	211
<b>Figure VI-8.</b> EI-MS of <i>N</i> -methyl- <i>N,N'</i> -bis[ <i>N</i> -(5-carboxy-1,3-phenylene-16-crown-5)propylamino]amine ( <b>VI-8</b> ).	212
<b>Figure VI-9.</b> <sup>1</sup> H NMR (300 MHz, CDCl <sub>3</sub> , ambient T) of <i>N</i> -methyl- <i>N,N'</i> -bis[ <i>N</i> -(5-carboxy-dibenzo-24-crown-8)propylamino]amine ( <b>VI-9</b> ).	213
<b>Figure VI-10.</b> Proposed synthesis of the pentafluorobenzene ester of dibenzo-24-crown-8 ( <b>VI-10</b> ).	214
<b>Figure VI-11.</b> <sup>1</sup> H NMR (300 MHz, CDCl <sub>3</sub> , ambient T) of <b>VI-13</b> , DAB- <i>dendr</i> -(amide-16C5) <sub>64</sub> .	216
<b>Figure VI-12.</b> <sup>1</sup> H NMR (300 MHz, CDCl <sub>3</sub> , ambient T) of product obtained from reaction of <b>VI-4</b> with <b>VI-11</b> , DAB- <i>dendr</i> -(NH <sub>2</sub> ) <sub>4</sub> . Impurities ( <b>i</b> ) have been marked.	216