

**Effects of acid hydrolysis conditions on cellulose nanocrystal yield and properties: A response surface methodology study**

Shuping Dong

Thesis submitted to the faculty of the  
Virginia Polytechnic Institute and State University  
in partial fulfillment of the requirements for the degree of

Master of Science  
In  
Macromolecular Science and Engineering

Maren Roman, Chair  
Timothy E. Long  
Scott Renneckar

April 24, 2014  
Blacksburg, VA

Keywords: cellulose nanocrystals, acid hydrolysis, central composite, Box Behnken

Copyright 2014 Shuping Dong

**Draft 09/01/2009**

(Questions? Concerns? Contact Gail McMillan, Director of the Digital Library and Archives at Virginia Tech's University Libraries: [gailmac@vt.edu](mailto:gailmac@vt.edu))

(Please ensure that Javascript is enabled on your browser before using this tool.)

## Virginia Tech ETD Fair Use Analysis Results

***This is not a replacement for professional legal advice but an effort to assist you in making a sound decision.***

Name: Shuping Dong

Description of item under review for fair use: Figure 2.2. Crystal structures of cellulose 1 and cellulose 2. Source: Ind. Eng. Chem., Anal. Ed., 1954, 16, 351. Figure 2.4. Mechanism of acid catalyzed hydrolysis of cellulose. Source: Industrial& Engineering Chemistry, 1956, 48, 333-335 Figure 2.5. Electron microscopy image of CNCs from cotton fibers. Source: Biochimica et Biophysica Acta, 1953, 10, 499-511 Figure 2.7. (1) Phase separation of cellulose suspension in pure water at different crystallite concentrations.(2) Chiral nematic texture of the anisotropic phase of a cellulose suspension. Source: Langmuir, 1997, 13, 2404-2409 Figure 2.8 Dependence of cellulose solubility and degree of crystallinity, X, of isolated particles on sulfuric acid concentration. Source: Nanoscience and Nanotechnology, 2012, 2(2), 9-13 Figure 2.9 Dependence of average DP and yield of isolated particles on sulfuric acid concentration. Source: Nanoscience and Nanotechnology, 2012, 2(2), 9-13 Figure 2.10. Geometry of a three-factor rotatable CCD with factorial design points, axial points and a center point. Source: Chemometrics and Intelligent Laboratory Systems, 1998, 42, 3-40 Figure 2.11 Geometry of a three-level, three-factor BBD. Source: Analytica Chimica Acta, 2007, 597, 179-186

Report generated on: 05-05-2014 at : 15:41:00

### **Based on the information you provided:**

#### **Factor 1**

Your consideration of the purpose and character of your use of the copyright work weighs: *in favor of fair use*

#### **Factor 2**

Your consideration of the nature of the copyrighted work you used weighs: *in favor of fair use*

#### **Factor 3**

Your consideration of the amount and substantiality of your use of the copyrighted work weighs: *in favor of fair use*

#### **Factor 4**

Your consideration of the effect or potential effect on the market after your use of the

copyrighted work weighs: *in favor of fair use*

**Based on the information you provided, your use of the copyrighted work weighs: *in favor of fair use***

