Virginia Cooperative Extension

Virginia Tech • Virginia State University

. www.ext.vt.edu

2901-1070

Southern Waxmyrtle (Myrica cerifera)

Alex X. Niemiera, Professor, School of Plant and Environmental Sciences, Virginia Tech

Summary:

Foliage: Deciduous broadleaf Height: 15 feet Spread: 15 feet Shape: Upright, multi-stem Southern waxmyrtle is a large evergreen shrub or small tree depending on how one prunes it. This species tolerates wet and dry soils and females have a somewhat showy display of gray berries in the fall/winter.

Plant Needs:

Zone: 7 (6b) to 11 Moisture: Moist to dry pH Range: 3.7 to 6.5 Light: Partial shade to full sun Soil Type: Sandy, loam, or clay

Functions:

Suggested uses for this plant include border, massing, foundation, and screen.

Planting Notes:

Plant both male and female plants for berries.

Extremely adaptable. Thrives in poor sandy or heavy clay soils. Will generally produce more fruit when grown in poor soil. Salt tolerant.

Care:

Easy to maintain. Test soil and adjust soil pH if it is too high.

Problems:

No serious pests. Iron chlorosis is a problem in high pH soils.

Alternatives:

Consult local garden centers, historic or public gardens and arboreta regarding cultivars and related species that grow well in your area.

Related species:

2018

Northern bayberry (*Myrica pensylvanica*) is a large shrub with a similar culture to southern waxmyrtle but differs in its hardiness zone range which is zone 3 to 6. The fruit display is showier on northern bayberry compared to southern waxmyrtle. Both species fix nitrogen which means that bacteria that reside on roots can covert atmospheric (gaseous) nitrogen into a fertilizer form that

Virginia Cooperative Extension programs and employment are open to all, regardless of age, color, disability, gender, gender identity, gender expression, national origin, political affiliation, race, religion, sexual orientation, genetic information, veteran status, or any other basis protected by law. An equal opportunity/affirmative action employer. Issued in furtherance of Cooperative Extension work, Virginia Polytechnic Institute and State University, Virginia State University, Virginia State University, and the U.S. Department of Agriculture cooperating. Edwin J. Jones, Director, Virginia Cooperative Extension, Virginia Tech, Blacksburg; M. Ray McKinnie, Administrator, 1890 Extension Program, Virginia State University, Petersburg.

Virginia Tech

plants can use. Both species will sucker and spread throughout a growing bed (especially northern bayberry). In terms of soil stabilization, this is an advantage. In terms of a garden bed, this may be a disadvantage.

Comments:

In addition to the salt, wet and dry soil tolerance, bayberries are grown for their aromatic leaves and its waxy, gray berries.

Because of their fragrance, the fruit were historically used in making barberry-scented candles.

This material was developed by Carol Ness as part of the Interactive Design and Development Project funded by the Kellogg Foundation.

