

## Colorado Potato Beetle

Thomas P. Kuhar, Department of Entomology, Virginia Tech

Eric. R. Day, Manager, Insect Identification Lab, Department of Entomology, Virginia Tech

Adam M. Alford, Department of Entomology, Virginia Tech

**Scientific Name:** Coleoptera: Chrysomelidae, *Leptinotarsa decemlineata*

**Description:** Adults are convex beetles ~ 3/8 inch long by 1/4 inch wide. The adult thorax is orange with black spots and the wing covers have five yellowish white and five black alternating stripes running lengthwise (Fig. 1). The larvae are smooth, soft-bodied, humpbacked, and are reddish in color with two rows of black spots along each side (Fig. 2); Large larvae reach a length of ~1/2 inch. Eggs are yellow and deposited in masses resembling small sausages standing on end (Fig. 3).

**Range and Plants Attacked:** The Colorado potato beetle (CPB) is found in most regions of the United States except for the Pacific Coast. It feeds exclusively on the foliage of cultivated and wild plants in the nightshade family (Solanaceae). It is a major pest of potatoes and eggplant, and a minor pest of tomatoes.

**Life cycle:** The CPB overwinters in the soil as an adult. In the spring, the emerging beetles seek suitable host plants (such as potato) by walking, but are capable of flying long distances when temperatures exceed 80°F. A female beetle may lay several hundred eggs in her lifetime. Oviposition usually begins by early May, and the eggs are laid in tight clusters of 30 to 60, usually on the underside of host leaves (Fig. 3). The eggs hatch within four to nine days. Larvae are usually found feeding in groups on the undersides of the leaves as they pass through four instars or molts.



Figure 1 Adult Colorado potato beetle. Ward Upham, Kansas State University, Bugwood.org



Figure 2 Larval Colorado potato beetle. Jessica Louque, Smithers Viscient, Bugwood.org



Figure 3 Colorado potato beetle eggs. Jack Rabin, Rutgers NJ Agric Expt. Station, Bugwood.org

Larvae complete their growth in two to three weeks, and peak larval populations develop by mid- to late May when the mature larvae pupate underground in earthen cells. The pupal stage lasts from five to ten days before the summer, or first-generation, adults emerge in June.

The rate of development for each stage depends on the temperature. In eastern Virginia, potatoes are harvested in late June and July; and therefore, most of these first-generation adults will feed on the remaining potato foliage and then enter the soil to overwinter. Some of these adults, however, will produce a partial second generation on tomato, eggplant, fall potato, or other available, suitable foliage. In other states in which potatoes are grown continuously all summer, there is a substantial second generation. There is much overlap within a generation, and typically eggs, larvae, and adults can be found on host foliage at any given time within the season, although generally one stage predominates at a given time.

**Description of Damage:** Adults and larvae feed on foliage in the same manner (Fig. 4). If CPB adults are present early in the season, they will clip small tomato or eggplant transplants or the emerging potato shoots at the ground level. As the larvae grow, they disperse through the plant canopy and consume large portions of the foliage. Large (third and fourth instar) larvae and first generation adults are the stages that do the most damage. If population pressure is heavy, the larvae and adults will completely defoliate the host plant and then feed on the stems. Loss of foliage weakens the plant and consequently results in reduction of marketable yield (tubers, fruit).

#### Organic/Biological Control:

**Control:** The spined soldier bug, *Podisus maculiventris* (Fig. 5), and the two-spotted stinkbug, *Perillus bioculatus* (Fig. 6), prey upon CPB eggs and larvae. Some beetles, specifically the ladybird beetle, *Coleomegilla maculata* (Fig.



Figure 4 CPB feeding damage to potato. Whitney Cranshaw, Colorado State University, Bugwood.org

7), and carabid beetles in the genus *Lebia* also prey on the eggs. Soilborne fungal pathogens such as *Beauveria*





Figure 5 The spined soldier bug (*Podisus maculiventris*), a predator of CPB. Phil Sloderbeck, Kansas State University, Bugwood.org.



Figure 6 The two-spotted stinkbug (*Perillus bioculatus*), a common predator of CPB larvae in VA. Whitney Cranshaw, Colorado State University, Bugwood.org



Figure 7 The spotted lady beetle (*Coleomegilla maculata*) eating Colorado potato beetle eggs. Whitney Cranshaw, Colorado State University, Bugwood.org

some populations if the wrong products are used. Most commercial growers use at planting (or seed-piece) applications of neonicotinoids such as imidacloprid or thiamethoxam, which move systemically from roots to above-ground plant parts to provide control of beetles for several weeks. Several other products are available for use as foliar sprays. A complete list of insecticide options can be found in the Commercial Vegetable Production Recommendations. Virginia Coop. Ext. Pub. No. 456-420. Options for organic growers include: Entrust (Spinosad) azadirachtins, and products containing *Bt tenebrionis*.

*bassiana* may cause high mortality of pupae and overwintering adults. None of these organisms, however, is capable of making an economic impact on large populations.

**Cultural Control:** Because CPB feeds exclusively on solanaceous crops and disperses by walking when temperatures are cool, rotation to a nonsolanaceous crop is one of the most effective cultural control measures a producer can take. This is especially true in eastern Virginia, where the beetles often overwinter in the same fields in which they developed. Plastic-lined

trenches along the side of a potato field where overwintering adults will enter are effective early in the season. Trenches should be at least 12 inches deep, with the sides having a slope of at least 46 degrees. A feasible option for small potato fields is spreading a thick layer of straw mulch after planting to create a favorable environment for the potatoes and an effective barrier to adult beetles. Mechanical control methods such as flaming and handpicking beetles have also been developed.

**Chemical Control and Resistance Management:** CPB has developed resistance to numerous insecticide and chemical control can be difficult for

