

Comparison of the USSES terminal-sire and Siremax composite breeds with the Suffolk breed as terminal sires in an extensive production system: carcass characteristics

Heather N. McKibben,^{*} David R. Notter,[†] Whitney C. Stewart,^{*,1} Warrie J. Means,^{*}
Natalie L. Pierce,[‡] and Joshua B. Taylor[‡]

^{*}Department of Animal Science, University of Wyoming, Laramie, WY 82071; [†]Department of Animal and Poultry Sciences, Virginia Tech, Blacksburg, VA 24061; and [‡]USDA, ARS, U.S. Sheep Experiment Station, Dubois, ID 83423

© The Author(s) 2018. Published by Oxford University Press on behalf of the American Society of Animal Science.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com.

Transl. Anim. Sci. 2018.2:S155–S158
doi: 10.1093/tas/txy042

INTRODUCTION

Crossbreeding can provide the Western U.S. sheep industry with needed improvements in quality and consistency of lamb carcass traits (Leymaster, 2002; Mousel et al., 2012). The U.S. Sheep Experiment Station (USSES) recently compared Suffolk, Texel, and Columbia breeds as terminal sires for improving carcass traits in cross-breed lambs reared in a rangeland system. Suffolk-sired lambs had the greatest hot and chilled carcass weights. Both Suffolk- and Texel-sired lambs had the largest LM area, which indicated greater yielding, more valuable carcasses (Notter et al., 2014). Columbia-sired lambs had the least body wall thickness, an indicator of closely trimmed retail cuts (Mousel et al., 2012). It was readily apparent that no single breed excelled in all carcass traits.

The USSES developed a terminal-sire composite breed (TSC) from the Suffolk, Columbia, and Texel breeds to determine if another modern multi-breed composite could further enhance lamb survival, growth, efficiency, and carcass traits compared with the Suffolk breed. Likewise, the industry developed the Siremax breed from Suffolk, Columbia, Texel, and Hampshire breeds to excel in structural soundness, longevity, and lean growth. To date, the TSC and

Siremax breeds have not been compared with breeds traditionally used as terminal sires in the Western U.S. We hypothesized that when used as terminal sires in a range-ewe production system, the composite breeds would result in crossbred lambs having in similar consistency and yield of carcass characteristics as cross-breed lambs sired by the Suffolk breed. The objective of this study was to compare the TSC and Siremax composite breeds with the Suffolk breed, when used as terminal sires mated to range-type ewes, on carcass characteristics of lambs. The experiment described herein is 1 yr of a 3-yr study having a broader focus of comparison, which includes survival, growth, efficiency, and retail traits.

MATERIALS AND METHODS

The USSES Institutional Animal Care and Use Committee (Dubois, ID) reviewed and approved all animal handling, treatment, and management procedures used in this study.

The experimental design, animal management, and sampling protocols for this study were previously described in Leeds et al. (2012) and Notter et al. (2012). Briefly, mature (2- to 7-yr old) USSES Targhee ($n = 80$), Polypay ($n = 80$), and Rambouillet ($n = 80$) ewes were single-sire mated (10 to 15 ewes per sire) to Suffolk ($n = 8$), Siremax ($n = 8$), or TSC ($n = 8$) rams in October, 2016. The goal was to generate at least 12 crossbred lambs from each ram. Suffolk and Siremax sires were purchased from flocks that partic-

¹Corresponding author: whit.stewart@uwyo.edu

Received March 16, 2018.

Accepted April 14, 2018.

ipated in the National Sheep Improvement Program. The TSC sires were developed at the USSES. Lambs were born in March and April, 2017, and male lambs were castrated shortly after birth. Ewes and their lambs were managed on sagebrush steppe and sub-alpine range through the spring and summer. Lambs were weaned at 101 to 127 d of age. Lambs were stratified by sex and sire breed and assigned to 1 of 24 pens, with 10 to 12 lambs per pen. Lambs of the same-sex and sire breed were assigned to one of four replicate pens. Individual sires and dam breeds were represented in each pen. Lambs were fed a finishing diet (free choice; 78% concentrate, DM basis) and were monitored and treated for illnesses as previously described in [Notter et al. \(2012\)](#).

Within each pen, lambs were randomly assigned to one of three slaughter groups to achieve a target mean harvest BW of 54.4, 61.2, or 68.0 kg. Lambs were weighed weekly and scheduled for harvest when predicted BW of remaining lambs reached the target BW. At harvest, lambs were weighed and transported by a commercial hauler to Mountain States Rosen abattoir in Greeley, CO. A total of 286 lambs (147 ewes and 139 wethers) born in 2017 were evaluated.

After harvest, HCW was recorded for each lamb, and carcasses were stored in a cooler (2–4 °C) for approximately 24 h. After chilling, each carcass was ribbed between the 12th and 13th ribs. Trained personnel subjectively assigned muscling scores, flank streaking, and maturity scores according to [USDA \(1992\)](#) guidelines. Fat depth was measured at the midpoint of the LM and body wall thickness was measured at approximately 12.7 cm from the dorsal midline of both right and left sides. Right and left LM perimeters were traced on acetate paper and later measured with a pork grid to obtain LM area. Measurements for fat depth, body wall thickness, and LM area of the left and right sides were averaged. The calculated percentage of boneless, closely trimmed retail cuts (**calculated %BCTRC**) were calculated as $\%BCTRC = 49.936 - (.0848 \times \text{HCW, pounds}) - (4.376 \times \text{12th rib fat, inches}) - (3.53 \times \text{body wall thickness, inches}) + (2.456 \times \text{LM area, square inches})$ ([Boggs, 2006](#)). Estimated weights of boneless, closely trimmed retail cuts (**EstBCTRCwt**) were estimated by multiplying calculated %BCTRC by HCW. USDA yield grades were assigned on unribbed carcasses taken by trained personnel using [USDA \(1992\)](#) guidelines. Following ribbing of the carcasses, a calculated yield grade was also assigned using the formula (12th rib fat (inches) \times 10) + 0.4 ([USDA, 1992](#)).

HCW, calculated and USDA yield grades, 12th rib back fat, body wall thickness, LM area,

calculated %BCTRC, and EstBCTRCwt were analyzed using the Mixed Model Procedure of SAS (v9.4; SAS Inst. Inc., Cary, NC). The model included fixed effects of sex, sire breed, dam breed, slaughter group, and dam breed \times slaughter group interaction and a random effect of sire nested within sire breed. Orthogonal linear contrasts were used to compare Suffolk-sired lambs to the average of Siremax- and TSC-sired lambs and to compare lambs sired by the composite breeds. Other two-way interactions among fixed effects and the random effect of pen (nested within sex and sire breed) were tested in preliminary analyses but were not significant for any trait and were removed from the final models. Results were interpreted using an alpha of 0.10.

RESULTS AND DISCUSSION

The dam breed \times slaughter group interaction was significant ($P = 0.07$ to 0.09) for HCW, 12th rib fat depth, EstBCTRCwt, and calculated yield grade, but was not significant ($P > 0.18$) for all other measures. The effect of slaughter group was significant ($P < 0.0001$) for HCW, 12th rib fat depth, body wall thickness, LM area, calculated %BCTRC, EstBCTRCwt, and calculated and USDA yield grades ([Table 1](#)). These results were expected, with differences in average lamb age of approximately 1 mo between adjacent slaughter groups. Sire breed effects were significant ($P = 0.08$) for HCW and EstBCTRCwt, but were not significant ($P > 0.13$) for 12th rib fat depth, body wall thickness, LM area, calculated %BCTRC, and calculated and USDA yield grades ([Table 2](#)). Sex effects were significant ($P \leq 0.10$) for HCW, 12th rib fat depth, body wall thickness, LM area, EstBCTRCwt, and calculated and USDA yield grades, while sex had no effect ($P = 0.11$) on EstBCTRCwt. The effect of dam breed was not significant ($P > 0.25$) for any measures.

Beneficial effects have resulted from the use of crossbreeding. Specifically, the use of terminal-sire crossbreeding can improve weaning BW, daily gain, and carcass weights of the lambs ([Sidwell, 1971](#); [Mousel et al., 2012](#)). For this study year, Suffolk-sired lambs had 0.43 kg (3.01%) heavier EstBCTRCwt ($P = 0.02$) and 1.06 kg (3.43%) heavier ($P = 0.08$) HCW compared with TSC- and Siremax-sired lambs ([Table 2](#)). These results of greater HCW from Suffolk-sired lambs are consistent with the previous results reported by [Kirton et al. \(1995\)](#) and [Leymaster \(1981\)](#). While HCW and EstBCTRCwt were greater in Suffolk- sired lambs,

Table 1. Least-squares means and SE by slaughter group for carcass characteristics

Trait ^a	Slaughter group			SE	P value
	1	2	3		
HCW, kg	25.63	29.98	34.76	0.36	<0.0001
12th rib fat, mm	4.20	5.92	6.92	0.26	<0.0001
Body wall thickness, mm	18.34	27.99	30.68	0.48	<0.0001
LM area, cm ²	14.67	17.57	19.03	0.22	<0.0001
Calculated %BCTRC	47.46	46.04	45.25	0.15	<0.0001
EstBCTRCwt	12.14	13.76	15.68	0.15	<0.0001
Calculated Yield Grade	2.05	2.73	3.12	0.10	<0.0001
USDA Yield Grade	1.87	3.19	2.64	0.05	<0.0001

^aMeasures of LM area, 12th rib fat, and body wall thickness were taken between the 12th and 13th ribs of both sides of the carcass and averaged for analysis.

Table 2. Least-squares means and SE for carcass characteristics

Trait ^a	Sire breed			SE	P value	Contrasts (P value)	
	Suffolk	Siremax ^b	TSC ^c			Suffolk vs. composites	TSC vs. Siremax
HCW, kg	30.83	29.77	29.77	0.36	0.08	0.03	0.99
12th Rib fat, mm	5.84	5.61	5.58	0.31	0.81	0.52	0.96
Body wall thickness, mm	25.75	25.82	25.43	0.57	0.87	0.87	0.63
LM area, cm ²	17.17	17.28	16.83	0.22	0.31	0.68	0.15
Calculated % BCTRC	46.13	46.34	46.28	0.18	0.68	0.41	0.79
EstBCTRCwt	14.15	13.72	13.72	0.15	0.08	0.02	0.99
Calculated yield grade	2.70	2.61	2.59	0.12	0.81	0.52	0.96
USDA yield grade	2.50	2.61	2.59	0.05	0.31	0.13	0.74

^aMeasures of LM area, 12th rib fat, and body wall thickness were taken between the 12th and 13th ribs of both sides and averaged for analysis.

^bSiremax was derived from crosses among the Suffolk, Columbia, Texel, and Hampshire breeds.

^cThe USSES TSC was developed at the USDA, ARS, USSES, Dubois, ID, by crossing the Suffolk, Columbia, and Texel breeds.

the TSC-sired lambs produced similar ($P = 0.99$) results to Siremax-sired lambs for all other carcass characteristics measured (Table 2). Based on this single year of data, we reject our initial hypothesis, as the TSC- and Siremax-sired lambs did not match the performance of Suffolk-sired lambs.

IMPLICATIONS

Based on this study year, these data indicate that the development of the TSC and Siremax breeds should continue with increased focus placed on traits associated with carcass weight. Apart from measurable carcass characteristics, other economically relevant whole-production cycle measures such as weight of lamb weaned per ewe exposed, which greatly impact the overall profitability of any sheep enterprise, are currently being analyzed. These data in conjunction with forthcoming preweaning and postweaning production data will clarify the overall utility of the TSC and Siremax breeds as terminal-sire breeds for extensive, range sheep productions systems in the Western U.S.

LITERATURE CITED

- Boggs, D. L. 2006. Livestock and carcasses: an integrated approach to evaluation, grading and selection. Dubuque (IA): Kendall/Hunt Pub. Co.
- Kirton, A. H., A. H. Carter, J. N. Clarke, D. P. Sinclair, G. J. K. Mercer, and D. M. Duganzich. 1995. A comparison between 15 ram breeds for export lamb production I. Live weights, body components, carcass measurements, and composition. *New Zealand J. Agric. Res.* 38:347–360. doi:10.1080/00288233.1995.9513136
- Leeds, T. D., D. R. Notter, K. A. Leymaster, M. R. Mousel, and G. S. Lewis. 2012. Evaluation of Columbia, USMARC-composite, Suffolk, and Texel rams as terminal sires in an extensive rangeland production system: I. Ewe productivity and crossbred lamb survival and preweaning growth. *J. Anim. Sci.* 90:2931–2940. doi:10.2527/jas.2011-4640
- Leymaster, K. A. 2002. Fundamental aspects of crossbreeding of sheep: use of breed diversity to improve efficiency of meat production. *Sheep Goat Res. J.* 17:50–59.
- Leymaster, K. A., and G. M. Smith. 1981. Columbia and Suffolk terminal sire breed effects. *J. Anim. Sci.* 53:1225–1235.
- Mousel, M. R., D. R. Notter, T. D. Leeds, H. N. Zerby, S. J. Moeller, and G. S. Lewis. 2012. Evaluation of Columbia, USMARC-composite, Suffolk, and Texel rams as terminal sires in an extensive rangeland production system: III. Prefabrication carcass traits and organ weights. *J. Anim. Sci.* 90:2953–2962. doi:10.2527/jas.2011-4767

- Notter, D. R., T. D. Leeds, M. R. Mousel, J. B. Taylor, D. P. Kirschten, and G. S. Lewis. 2012. Evaluation of Columbia, USMARC-composite, Suffolk, and Texel rams as terminal sires in an extensive rangeland production system: II. Postweaning growth and ultrasonic measures of composition for lambs fed a high-energy feedlot diet. *J. Anim. Sci.* 90:2941–2952. doi:10.2527/jas.2011-4641
- Notter, D. R., M. R. Mousel, T. D. Leeds, H. N. Zerby, S. J. Moeller, G. S. Lewis, and J. B. Taylor. 2014. Evaluation of Columbia, U.S. Meat animal research center composite, Suffolk, and Texel rams as terminal sires in an extensive rangeland production system: VI. Measurements of live-lamb and carcass shape and their relationship to carcass yield and value. *J. Anim. Sci.* 92:1980–1994. doi:10.2527/jas.2013-7154
- Sidwell, G. M., and L. R. Miller. 1971. Production in some pure breeds of sheep and their crosses. II. Birth weights and weaning weights of lambs. *J. Anim. Sci.* 32:1090–1094.
- USDA. 1992. Official United States standards for grade of lamb, yearling mutton, and mutton carcasses. Washington (DC): AMS, USDA.