

Decomposition squared

E. Sharpe  and H. Zhang 

*Physics Department, Virginia Tech,
850 West Campus Drive, Blacksburg, VA. 24061, U.S.A.*

E-mail: ersharpe@vt.edu, h Zhang96@vt.edu

ABSTRACT: In this paper, we test and extend a proposal of Gu, Pei, and Zhang for an application of decomposition to three-dimensional theories with one-form symmetries and to quantum K theory. The theories themselves do not decompose, but, OPEs of parallel one-dimensional objects (such as Wilson lines) and dimensional reductions to two dimensions do decompose, sometimes in two independent ways. We apply this to extend conjectures for quantum K theory rings of gerbes (realized by three-dimensional gauge theories with one-form symmetries) via both orbifold partition functions and gauged linear sigma models.

KEYWORDS: Chern-Simons Theories, Differential and Algebraic Geometry, Global Symmetries, Supersymmetric Gauge Theory

ARXIV EPRINT: [2405.12269](https://arxiv.org/abs/2405.12269)

Contents

1	Introduction	1
2	Prediction	4
3	Presentations as global orbifolds	8
3.1	Example with central trivially-acting group	9
3.2	First nonbanded example	9
3.3	Second nonbanded example	10
3.4	Third nonbanded example	11
3.5	First trivially-acting nonabelian group example	13
3.6	Second trivially-acting nonabelian group example	14
3.7	Third trivially-acting nonabelian group example	15
4	Quantum K theory via GLSM computations	16
4.1	Warmup: gerbes on projective spaces	16
4.2	General \mathbb{Z}_ℓ gerbes over Grassmannians	19
4.3	Gerbes via weighted Grassmannians	22
4.4	More general weighted Grassmannians	25
4.5	Gerbes on flag manifolds	26
4.6	More general levels: projective spaces	28
5	Conclusions	29
A	Bundles on stacks and gerbes	29

1 Introduction

Decomposition is the observation that a local d -dimensional quantum field theory with a global $(d - 1)$ -form symmetry is equivalent to (“decomposes” into) a disjoint union of local theories without global $(d - 1)$ -form symmetries. Decomposition was first described in 2006 in [1] as part of efforts to resolve some of the technical difficulties in making sense of string propagation on stacks and gerbes [2–4], and it has been documented and confirmed since in many examples, many kinds of examples, in different dimensions, see for example [5–32] for a sample of the literature, and see also [33–36] for introductions and reviews.

One of the original predictions of decomposition [1] was that the Gromov-Witten invariants and quantum cohomology rings of gerbes should be equivalent to the those of disjoint unions of spaces. Gerbes are essentially bundles of one-form symmetries, hence a two-dimensional sigma model whose target space is a gerbe should admit a global one-form symmetry (corresponding to translation along the fibers of the target), and so decompose. This prediction for Gromov-Witten theory was checked in e.g. [37–42].

In this paper, we extend a proposal of [16, section 4], [17, section 3], and discuss extensions of such notions to three-dimensional theories with one-form symmetries, and corresponding predictions of decomposition for quantum K theory. In particular, quantum K theory rings can be computed using three-dimensional gauge theories [43, section 2.4], [44–58], and for the same reasons as above, the three-dimensional gauge theories for gerbes have one-form symmetries. Now, such three-dimensional theories themselves do not decompose, as that would require a two-form symmetry. (A three-dimensional sigma model whose target is a 2-gerbe, on the other hand, would have a global two-form symmetry and so decompose.) However, the quantum K theory rings are computed as OPEs of unlinked parallel Wilson lines, wrapped on the same S^1 , which leads to two parallel effects, potentially two different decompositions:

- Those Wilson lines can be acted upon by the generator of the global one-form symmetry to produce other Wilson lines, so that there is a multiplicity, which results in one decomposition.
- For much the same reasons that electromagnetism of infinite parallel planes in three dimensions reduces to an effectively one-dimensional problem, here the pertinent aspects of the three-dimensional theory are captured by an effective two-dimensional theory. Each such two-dimensional theory has a one-form symmetry, and so decomposes.

More technically, given a BK (one-form) symmetry in three dimensions, after Kaluza-Klein reduction on S^1 , the two-dimensional theory has a $BK \times K$ symmetry:

- Wilson lines wrapped along the S^1 generate dimension-zero defects (and the one-form symmetry BK) in the two-dimensional theory, This leads to one level of decomposition in two dimensions.
- Real-codimension-one defects perpendicular to S^1 (corresponding, as we will argue later, to the zero-form symmetry K) couple to states in copies of the two-dimensional space. (In orbifold constructions, these are twisted sectors along the S^1 .) These are invisible in a dimensional reduction, but appear in a more complete Kaluza-Klein reduction. These are better understood as superselection sectors rather than decomposition, but for the purpose of understanding IR phenomena such as quantum K theory, the effect is very similar.

As a result, deep in the IR, there are potentially two different notions of decomposition operating in such examples. If there is no 't Hooft anomaly between the two-dimensional BK and K (in three dimensions, if there is no self-'t Hooft anomaly in the BK), then both mechanisms operate independently, so that for a (banded) \mathbb{Z}_ℓ gerbe, deep in the IR, one gets ℓ^2 universes, as first remarked in [16, section 4], a result we will see explicitly in both orbifold partition functions and also in physics computations of quantum K theory rings.

In particular, we will use three-dimensional gauge theories to justify and illustrate the conjecture that the quantum K theory ring of a gerbe is equivalent to the quantum K theory ring of a disjoint union (of squared order), as expected from the physics of decomposition.

We should emphasize that this phenomenon is not specific to one-form symmetries in three-dimensional theories. For example, schematically, given a d -dimensional theory with

a \mathbb{Z}_ℓ $(d - k - 1)$ -form symmetry, say, one can construct projection operators on parallel k -dimensional objects, and by doing a Kaluza-Klein reduction along a k -dimensional factor in the spacetime manifold, one potentially gets a disjoint union of ℓ low-energy theories of dimension $d - k$ with a \mathbb{Z}_ℓ $(d - k - 1)$ -form symmetry, each of which separately decomposes, for a total of ℓ^2 universes. (To get this additional structure assumes no mixed 't Hooft anomalies, and also may require, on dimensional grounds, that $d - k - 1 \leq k$, or $2k \geq d - 1$, so that the $(d - k - 1)$ -form symmetry may reduce to a zero-form symmetry.) Other variations also exist, and will be discussed in future work. Related ideas have also appeared in discussions of compactifications of six-dimensional $(2,0)$ theories, see for example [59, section 2.1].

We should also emphasize that to see this phenomenon requires keeping track of modes wrapped on the S^1 . In other words, the point of this paper is to discuss a phenomenon arising in Kaluza-Klein reductions. By contrast, in a dimensional reduction on an S^1 , when all dependence on the S^1 is merely truncated, we do not expect these phenomena to arise.

We begin in section 2 by making a prediction for dimensional reductions and OPEs of parallel one-dimensional objects in three-dimensional G gauge theories with trivially-acting $K \subset G$ (and hence a one-form symmetry). The previous papers [16, 17] considered the case that K is a subset of the center of G , and we extend the proposal to more general K , not necessarily central — meaning, not-necessarily-banded gerbes. In such more general cases, the statement of decomposition is more complex than in cases in which K is central.

In section 3 we discuss that prediction in the case of global orbifolds by finite groups. Our construction of the dimensional reduction explicitly reproduces the form of the prediction of section 2, but we think it useful to illustrate the consequences in a number of different kinds of examples.

In section 4 we turn to gauged linear sigma model (GLSM) computations. In the global orbifolds of the previous section, we could only discuss the form of the decomposition (the disjoint union), but not the quantum K theory rings of the separate universes. Using GLSM methods, we are able to discern both the decomposition as well as the quantum K theory rings of the individual universes. In each case we discuss the quantum cohomology of a two-dimensional GLSM and the quantum K-theory ring from a three-dimensional GLSM, and in each case, discuss how the decomposition can be seen explicitly. Specifically, in subsection 4.1, we review gerbes on projective spaces, for which results already exist in the literature. In subsection 4.2 we discuss general \mathbb{Z}_ℓ gerbes on Grassmannians $G(k, n)$, and in section 4.3, we turn to \mathbb{Z}_ℓ gerbes presented as weighted Grassmannians, analogues of weighted projective spaces. Mathematically, these are special cases of the gerbes in the previous section, but their physical presentation is different, so we repeat the analysis here, and outline how the physical predictions for quantum K theory rings are consistent with expectations of decomposition.

For completeness, in section 4.4 we discuss general weighted Grassmannians and predictions for their quantum K theory rings. In section 4.5 we perform the same analyses for \mathbb{Z}_ℓ gerbes on flag manifolds.

The same methods can be applied to study other spaces beyond those above — for example, gerbes on Fano toric varieties. However, the methods and analyses are essentially the same as that discussed here, so for the purposes of this paper, we feel that the examples above should suffice to set up the conjecture that quantum K theory of gerbes is equivalent to quantum K theory of disjoint unions of spaces, via longitudinal decomposition.

In appendix A we discuss bundles on stacks and gerbes, as relevant for discussions of quantum K theory.

In passing, the fact that dimensional reduction can yield disjoint unions plays an essential role in this paper, and has also been discussed in a different context in [60].

2 Prediction

First, we briefly recall decomposition in two-dimensional gauge theories, before turning to three-dimensional examples. Consider a two-dimensional G gauge theory (which we denote $[X/G]$, in obvious reference to orbifolds, but the idea holds more generally) in which a subgroup K acts trivially. This theory has¹ a one-form symmetry, and so one expects a decomposition. Then [1]

$$\text{QFT}_{2d}([X/G]) = \text{QFT}_{2d}\left(\left[\frac{X \times \hat{K}}{G/K}\right]_{\omega}\right), \tag{2.1}$$

where \hat{K} denotes the set of irreducible representations of K , and ω denotes discrete torsion described in [1]. In the special case that the effectively-acting group G/K acts trivially on \hat{K} , the right-hand-side becomes a disjoint union of G/K gauge theories, as many as irreducible representations of K .

Next, consider a three-dimensional G gauge theory, again denoted $[X/G]$, with a trivially-acting subgroup $K \subset G$. For the moment, we assume the theory does not have a Chern-Simons term, and return to such terms later. This theory has a one-form symmetry, but in three dimensions, this does not predict a decomposition. However, if we consider either a dimensional reduction to two dimensions, or alternatively consider a theory of parallel one-dimensional objects (such as Wilson lines, as relevant to quantum K theory), then the low-energy effective two-dimensional theory decomposes, and we predict that at low energies, decomposition has the form

$$\text{QFT}_{3d\text{eff}}([X/G]) = \text{QFT}_{2d}\left(\coprod_{[g]} \left[\frac{X \times \hat{K}_g}{C(g)/K_g}\right]_{\omega}\right), \tag{2.2}$$

where the disjoint union is over trivially-acting conjugacy classes $[g]$ of G , $C(g) \subset G$ denotes the centralizer of a representative $g \in G$, $K_g \subset C(g)$ is the trivially-acting subgroup of the centralizer, and finally ω denotes discrete torsion, the same discrete torsion that would arise in a two-dimensional $C(g)$ gauge theory with the same matter, as described in [1].

As an important special case, suppose that the trivially-acting subgroup of the (original) group G lies within the center of G : $K \subset Z(G) \subset G$. Then, the set of trivially-acting conjugacy classes is equivalent to the set of elements of K , and the low-energy decomposition above reduces to the statement that

$$\text{QFT}_{3d\text{eff}}([X/G]) = \text{QFT}_{2d}\left(\prod_{g \in K} \prod_{\rho \in \hat{K}} [X/(G/K)]_{\omega(\rho)}\right), \tag{2.3}$$

¹Strictly speaking, one speaks of higher-form symmetries only for abelian groups. However, decomposition is slightly more general — there is a decomposition even if the trivially-acting group is nonabelian.

which has $|K|^2$ universes,² rather than $|K|$ as in the analogous two-dimensional case. This special case was discussed in [16, 17]; part of the point of this paper is to extend that to more general cases.

We can understand this as a consequence of two orthogonal effects, both arising from the one-form symmetry BK of the gauge theory in three dimensions, on a three-manifold of the form $S^1 \times \Sigma$:

- The line operators for the BK along the S^1 reduce to pointlike operators on Σ , and so reduce to a one-form symmetry on Σ , responsible for one decomposition.
- In addition, there are twisted sector states along the S^1 , arising in the gauge theory. In three dimensions, twisted sector states are supported along two-dimensional surfaces, whose intersection with Σ corresponds to line operators for K in the two-dimensional theory, or equivalently line operators for BK in the three-dimensional theory. Strictly speaking, since these do not arise from a separate one-form symmetry but rather a zero-form symmetry, it is better to understand the resulting sectors as superselection sectors.

As a result, schematically the theory has a

$$(\text{decomposition}) \times (\text{superselection sectors}) \tag{2.4}$$

structure, rather than a

$$(\text{decomposition})^2 \tag{2.5}$$

structure per se. However, much of our interest will focus on quantum K theory and other IR effects for which the distinction is moot (hence our focus on low-energy behaviors).

So far, we have discussed theories without Chern-Simons terms. A Chern-Simons term will modify the one-form symmetry, and hence the structure of the decomposition. We compare several cases to illustrate this. We focus on abelian theories, both as prototypes for more general cases, and also because in GLSM computations, generically on the Coulomb branch the gauge symmetry is abelian.

- First, consider a three-dimensional $U(1)$ Chern-Simons theory at level m , a $U(1)_m$ theory. This theory has a $B\mathbb{Z}_m$ (one-form) symmetry, with line operators given by Wilson lines of various charges. If m is even, there are m distinct line operators of the form

$$W_n = \exp\left(in \oint A\right), \tag{2.6}$$

related by $n \cong n \pmod m$, often conventionally labelled [61, appendix C.1]

$$n = 0, \pm 1, \dots, \pm \frac{m-2}{2}, + \frac{m}{2}. \tag{2.7}$$

²As is discussed elsewhere, this is due to a combination of decomposition and superselection sectors, so deep in the IR this is $|K|^2$ universes, but should be more invariantly understood as a combination of decomposition and superselection sectors rather than just decomposition per se.

If m is odd, the theory can be defined only if the underlying three-manifold is spin, and there are $2m$ line operators of the form above (n is no longer quite equivalent to $n + m$). (See [61, appendix C], [62, appendix A], [63, appendix C], [64, section 2.2], [24, section 5.9], [65–67] for more information.)

For more information on the identifications above, see for example [68].

- A three-dimensional $U(1)$ gauge theory with matter fields of charges all multiples of k , and no Chern-Simons term, has a $B\mathbb{Z}_k$ (one-form) symmetry. In this case, the periodicity arises because a Wilson line W_k can end on a field of charge k , so we can use those perturbative fields to ‘break’ Wilson lines, so that $W_k \cong 1$. Since we can write any $W_n = W_{n-k} \otimes W_k$, this results in a periodicity $W_n \cong W_{n-k}$.
- Next, we combine these cases. Consider a $U(1)$ gauge theory with matter fields of charges all multiples of k , and also with a Chern-Simons term at level m . This theory has a $B\mathbb{Z}_{\text{gcd}(m,k)}$ (one-form) symmetry. To see this, we use Bézout’s identity, which says that there exist integers a, b such that

$$am + bk = \text{gcd}(m, k), \tag{2.8}$$

and moreover, integral linear combinations of m and k are multiples of $\text{gcd}(m, k)$. As a result, by using combinations of the two periodicity mechanisms above, the Wilson lines W_n are only distinct for $n \pmod{\text{gcd}(m, k)}$.

For simplicity we restrict to the case m is even. Suppose $m = 2, k = 3$, which have $\text{gcd}(m, k) = 1$. The distinct Wilson lines of the Chern-Simons theory at level 2 have $n = 0, 1$. Now,

$$n = 1 \cong n = 4 \text{ using the } k \text{ periodicity,} \tag{2.9}$$

$$\cong n = 0 \text{ using the } m \text{ periodicity,} \tag{2.10}$$

and so there are no nontrivial Wilson lines — all are equivalent to the identity, as expected from the gcd.

For another example, suppose $m = 6, k = 4$, which have $\text{gcd}(m, k) = 2$. The allowed Wilson lines of the Chern-Simons theory at level 6 have

$$n = 0, \pm 1, \pm 2, +3. \tag{2.11}$$

Now,

$$n = 3 \cong n = 7 \text{ using the } k \text{ periodicity,} \tag{2.12}$$

$$\cong n = 1 \text{ using the } m \text{ periodicity,} \tag{2.13}$$

$$n = 2 \cong n = 6 \text{ using the } k \text{ periodicity,} \tag{2.14}$$

$$\cong n = 0 \text{ using the } m \text{ periodicity,} \tag{2.15}$$

$$n = -1 \cong n = 5 \text{ using the } m \text{ periodicity,} \tag{2.16}$$

$$\cong n = 1 \text{ using the } k \text{ periodicity,} \tag{2.17}$$

$$n = -2 \cong n = 4 \text{ using the } m \text{ periodicity,} \tag{2.18}$$

$$\cong n = 0 \text{ using the } k \text{ periodicity.} \tag{2.19}$$

so that the $U(1)_6$ Chern-Simons theory effectively only has two distinct Wilson lines (W_0, W_1) in the presence of charge 4 matter, as expected from the gcd.

As a result, in the presence of Chern-Simons terms, we must modify our prediction. To further complicate matters, for G gauge theories in which the trivially-acting subgroup $K \subset G$ is not abelian, the decomposition is not solely understandable in terms of one-form symmetries, as BK is only defined for K abelian. In this paper, in the presence of Chern-Simons terms, we only discuss decomposition for G gauge theories in which the trivially-acting subgroup is abelian. (Decomposition will exist more generally, but we leave the matter of straightening out a precise prediction for future work.)

So far we have discussed conditions for the presence of a one-form symmetry that could generate one level of decomposition. To get a second level of decomposition (or rather, independently operating superselection sectors), the two effects much act independently. This means we must also require that the self-'t Hooft anomaly of that one-form symmetry in three dimensions, or equivalently the 't Hooft anomaly in two dimensions between the one-form symmetry and the corresponding reduced zero-form symmetry, vanish.

This 't Hooft anomaly was computed in, for example, [69, section 5.1]. For the level m $U(1)$ Chern-Simons theory with matter of charge k outlined above, it was argued that the 't Hooft anomaly is proportional to

$$\frac{k}{m^2} \pmod{1}. \tag{2.20}$$

Now, consider a G gauge theory with trivially-acting abelian subgroup $K \subset Z(G) \subset G$. Assume that, in the presence of Chern-Simons terms, there is a one-form symmetry BL for $L \subset K$, and let us assume that there is no self-'t Hooft anomaly of BL in three dimensions. Then, in this case, we predict that

$$\text{QFT}_{3\text{d eff}'}([X/G]) = \text{QFT}_{2\text{d}}\left(\prod_{g \in L} \prod_{\rho \in \hat{L}} [X/(G/L)]_{\omega(\rho)}\right), \tag{2.21}$$

which has $|L|^2$ universes. Although a larger subgroup K acts trivially, only $L \subset K$ will result in a decomposition, due to the presence of the Chern-Simons terms. (We leave a systematic prediction for more general cases to future work.)

Now, we turn to quantum K theory, for a gerbe presented as a quotient $[X/G]$ where a subgroup $K \subset G$ acts trivially. Quantum K theory is realized physically in a three-dimensional gauge theory on a three-manifold $\Sigma \times S^1$. The quantum K theory ring is the OPE ring of parallel Wilson lines wrapped on the S^1 , as discussed in [47–58]. In order to reproduce the quantum K theory ring appearing in mathematics, there are also Chern-Simons terms. For a gauge theory with abelian gauge group and no superpotential, to match mathematics, the levels are given by [57, section 2]

$$k^{ab} = -\frac{1}{2}(R_i - 1) \sum_i Q_a^i Q_b^i, \tag{2.22}$$

where the R_i denotes the R -charge of the i th chiral superfield, and Q_a^i is the charge of the i th chiral superfield under the a th $U(1)$ factor in a maximal torus of the gauge group. (If all R charges vanish, this reduces to the $U(1)_{-1/2}$ quantization described in e.g. [52, section 2.2].)

For our purposes, we observe that if there is a trivially-acting \mathbb{Z}_k in the center of the gauge group, then the charges Q_a^i are divisible by k , and so the levels used in computing quantum K theory are divisible by k . As a result, although quantum K theory is computed by a three-dimensional gauge theory with Chern-Simons terms, the Chern-Simons terms do not reduce any one-form symmetry arising from a subgroup of the gauge group acting trivially. Furthermore, the levels are divisible by the square of the charges, so there is no 't Hooft anomaly. In effect, for purposes of understanding decomposition, we can ignore the presence of the Chern-Simons terms.

Thus, for the quantum K theory of a gerbe presented as a quotient $[X/G]$ where a subgroup $K \subset G$ acts trivially, we predict

$$\text{QK}([X/G]) = \text{QK}\left(\prod_{[g]} \left[\frac{X \times \hat{K}_g}{C(g)/K_g} \right]_{\omega}\right), \tag{2.23}$$

Next, we shall check this prediction explicitly, in theories presented as global orbifolds in section 3, and in gauged linear sigma models in section 4.

3 Presentations as global orbifolds

Consider dimensional reductions of orbifolds $[X/G]$ from three dimensions to two-dimensions. As this is a dimensional reduction, we omit dependence on the third dimension, hence we omit analogues of twisted sectors resulting from nontrivially-acting elements of G . However, (conjugacy classes³ of) trivially-acting elements of G can still contribute along the third direction. Then, for any one group element $g \in G$ (representing a conjugacy class) along the third direction, we are left with a two-dimensional orbifold by the centralizer $C(g)$. As a result, we describe the partition function of a dimensionally-reduced orbifold $[X/G]$, on three-manifold $S^1 \times \Sigma$, as

$$\sum_{[g]} [X/C(g)], \tag{3.1}$$

where the sum is over conjugacy classes of G that act trivially on X .

Using known results for two-dimensional decompositions [1], this immediately reproduces the structure of the three-dimensional decomposition in section 2. In this section we will compute the result in a number of examples, to illustrate the range of phenomena that arise. In each case, we will compute the partition function, after dimensional reduction, on a T^2 .

In this section there will be no Chern-Simons terms in the three-dimensional theory to complicate the analysis. We shall consider examples with Chern-Simons terms later in section 4.

In passing, in the special case that all of G acts trivially, this has also been described in [70, 71], in discussions of dimensionally-reducing three-dimensional Dijkgraaf-Witten theory. Analogous results in the condensed matter literature (in a different number of dimensions) are also discussed in [72, section 3.C].

³Conjugacy classes, conjugating by elements of G , instead of group elements, because a gauge transformation will conjugate by elements of G . As any trivially-acting subgroup is normal, conjugation will always map a conjugacy class to itself.

3.1 Example with central trivially-acting group

Consider $[X/D_4]$, where D_4 is the eight-element dihedral group, with trivially-acting central $\mathbb{Z}_2 \subset D_4$, as in [2, section 2.0.1], [1, section 5.2]. We can write $D_4 = \langle a, b \rangle$ where

$$a^2 = 1, \quad b^2 = z, \quad b^4 = 1, \quad z^2 = 1, \quad ba = abz, \quad (3.2)$$

and z generates the \mathbb{Z}_2 center.

Next, we will compute T^2 partition functions after dimensional reduction.

In either of the $1, z$ sectors (meaning, cases in which $1, z$ are inserted along the third S^1), the commuting pairs in D_4 which commute with that third group element form all of the commuting pairs in an ordinary two-dimensional orbifold, namely

$$1, z \begin{array}{|c|} \hline \square \\ \hline 1, z \\ \hline \end{array}, \quad 1, z \begin{array}{|c|} \hline \square \\ \hline a, az \\ \hline \end{array}, \quad 1, z \begin{array}{|c|} \hline \square \\ \hline b, bz \\ \hline \end{array}, \quad 1, z \begin{array}{|c|} \hline \square \\ \hline ab, ba \\ \hline \end{array}, \quad a, az \begin{array}{|c|} \hline \square \\ \hline 1, z \\ \hline \end{array}, \quad b, bz \begin{array}{|c|} \hline \square \\ \hline 1, z \\ \hline \end{array}, \quad ab, ba \begin{array}{|c|} \hline \square \\ \hline 1, z \\ \hline \end{array}, \quad a, az \begin{array}{|c|} \hline \square \\ \hline a, az \\ \hline \end{array}, \quad b, bz \begin{array}{|c|} \hline \square \\ \hline b, bz \\ \hline \end{array}, \quad ab, ba \begin{array}{|c|} \hline \square \\ \hline ab, ba \\ \hline \end{array}, \quad (3.3)$$

which contribute

$$\begin{aligned} & \frac{4}{|D_4|} \left[Z_{1,1} + Z_{1,\bar{a}} + Z_{1,\bar{b}} + Z_{1,\bar{a}\bar{b}} + Z_{\bar{a},1} + Z_{\bar{b},1} + Z_{\bar{a}\bar{b},1} + Z_{\bar{a},\bar{a}} + Z_{\bar{b},\bar{b}} + Z_{\bar{a}\bar{b},\bar{a}\bar{b}} \right] \\ & = Z \left([X/\mathbb{Z}_2 \times \mathbb{Z}_2] \prod [X/\mathbb{Z}_2 \times \mathbb{Z}_2]_{\text{d.t.}} \right). \end{aligned} \quad (3.4)$$

As there are two such sectors (one for each of $1, z$), we see that the three-dimensional effective theory decomposes into

$$\prod_2 \left([X/\mathbb{Z}_2 \times \mathbb{Z}_2] \prod [X/\mathbb{Z}_2 \times \mathbb{Z}_2]_{\text{d.t.}} \right) = \prod_2 [X/\mathbb{Z}_2 \times \mathbb{Z}_2] \prod_2 [X/\mathbb{Z}_2 \times \mathbb{Z}_2]_{\text{d.t.}}. \quad (3.5)$$

As observed earlier, this construction reproduces the decomposition prediction of section 2.

3.2 First nonbanded example

Consider the orbifold $[X/\mathbb{H}]$, where \mathbb{H} is the eight-element group of unit quaternions $\{\pm 1, \pm i, \pm j, \pm k\}$, with trivially-acting subgroup $\langle i \rangle \cong \mathbb{Z}_4$, as in [2, section 2.0.4], [1, section 5.4.1]. The group \mathbb{H} has conjugacy classes

$$\{1\}, \quad \{-1\}, \quad \{\pm i\}, \quad \{\pm j\}, \quad \{\pm k\}. \quad (3.6)$$

We construct the partition function on a torus by listing results for group elements along the third S^1 :

- ± 1 : in each of these two sectors, all commuting pairs in \mathbb{H} contribute:

$$\pm 1, \pm i \begin{array}{|c|} \hline \square \\ \hline \pm 1, \pm i \\ \hline \end{array}, \quad \pm 1 \begin{array}{|c|} \hline \square \\ \hline \pm j, \pm k \\ \hline \end{array}, \quad \pm j, \pm k \begin{array}{|c|} \hline \square \\ \hline \pm 1 \\ \hline \end{array}, \quad \pm j \begin{array}{|c|} \hline \square \\ \hline \pm j \\ \hline \end{array}, \quad \pm k \begin{array}{|c|} \hline \square \\ \hline \pm k \\ \hline \end{array} \quad (3.7)$$

which contribute

$$\frac{1}{|\mathbb{H}|} [(16)Z_{1,1} + (8)Z_{1,\xi} + (8)Z_{\xi,1} + (2)(4)Z_{\xi,xi}] = Z \left(X \prod_2 [X/\mathbb{Z}_2] \right), \quad (3.8)$$

where ξ represents the effectively-acting \mathbb{Z}_2 , matching the usual result for decomposition in the corresponding two-dimensional orbifold.

- $\{\pm i\}$: in this sector, the allowed commuting pairs are

$$\begin{matrix} \pm 1, \pm i & \square \\ & \pm 1, \pm i \end{matrix}, \tag{3.9}$$

which contribute

$$\frac{1}{|C(i)|} [(16)Z_{1,1}] = (4)Z(X) = Z\left(\coprod_4 X\right). \tag{3.10}$$

There are no contributions from $\{\pm j\}$, $\{\pm k\}$ along the third S^1 , as they do not act trivially on X . Altogether, the results from the $\{1\}$, $\{-1\}$, $\{\pm i\}$ sectors are consistent with a decomposition into

$$\coprod_6 X \coprod_4 [X/\mathbb{Z}_2]. \tag{3.11}$$

As observed earlier, this necessarily reproduces the prediction of section 2.

Note as a quick consistency check that if we think of each X as a double cover of $[X/\mathbb{Z}_2]$, then this is locally

$$(6)(2) + 4 = 16 = 4^2 = |\mathbb{Z}_4|^2 \tag{3.12}$$

copies of $[X/\mathbb{Z}_2]$, as expected.

3.3 Second nonbanded example

Next we consider $[X/A_4]$, where A_4 is the 12-element alternating group on four indeterminates, with trivially-acting normal subgroup $\mathbb{Z}_2 \times \mathbb{Z}_2$, as in [2, section 2.0.5], [1, section 5.5]. We can write

$$\mathbb{Z}_2 \times \mathbb{Z}_2 = \{1, \alpha, \beta, \gamma\}, \tag{3.13}$$

where

$$\alpha = (14)(23), \quad \beta = (13)(24), \quad \gamma = (12)(34), \tag{3.14}$$

and all the elements of A_4 , arranged into $A_4/\mathbb{Z}_2 \times \mathbb{Z}_2 = \mathbb{Z}_3$ cosets, are

$$A_4 = \{1, \alpha, \beta, \gamma, \tag{3.15}$$

$$(123), (142), (243), (134), \tag{3.16}$$

$$(132), (124), (234), (143)\}. \tag{3.17}$$

The conjugacy classes in A_4 are

$$\{1\}, \quad \{\alpha, \beta, \gamma\}, \quad \{(123), (142), (243), (134)\}, \quad \{(132), (124), (234), (143)\}. \tag{3.18}$$

As before, we enumerate contributions to a T^2 partition function from each sector defined by a loop around the third S^1 with a trivially-acting group element (representing a conjugacy class) inserted.

- 1: first, we consider the case that the identity is inserted along the third S^1 . Then, we simply count commuting pairs in A_4 , which are

$$\begin{aligned}
 & \begin{matrix} 1, \alpha, \beta, \gamma \\ 1, \alpha, \beta, \gamma \end{matrix} \begin{matrix} \square \\ \square \end{matrix}, \quad \begin{matrix} 1 \\ (123), \dots \end{matrix} \begin{matrix} \square \\ \square \end{matrix}, \quad \begin{matrix} 1 \\ (132), \dots \end{matrix} \begin{matrix} \square \\ \square \end{matrix}, \quad \begin{matrix} (123), \dots \\ 1 \end{matrix} \begin{matrix} \square \\ \square \end{matrix}, \quad \begin{matrix} (132), \dots \\ 1 \end{matrix} \begin{matrix} \square \\ \square \end{matrix}, \quad (3.19) \\
 & \begin{matrix} (123), (132) \\ (123), (132) \end{matrix} \begin{matrix} \square \\ \square \end{matrix}, \quad \begin{matrix} (142), (124) \\ (142), (124) \end{matrix} \begin{matrix} \square \\ \square \end{matrix}, \quad \begin{matrix} (243), (234) \\ (243), (234) \end{matrix} \begin{matrix} \square \\ \square \end{matrix}, \quad \begin{matrix} (134), (143) \\ (134), (143) \end{matrix} \begin{matrix} \square \\ \square \end{matrix} \quad (3.20)
 \end{aligned}$$

which contribute

$$\begin{aligned}
 & \frac{1}{|A_4|} [(16)Z_{1,1} + (4)Z_{1,\xi} + (4)Z_{1,\xi^2} + (4)Z_{\xi,1} + (4)Z_{\xi^2,1} \\
 & + (4)Z_{\xi,\xi} + (4)Z_{\xi,\xi^2} + (4)Z_{\xi^2,\xi} + (4)Z_{\xi^2,\xi^2}] \\
 & = Z(X \amalg [X/\mathbb{Z}_3]), \quad (3.21)
 \end{aligned}$$

where ξ denotes the generator of the effectively-acting $\mathbb{Z}_3 = A_4/\mathbb{Z}_2 \times \mathbb{Z}_2$, as expected from decomposition in the corresponding two-dimensional orbifold [1, section 5.5].

- $\{\alpha, \beta, \gamma\}$: in this sector, the conjugacy class represented by α , the commuting pairs which also commute with α are

$$\begin{matrix} 1, \alpha, \beta, \gamma \\ 1, \alpha, \beta, \gamma \end{matrix} \begin{matrix} \square \\ \square \end{matrix}, \quad (3.22)$$

which contribute

$$\frac{1}{|C(\alpha)|} [(16)Z_{1,1}] = (4)Z(X). \quad (3.23)$$

Putting this together, the sum of the results for the $\{1\}, \{\alpha, \beta, \gamma\}$ sectors are consistent with a decomposition into

$$\left(\amalg_5 X \right) \amalg [X/\mathbb{Z}_3]. \quad (3.24)$$

As observed earlier, this reproduces the decomposition prediction of section 2.

As a consistency check, note that if we interpret X as a triple cover of $[X/\mathbb{Z}_3]$, then locally this is

$$(5)(3) + 1 = 16 = |\mathbb{Z}_2 \times \mathbb{Z}_2|^2 \quad (3.25)$$

copies of $[X/\mathbb{Z}_3]$.

3.4 Third nonbanded example

Next, consider the orbifold $[X/D_n]$, with trivially-acting $\mathbb{Z}_n \subset D_n$, as in [1, section 5.6]. Here, D_n denotes the $2n$ -element dihedral group, generated by a, b , where

$$a^2 = 1, \quad b^n = 1, \quad aba = b^{-1}. \quad (3.26)$$

The trivially-acting subgroup $\mathbb{Z}_n = \langle b \rangle$, and $D_n/\mathbb{Z}_n = \mathbb{Z}_2$. In the special case that n is even, D_n has a \mathbb{Z}_2 center, generated by $z = b^{n/2}$. For completeness, the elements of D_n can be enumerated as

$$D_n = \{1, b, b^2, \dots, b^{n-1}, a, ab, \dots, ab^{n-1}\}. \quad (3.27)$$

If n is even, then D_n has

$$\frac{n}{2} + 3 \tag{3.28}$$

conjugacy classes, which are, explicitly

$$\{1\}, \quad \{a, ab^2, ab^4, \dots, ab^{n-2}\}, \quad \{ab, ab^3, ab^5, \dots, ab^{n-1}\}, \tag{3.29}$$

$$\{b^j, b^{-j}\} \quad \text{for } 1 \leq j \leq n/2. \tag{3.30}$$

If n is odd, then D_n has

$$\frac{n+3}{2} \tag{3.31}$$

conjugacy classes, which are, explicitly

$$\{1\}, \quad \{a, ab, ab^2, ab^3, ab^4, \dots, ab^{n-1}\}, \tag{3.32}$$

$$\{b^j, b^{-j}\} \quad \text{for } 1 \leq j \leq \frac{n-1}{2}. \tag{3.33}$$

As before, we consider contributions to a T^2 partition function from sectors with trivially-acting conjugacy classes on the third S^1 .

- 1: in this sector, all commuting pairs in D_n contribute. For n odd, these are

$$\begin{matrix} \langle b \rangle & \square & 1 & \square & a, ab, \dots & \square & ab^i & \square \\ \langle b \rangle & & a, ab, \dots & & 1 & & ab^i & \end{matrix} \tag{3.34}$$

which contribute

$$\frac{1}{|D_n|} \left[(n^2)Z_{1,1} + (n)Z_{1,\xi} + (n)Z_{\xi,1} + (n)Z_{\xi,xi} \right] \tag{3.35}$$

$$= \frac{1}{2n} \left[(n^2 - n)Z_{1,1} + (n)(Z_{1,1} + Z_{1,\xi} + Z_{\xi,1} + Z_{\xi,\xi}) \right], \tag{3.36}$$

$$= Z \left(\prod_{(n-1)/2} X \prod_1 [X/\mathbb{Z}_2] \right), \tag{3.37}$$

where ξ generates the effectively-acting $\mathbb{Z}_2 = D_n/\mathbb{Z}_n$.

For n even, in addition to the commuting pairs above, there are also

$$\begin{matrix} z & \square & a, ab, \dots & \square & ab^{i+n/2} & \square \\ & a, ab, \dots & z & & ab^i & \end{matrix} \tag{3.38}$$

and the total contribution becomes

$$\frac{1}{|D_n|} \left[(n^2)Z_{1,1} + (2n)Z_{1,\xi} + (2n)Z_{\xi,1} + (2n)Z_{\xi,xi} \right] \tag{3.39}$$

$$= \frac{1}{2n} \left[(n^2 - 2n)Z_{1,1} + (2n)(Z_{1,1} + Z_{1,\xi} + Z_{\xi,1} + Z_{\xi,\xi}) \right], \tag{3.40}$$

$$= Z \left(\prod_{(n-2)/2} X \prod_2 [X/\mathbb{Z}_2] \right). \tag{3.41}$$

- z : if n is even, then $z = b^{n/2}$ is in the center of D_n and defines its own conjugacy class. In this sector, there are the same contributions as for the 1 sector above, and so we get the contribution

$$Z \left(\prod_{(n-2)/2} X \prod_2 [X/\mathbb{Z}_2] \right). \tag{3.42}$$

- $\{b^i, b^{-i}\}, i \neq 0, n/2$: in these sectors, there are contributions from the commuting pairs

$$\langle b \rangle \square_{\langle b \rangle}, \tag{3.43}$$

which contribute

$$\frac{1}{|C(b^i)|} (n^2 Z_{1,1}) = nZ(X) = Z \left(\prod_n X \right). \tag{3.44}$$

We summarize the total contribution as follows. If n is odd, the results are consistent with a decomposition into

$$\prod_{(n+1)(n-1)/2} X \prod_1 [X/\mathbb{Z}_2]. \tag{3.45}$$

As observed earlier, this reproduces the decomposition prediction of section 2.

As a consistency check, if we interpret X as a double cover of $[X/\mathbb{Z}_2]$, then this is locally

$$(2) \frac{(n+1)(n-1)}{2} + 1 = n^2 = |\mathbb{Z}_n|^2 \tag{3.46}$$

copies of $[X/\mathbb{Z}_2]$.

If n is even, the results are consistent with a decomposition into

$$\prod_{(n-2)(n+2)/2} X \prod_4 [X/\mathbb{Z}_2]. \tag{3.47}$$

As a consistency check, if we interpret X as a double cover of $[X/\mathbb{Z}_2]$, then this is locally

$$(2) \frac{(n-2)(n+2)}{2} + 4 = n^2 = |\mathbb{Z}_n|^2 \tag{3.48}$$

copies of $[X/\mathbb{Z}_2]$.

As observed earlier, for n both even and odd, this reproduces the decomposition prediction of section 2.

3.5 First trivially-acting nonabelian group example

Next, consider the orbifold $[X/S_3]$, where all of S_3 acts trivially. We enumerate the elements of S_3 as

$$S_3 = \{1, (12), (23), (13), (123), (132) = (123)^2\}. \tag{3.49}$$

The conjugacy classes are

$$\{1\}, \quad \{(12), (23), (13)\}, \quad \{(123), (132)\}. \tag{3.50}$$

As before, we itemize contributions to a T^2 partition function by sectors corresponding to conjugacy classes of trivially-acting group elements on the third S^1 . For each conjugacy class, we pick one representative below.

- 1: in this sector, $C(1) = S_3$, so all commuting pairs in S_3 contribute. These are

$$1 \begin{array}{|c|} \hline \square \\ \hline 1 \end{array}, \quad 1 \begin{array}{|c|} \hline \square \\ \hline g \neq 1 \end{array}, \quad g \neq 1 \begin{array}{|c|} \hline \square \\ \hline 1 \end{array}, \quad \begin{array}{|c|} \hline (12) \square \\ \hline (12) \end{array}, \quad \begin{array}{|c|} \hline (23) \square \\ \hline (23) \end{array}, \quad \begin{array}{|c|} \hline (13) \square \\ \hline (13) \end{array}, \quad \begin{array}{|c|} \hline (123), (132) \square \\ \hline (123), (132) \end{array} \quad (3.51)$$

which contribute

$$\frac{1}{|S_3|} [1 + 5 + 5 + 1 + 1 + 1 + 4] Z_{1,1} = (3)Z(X), \quad (3.52)$$

consistent with the prediction of decomposition in the corresponding two-dimensional orbifold (as S_3 has three irreducible representations).

- (12): in this sector, $C((12)) = \{1, (12)\}$, so the pertinent commuting pairs are

$$\begin{array}{|c|} \hline 1, (12) \square \\ \hline 1, (12) \end{array} \quad (3.53)$$

which contribute

$$\frac{1}{|C((12))|} (4) Z_{1,1} = (2)Z(X). \quad (3.54)$$

- $\{(123), (132)\}$: in this sector, $C((123)) = \{1, (123), (132)\}$, and the pertinent commuting pairs are

$$\begin{array}{|c|} \hline 1, (123), (132) \square \\ \hline 1, (123), (132) \end{array} \quad (3.55)$$

which contribute

$$\frac{1}{|C((123))|} (9) Z_{1,1} = (3)Z(X). \quad (3.56)$$

Altogether, adding the contributions from the different sectors gives $(8)Z(X)$, which is consistent with a decomposition into

$$\coprod_8 X. \quad (3.57)$$

As observed earlier, this reproduces the decomposition prediction of section 2.

3.6 Second trivially-acting nonabelian group example

Next, consider the orbifold $[X/\mathbb{H}]$, where all of $\mathbb{H} = \{\pm 1, \pm i, \pm j, \pm k\}$ acts trivially. The conjugacy classes are

$$\{1\}, \quad \{-1\}, \quad \{\pm i\}, \quad \{\pm j\}, \quad \{\pm k\}. \quad (3.58)$$

As before, we itemize contributions to a T^2 partition function by sectors corresponding to group elements representing trivially-acting conjugacy classes on the third S^1 :

- ± 1 : in these two sectors, all commuting pairs in \mathbb{H} contribute. These are

$$\begin{array}{|c|} \hline \pm 1 \square \\ \hline \pm 1 \end{array}, \quad \begin{array}{|c|} \hline \pm 1 \square \\ \hline g \neq \pm 1 \end{array}, \quad g \neq \pm 1 \begin{array}{|c|} \hline \square \\ \hline \pm 1 \end{array}, \quad \begin{array}{|c|} \hline \pm i \square \\ \hline \pm 1 \end{array}, \quad \begin{array}{|c|} \hline \pm j \square \\ \hline \pm j \end{array}, \quad \begin{array}{|c|} \hline \pm k \square \\ \hline \pm k \end{array}, \quad (3.59)$$

which contribute

$$\frac{1}{|\mathbb{H}|} (4 + (2)(6) + (2)(6) + 4 + 4 + 4) Z_{1,1} = (5)Z(X), \quad (3.60)$$

consistent with the prediction of decomposition in the corresponding two-dimensional orbifold (as \mathbb{H} has five irreducible representations).

- $\{\pm i\}$: in this sector, the pertinent commuting pairs are

$$\pm 1, \pm i \begin{array}{|c|} \hline \square \\ \hline \end{array}, \quad (3.61)$$

which contribute

$$\frac{1}{|C(i)|} (4^2) Z_{1,1} = \frac{16}{4} Z(X) = (4) Z(X). \quad (3.62)$$

- $\{\pm j\}, \{\pm k\}$: these two sectors each give the same results as the $\{\pm i\}$ sector.

Altogether, adding the contributions from the different sectors gives

$$(22) Z(X),$$

which is consistent with a decomposition into

$$\coprod_{22} X. \quad (3.63)$$

As observed earlier, this reproduces the decomposition prediction of section 2.

3.7 Third trivially-acting nonabelian group example

Next, consider the orbifold $[X/D_4]$, where all of the eight-element dihedral group D_4 acts trivially. The conjugacy classes of D_4 are

$$\{1\}, \quad \{z\}, \quad \{a, az\}, \quad \{b, bz\}, \quad \{ab, ba\}. \quad (3.64)$$

As before, we itemize contributions to a T^2 partition function by sectors corresponding to trivially-acting group elements on the third S^1 :

- $1, z$: in these two sectors, all commuting pairs in D_4 contribute. These are

$$1, z \begin{array}{|c|} \hline \square \\ \hline \end{array}, \quad 1, z \begin{array}{|c|} \hline \square \\ \hline \end{array}, \quad g \neq 1, z \begin{array}{|c|} \hline \square \\ \hline \end{array}, \quad a, az \begin{array}{|c|} \hline \square \\ \hline \end{array}, \quad b, bz \begin{array}{|c|} \hline \square \\ \hline \end{array}, \quad ab, ba \begin{array}{|c|} \hline \square \\ \hline \end{array}, \quad (3.65)$$

which contribute

$$\frac{1}{|D_4|} [4 + (2)(6) + (2)(6) + 4 + 4 + 4] Z_{1,1} = (5) Z(X), \quad (3.66)$$

consistent with the prediction of decomposition in the corresponding two-dimensional orbifold (as D_4 has five irreducible representations).

- $\{a, az\}$: in this sector, the pertinent commuting pairs are

$$1, z \begin{array}{|c|} \hline \square \\ \hline \end{array}, \quad 1, z \begin{array}{|c|} \hline \square \\ \hline \end{array}, \quad a, az \begin{array}{|c|} \hline \square \\ \hline \end{array}, \quad a, az \begin{array}{|c|} \hline \square \\ \hline \end{array}, \quad (3.67)$$

which contribute

$$\frac{1}{|C(a)|} [4 + 4 + 4 + 4] Z_{1,1} = (4) Z(X). \quad (3.68)$$

- $\{b, bz\}, \{ab, ba\}$: contributions from each of these two sectors follow the same form as those from the $\{a, az\}$ sector.

Altogether, adding the contributions from the different sectors gives $(22) Z(X)$, which is consistent with a decomposition into

$$\coprod_{22} X. \quad (3.69)$$

As observed earlier, this reproduces the decomposition prediction of section 2.

4 Quantum K theory via GLSM computations

In this section, we turn to a different class of presentations, namely gauged linear sigma models, rather than orbifolds, and verify the structure predicted in section 2. Here, the trivially-acting subgroup K of the gauge group will always be a subset of the center. Furthermore, in all subsections except 4.6, the two-dimensional theory will have both a BK (one-form) and K (zero-form) symmetry, without a 't Hooft anomaly between them. As a result, in all subsections except 4.6, we will see a decomposition of the effective two-dimensional theory into $|K|$ copies of a two-dimensional GLSM with a one-form symmetry, each copy of which again separately decomposes, giving altogether a decomposition into a total of $|K|^2$ universes, as originally predicted for such cases in [16, 17]. The first level of decomposition — the choice of Wilson line — is visible in the fact that the roots are symmetric under multiplication by ℓ th root of unity. As the Coulomb branch parameters are the Wilson lines, this is precisely a symmetry between possible choices of Wilson line. Furthermore, by using GLSM methods, we are able to make predictions for the quantum K theory rings, not just the structure of the decomposition.

In section 4.6, we consider more general Chern-Simons levels and 't Hooft anomalies, and discuss how the decomposition story is modified.

We should emphasize that, as is typical in such computations, we are making mathematical conjectures via an interpretation of the Coulomb branch equations, and a proper treatment of the mathematics should await a more rigorous mathematical analysis of the quantum K theory. Our point, however, is that an interpretation of the Coulomb branch equations consistent with expectations from decomposition is possible.

4.1 Warmup: gerbes on projective spaces

4.1.1 General \mathbb{Z}_ℓ gerbes on projective spaces

We begin with general \mathbb{Z}_ℓ gerbes on \mathbb{P}^{n-1} , reviewing examples discussed in [16, section 4]. Following e.g. [4], such gerbes can be described by a $U(1)^2$ GLSM with fields x_i, z , and charges

$$\begin{array}{cc} x_i & z \\ 1 & -m \\ 0 & \ell \end{array}$$

As discussed in [4], this defines a \mathbb{Z}_ℓ gerbe over \mathbb{P}^{n-1} , with characteristic class $-m \pmod{\ell}$. The weighted projective space $\mathbb{P}_{[\ell, \dots, \ell]}^{n-1}$ is equivalent to the case $m = 1$.

Before computing the quantum K theory, we begin by reviewing the quantum cohomology. Following [4, section 3.2], the quantum cohomology ring computed from the two-dimensional GLSM above is

$$\mathbb{C}[\psi_1, \psi_2] / \langle \psi_1^n - \psi_2^m q_1, \psi_2^\ell - q_2 \rangle. \tag{4.1}$$

(This is computed using GLSM Coulomb branch methods, as the critical locus of the one-loop twisted effective superpotential given in e.g. [73].) In decomposition, we interpret the ψ_2 as indexing ℓ different universes, each a copy of the ordinary supersymmetric \mathbb{P}^{n-1} model with q 's in different universes slightly different, related by shifts of the B field.

Let us next compute the quantum K theory of these examples. Physically, this is the OPE ring of Wilson lines in a three-dimensional gauge theory on a 3-manifold of the form $S^1 \times \Sigma$, for Σ a Riemann surface, where the Wilson lines are wrapped on S^1 and move in parallel along Σ [43, 47–49]. This also can be computed from the critical locus of a twisted one-loop effective superpotential along the Coulomb branch, albeit in a two-dimensional gauge theory arising from a regularized Kaluza-Klein reduction of a three-dimensional gauge theory [44–46, 51–58].

The two-dimensional theory has an infinite tower of fields. For each field in the three-dimensional theory, transforming in representation R of the gauge group G , there is an infinite tower of massive fields in the two-dimensional theory, each also transforming in the same representation R of the gauge group. As a result, if the three-dimensional gauge theory has a one-form symmetry due to some subgroup of the gauge group acting trivially on all the matter, the two-dimensional theory arising from the Kaluza-Klein reduction will have the same one-form symmetry, a fact we shall use throughout this paper.

The twisted one-loop effective superpotential arising from those Kaluza-Klein towers is then regularized. Following [55, equ'n (2.1)], [46, equ'n (2.33)], [45, section 2.2.2], [50, section 2.2.1], [52, section 2], [53, equ'n (2.10)], [54, equ'n (2.2)], the regularized superpotential has the form

$$W = \frac{1}{2}k^{ab}(\ln X_a)(\ln X_b) + \sum_a (\ln q_a)(\ln X_a) + \sum_i \left[\text{Li}_2(X^{\rho_i}) + \frac{1}{4}(\ln(X^{\rho_i}))^2 \right], \quad (4.2)$$

where

$$X^{\rho_i} = \prod_a X_a^{Q_a^i}, \quad (4.3)$$

so that

$$X^x = X_1, \quad X^z = X_1^{-m} X_2^\ell, \quad (4.4)$$

and

$$k^{11} = -\frac{1}{2}(n + m^2), \quad k^{12} = k^{21} = \frac{1}{2}m\ell, \quad k^{22} = -\frac{1}{2}\ell^2, \quad (4.5)$$

using the general formula for Chern-Simons levels [57, equ'n (2.6)]

$$k^{ab} = \frac{1}{2} \sum_i (R_i - 1) Q_i^a Q_i^b, \quad (4.6)$$

as given by $U(1)_{-1/2}$ quantization, see e.g. [52, section 2.2], [55], as that is the choice that reproduces ordinary quantum K-theory.

Computing the critical locus, we find the quantum K theory ring relations

$$(1 - X_1)^n (1 - X_1^{-m} X_2^\ell)^{-m} = q_1, \quad (4.7)$$

$$(1 - X_1^{-m} X_2^\ell)^\ell = q_2. \quad (4.8)$$

In passing, note that shifting $m \mapsto m + \ell$ is equivalent to changing $q_1 \mapsto q_1 q_2^{-1}$. It is in this sense that the quantum K theory ring only depends upon the characteristic class of the gerbe, $m \bmod \ell$.

Define

$$y = 1 - X_1^{-m} X_2^\ell, \tag{4.9}$$

then we can write the ring relations above as

$$(1 - X_1)^n = q_1 y^m, \tag{4.10}$$

$$y^\ell = q_2. \tag{4.11}$$

As y is determined by X_2^ℓ , there is an ℓ -fold phase choice in solutions of X_2 for fixed y , which we interpret as reflecting ℓ copies of a theory, each copy of which itself decomposes into ℓ universes, for altogether ℓ^2 copies of the quantum K theory ring of the underlying space \mathbb{P}^{n-1} (from the first relation), indexed by the ℓ^2 values of y and X_2 .

4.1.2 Special case: weighted projective spaces

Next, we specialize to gerbes which can be presented as weighted projective spaces, a case previously discussed in [16, section 4.1]. These admit an additional, different, UV presentation, so it will be instructive to compute the quantum K theory ring and compare to the general case, reviewing the result of [16, section 4.1].

For a general stacky weighted projective space $\mathbb{P}_{[w_0, \dots, w_{n-1}]}^{n-1}$, following the prescription in e.g. [55, section 2], the quantum K theory ring relations arise as the critical locus of the superpotential

$$W = \frac{k}{2} (\ln X)^2 + (\ln q)(\ln X) + \sum_i \left[\text{Li}_2(X^{w_i}) + \frac{1}{4} (\ln(X^{w_i}))^2 \right], \tag{4.12}$$

with Chern-Simons level

$$k = -\frac{1}{2} \sum_i (w_i)^2. \tag{4.13}$$

It is straightforward to compute that the critical locus is given by

$$\prod_{j=0}^{n-1} (1 - X^{w_j})^{w_j} = q. \tag{4.14}$$

This result matches results for quantum K theory rings for weighted projective stacks given in [74, thm. 1.5, cor. 5.6], namely that for a weighted projective space $\mathbb{P}_{[w_0, \dots, w_{n-1}]}^{n-1}$. (See also [75, example 1.3], and [16] for a physics discussion.)

Now, \mathbb{Z}_ℓ gerbes on \mathbb{P}^{n-1} of characteristic class $-1 \pmod{\ell}$ are described as stacky weighted projective spaces $\mathbb{P}_{[\ell, \ell, \dots, \ell]}^{n-1}$, meaning that all weights w_i are equal to ℓ . (For a discussion of how the resulting physical theory in two dimensions differs from that of the ordinary supersymmetric \mathbb{P}^{n-1} model, see [2–4].)

In the case that the weights are all ℓ , the result above for the quantum K theory ring reduces to

$$(1 - X^\ell)^{\ell n} = q, \tag{4.15}$$

As a consistency check, write $X = \exp(-2\pi R\sigma)$, $q = R^{\ell n} q_{2d}$, then the relation (4.15) reduces to

$$\sigma^{\ell n} \propto q_{2d}, \tag{4.16}$$

which is the same quantum cohomology ring relation for $\mathbb{P}_{[\ell, \dots, \ell]}^{n-1}$ discussed in e.g. [2–4].

To help explain (4.15), consider the case that $\ell = 1$, describing an ordinary projective space \mathbb{P}^{n-1} . In this case, we interpret $X = \mathcal{O}(-1)$, then $1 - X = \mathcal{O}_H$, the class of a hyperplane. The product corresponds to generic intersection, and the intersection of n hyperplanes in general position in \mathbb{P}^{n-1} is empty, so that classically, $(1 - X)^n = 0$ in $K(\mathbb{P}^{n-1})$.

In the case $\ell > 1$, our understanding of [74, 76, 77] is that to interpret (4.15), one should interpret $X = \mathcal{O}(-1/\ell)$, a line bundle on the gerbe which is not a pullback from the underlying projective space, and that one uses an orbifold product on the inertia stack, a K-theoretic analogue of products in orbifold cohomology.

In terms of decomposition, taking an ℓ th root of (4.15), we get ℓ copies of the ordinary quantum K theory ring of \mathbb{P}^{n-1} , generated by X^ℓ , indexed by ℓ th roots of unity. The fact that the generator is X^ℓ can be interpreted as meaning we get the decomposition above for each of the ℓ roots of X^ℓ , giving a decomposition into a total of ℓ^2 universes, matching expectations. As a consistency check, we can compare to the ring relations for general \mathbb{Z}_ℓ gerbes as follows. Identify $q = q_1^\ell q_2^m$ and $X_1 = X^\ell$, then the ℓ th power of the relation (4.10) can be written

$$(1 - X^\ell)^{\ell n} = q. \tag{4.17}$$

which, for generator X^ℓ , is the ring relation for $m = 1$, matching (4.15).

In any event, again we see a decomposition in quantum K theory, as predicted.

4.2 General \mathbb{Z}_ℓ gerbes over Grassmannians

In this section we will discuss \mathbb{Z}_ℓ gerbes over Grassmannians, constructing them as \mathbb{C}^\times quotients of line bundles over ordinary Grassmannians, exactly as we reviewed for projective spaces in section 4.1.

It will be handy to recall that irreducible representations of $u(k)$ are characterized by a k -tuple of ordered integers

$$[\lambda_1, \lambda_2, \dots, \lambda_k], \quad \lambda_i \in \mathbb{Z} \tag{4.18}$$

with $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_k$. In this description, the corresponding $su(k)$ representation is

$$(\lambda_1 - \lambda_k, \lambda_2 - \lambda_k, \dots, \lambda_{k-1} - \lambda_k), \tag{4.19}$$

where the $\det u(k) = u(1)$ charge is given by $\lambda_1 + \lambda_2 + \dots + \lambda_k$.

4.2.1 Quick review of ordinary Grassmannians

Before describing gerbes on Grassmannians, and their quantum cohomology and quantum K theory, let us first quickly review ordinary Grassmannians, their quantum cohomology and quantum K theory.

First, an ordinary Grassmannian $G(k, n)$ can be constructed as $\mathbb{C}^{kn}/\mathrm{GL}(k)$, meaning a $U(k)$ GLSM with n chiral multiplets in the fundamental representation [78]. As a Fano space realized by a GLSM, both its quantum cohomology and quantum K theory can be computed using Coulomb branch methods.

Recall that for an ordinary Grassmannian $G(k, n)$, the nonequivariant relations in the quantum cohomology ring are

$$(-1)^{k-1}q = \sigma_a^n, \tag{4.20}$$

where $\{\sigma_a\}$ ($a \in \{1, \dots, k\}$) are local Coulomb branch coordinates on a Weyl-group orbifold, or more formally, Chern roots of the universal subbundle S . After symmetrization, these give rise to the relation

$$c(S)c(Q) = 1 + (-1)^{n-k}q. \tag{4.21}$$

This is the quantum cohomology ring relation (see e.g. [78]).

There is a similar description⁴ of the quantum K theory ring of a Grassmannian $G(k, n)$, derived again from a twisted one-loop effective superpotential. The Coulomb branch equations one derives are (see e.g. [55, equ'n (2.40)])

$$(-1)^{k-1}qX_a^k = (\det X)(1 - X_a)^n, \tag{4.22}$$

(where the $X_a = \exp(-2\pi R\sigma_a)$ for R the radius of the S^1 in the 3-manifold,) which after symmetrization become the quantum K-theory ring relation [56, theorem 1.1]

$$\lambda_y(S) \star \lambda_y(\mathbb{C}^n/S) = \lambda_y(\mathbb{C}^n) - y^{n-k} \frac{q}{1-q} \det(\mathbb{C}^n/S) \star (\lambda_y(S) - 1). \tag{4.23}$$

For the purposes of comparing to decomposition predictions, it will mostly be sufficient for our purposes to work only with Coulomb branch expressions.

4.2.2 Description of \mathbb{Z}_ℓ gerbes over Grassmannians

Mathematically, we can describe \mathbb{Z}_ℓ gerbes over Grassmannians $G(k, n)$ as quotients

$$[L^*/\mathbb{C}^*], \tag{4.24}$$

where $L \rightarrow G(k, n)$ is a line bundle, L^* is L minus its zero section, and \mathbb{C}^* acts with a trivially-acting \mathbb{Z}_ℓ subgroup. The resulting \mathbb{Z}_ℓ gerbe has characteristic class $c_1(L) \bmod \ell$.

In physics, we describe such gerbes as⁵ a $U(k) \times U(1)$ gauge theory with n chiral superfields in the fundamental representation of $U(k)$ and neutral under $U(1)$, together with another chiral superfield in the $U(k)$ representation $[-m', -m', \dots, -m']$ (equivalently, the charge $-m'k$ representation of $\det U(k)$) and of charge ℓ under the $U(1)$. This describes a \mathbb{Z}_ℓ gerbe on $G(k, n)$ of characteristic class $-km' \bmod \ell$.

⁴Quantum K theory of Grassmannians has been extensively discussed, see e.g. [44, 49, 54, 55, 79–82] for a few examples in both the mathematics and physics literatures.

⁵We would like to thank W. Gu for useful discussions.

4.2.3 Quantum cohomology

Using standard methods [73], the quantum cohomology ring of this Grassmannian is given by

$$\sigma_a^n \left(\ell \sigma_0 - m' \sum_b \sigma_b \right)^{-m'} = (-)^{k-1} \tilde{q}_1, \tag{4.25}$$

$$\left(\ell \sigma_0 - m' \sum_b \sigma_b \right)^\ell = \tilde{q}_0. \tag{4.26}$$

Define

$$\Upsilon = \ell \sigma_0 - m' \sum_b \sigma_b. \tag{4.27}$$

Rescaling $\tilde{q}_0 \rightarrow 1$ without⁶ loss of generality, we can write these equations as

$$\sigma_a^n = (-)^{k-1} \tilde{q}_1 \Upsilon^{m'}, \quad \Upsilon^\ell = 1. \tag{4.28}$$

This is precisely as expected for the quantum cohomology ring of a disjoint union of ℓ copies of $G(k, n)$, each with B fields / theta angles slightly shifted (as encoded in ℓ th roots of unity), the same pattern discussed in [4] for quantum cohomology rings of toric gerbes on projective spaces. (For completeness, we also mention that the same structure is visible in nonabelian mirror constructions [83].)

4.2.4 Quantum K theory

Next, we compute the quantum K theory ring relations. Using the results reviewed in section 4.1, we can write the effective twisted superpotential as follows,

$$\begin{aligned} W = & \frac{k}{2} \sum_{a=1}^k (\ln X_a)^2 - \frac{1}{2} \left(\sum_{a=1}^k \ln X_a \right)^2 + (\ln(-1)^{k-1} q_1) \sum_{a=1}^k \ln X_a \\ & + (\ln q_0)(\ln X_0) + n \sum_{a=1}^k \text{Li}_2(X_a) + \text{Li}_2((\det X)^{-m'} X_0^\ell). \end{aligned} \tag{4.29}$$

Taking the critical locus,

$$X_0 \frac{\partial W}{\partial X_0} = 0, \quad X_a \frac{\partial W}{\partial X_a} = 0, \tag{4.30}$$

we obtain,

$$q_0 = \left(1 - (\det X)^{-m'} X_0^\ell \right)^\ell, \tag{4.31}$$

$$(-1)^{k-1} q_1 X_a^k = (\det X)(1 - X_a)^n \left(1 - (\det X)^{-m'} X_0^\ell \right)^{-m'}. \tag{4.32}$$

Furthermore, this implies

$$(-1)^{\ell(k-1)} q_0^{m'} q_1^\ell X_a^{k\ell} = (\det X)^\ell (1 - X_a)^{\ell n}. \tag{4.33}$$

⁶It can be absorbed into q_1 with suitable redefinitions.

As a consistency check, note that when $\ell = 1$, we obtain

$$(-1)^{k-1}(q_0^{m'} q_1) X_a^k = (\det X)(1 - X_a)^n, \tag{4.34}$$

which agrees with the relations for ordinary Grassmannian (4.22) if we make the identification $q_0^{m'} q_1 = q$.

Now, to compare to the claimed decomposition, define

$$y = 1 - (\det X)^{-m'} X_0^\ell. \tag{4.35}$$

Then, the quantum K-theory ring relations (4.32) become

$$q_0 = y^\ell, \tag{4.36}$$

$$(-1)^{k-1} q_1 y^{m'} X_a^k = (\det X)(1 - X_a)^n, \tag{4.37}$$

which, given the ℓ -fold ambiguity in X_0 for fixed y , is clearly a total of ℓ^2 copies of the quantum K theory ring relations of $G(k, n)$, each with shifted q , shifted by an ℓ -th root of unity, consistent with expectations.

4.3 Gerbes via weighted Grassmannians

In this section we will construct a theory describing a \mathbb{Z}_{km+1} gerbe over the Grassmannian, as an analogue of a weighted projective space, and analyze physics predictions for its quantum K theory. In principle this is just a different presentation of one of the \mathbb{Z}_ℓ gerbes of the previous section, but it will be an instructive check to consider it in detail.

4.3.1 Construction of the theory

First, following [78], we describe an ordinary Grassmannian $G(k, n)$ by a $3d \mathcal{N} = 2$ $U(k)$ gauge theory with n chiral superfields in the fundamental representation, meaning the representation with highest weight

$$[1, \underbrace{0, \dots, 0}_{k-1}]. \tag{4.38}$$

Now, to describe the weighted Grassmannian describing a \mathbb{Z}_{km+1} gerbe,⁷ consider a $3d \mathcal{N} = 2$ $U(k)$ gauge theory with n chiral superfields in the $U(k)$ representation with highest weight

$$[m + 1, \underbrace{m, \dots, m}_{k-1}], \tag{4.39}$$

meaning, a tensor product of the fundamental representation with a charge m representation of $\det U(k)$. We can understand why this describes a gerbe as follows. First, since the determinant of $U(k)$ is itself a product of k $U(1)$ factors, namely the diagonal in a $k \times k$ matrix representation of $U(k)$, there is a trivially-acting \mathbb{Z}_k subgroup of $U(k)$ acting on anything of charge 1 under $\det U(k)$, defined by diagonal matrices with k th roots of unity

⁷We would like to thank W. Gu for useful discussions.

along the diagonal. A charge m representation of $\det U(k)$ is therefore invariant under a \mathbb{Z}_{mk} subgroup of $U(k)$. The chiral multiplets here are in a charge m representation of $U(k)$ tensored with the fundamental representation of $U(k)$, which is invariant under a \mathbb{Z}_{km+1} subgroup of $U(k)$. Thus, this gauge theory has a trivially-acting \mathbb{Z}_{km+1} subgroup, hence the gauge theory has a \mathbb{Z}_{km+1} one-form symmetry, and so describes a \mathbb{Z}_{km+1} gerbe over $G(k, n)$.

4.3.2 Quantum cohomology

Using standard methods [73], it is straightforward to compute that the quantum cohomology ring is given by

$$(-1)^{k-1}q = \left(\sigma_a + m \sum_b \sigma_b \right)^n \prod_{b=1}^k \left(\sigma_b + m \sum_c \sigma_c \right)^{nm}. \quad (4.40)$$

In principle, the σ fields should couple to the bundle describing the ‘minimal’ action of the gauge group. If S denotes the universal subbundle on $G(k, n)$, and π the projection from the gerbe to $G(k, n)$, then we take the σ fields to couple to \tilde{S} defined by

$$\tilde{S} = \pi^* S \otimes (\pi^* \det S)^{-1/k} \otimes (\pi^* \det S)^{1/(k(km+1))}, \quad (4.41)$$

$$= \pi^* S \otimes (\pi^* \det S)^{-m/(km+1)}. \quad (4.42)$$

(Note that since \tilde{S} is of rank k , the $(k(km+1))$ th root is well-defined on a \mathbb{Z}_{km+1} gerbe.) We justify this identification by the fact that

$$\det \tilde{S} = (\det \pi^* S)^{1/(km+1)}. \quad (4.43)$$

This tells us that the σ fields are coupling to a generator, roughly speaking.

Now define $\tau_a = \sigma_a + m \sum_b \sigma_b$. In terms of τ_a , equation (4.40) becomes

$$(-1)^{k-1}q = \tau_a^n (\det \tau)^{nm}. \quad (4.44)$$

To interpret this result, we note that (4.44) implies

$$\left(\prod_a \tau_a^n \right) (\det \tau)^{knm} = (-)^{k(k-1)} q^k = q^k, \quad (4.45)$$

or more simply

$$(\det \tau)^{n(km+1)} = q^k, \quad (4.46)$$

hence

$$(\det \tau)^{nm} = \xi^m \left[q^k \right]^{m/(km+1)}, \quad (4.47)$$

for ξ a $(km+1)$ th root of unity. Then, we can rewrite (4.44) as

$$\tau_a^n = \xi^{-m} (-)^{k-1} q^{+1/(km+1)}. \quad (4.48)$$

To interpret equation (4.48), recall that for ordinary Grassmannian, the nonequivariant relations in the quantum cohomology ring are

$$(-1)^{k-1}q = \sigma_a^n, \quad (4.49)$$

which give rise to the relation

$$c(S) c(Q) = 1 + (-1)^{n-k} q. \tag{4.50}$$

Here, the τ_a correspond to the Chern roots of

$$\tilde{S} \otimes (\det \tilde{S})^m = \pi^* S. \tag{4.51}$$

Thus, we see that for the \mathbb{Z}_{km+1} gerbe on the Grassmannian, the relation (4.48) should be interpreted as

$$c(\pi^* S) c(\pi^* Q) = 1 + (-1)^{n-k} \xi^{-m} q^{+1/(km+1)}, \tag{4.52}$$

or more simply, $km + 1$ copies of the ordinary quantum cohomology ring of $G(k, n)$, with θ angle shifts (encoded in the roots of unity ξ). This is as expected from decomposition, and also correctly reduces to results for gerbes on projective spaces in the special case $k = 1$.

4.3.3 Quantum K theory

Applying the same methods discussed earlier, the Coulomb branch equation is given by

$$(-1)^{k-1} q X_a^k = (1 - X_a (\det X)^m)^n \prod_{b=1}^k [X_b (1 - X_b (\det X)^m)^{nm}]. \tag{4.53}$$

We can quickly check that this expression has correct specializations:

- When $k = 1$, this specializes to the relation for $\mathbb{P}_{[\ell, \dots, \ell]}^{n-1}$, a \mathbb{Z}_ℓ gerbe on the projective space \mathbb{P}^{n-1} ,

$$(1 - X^\ell)^{n\ell} = q, \quad \text{with } \ell = m + 1, \tag{4.54}$$

matching equation (4.15) earlier.

- When $m = 0$, this specializes to the relation (4.22) for ordinary Grassmannians.

Now, let us work out how to describe this in terms of decomposition. Define

$$M_a = X_a (\det X)^m, \tag{4.55}$$

then the Coulomb branch equation (4.53) can be written

$$(-1)^{k-1} q \prod_{b=1}^k \left(\frac{M_a (1 - M_a)^{nm}}{M_b (1 - M_b)^{nm}} \right) = (1 - M_a)^{n(mk+1)}. \tag{4.56}$$

Taking a product over values of a , this implies

$$q^k = \prod_{a=1}^k (1 - M_a)^{n(mk+1)}, \tag{4.57}$$

hence,

$$\prod_{a=1}^k (1 - M_a)^{nm} = q^{mk/(mk+1)} \zeta^m, \tag{4.58}$$

where ζ is a $(mk + 1)$ th root of unity. Then we can write (4.53) as

$$(-1)^{k-1} q^{1/(mk+1)} M_a^k = (1 - M_a)^n \zeta^m \det M, \tag{4.59}$$

using the fact that, for example, $\det M = (\det X)^{1+km}$.

Comparing to the quantum K theory ring relations (4.22) for the ordinary Grassmannian $G(k, n)$, we see that, as expected, the relation (4.59) describes $km + 1$ copies of the quantum K-theory ring relation of $G(k, n)$, indexed by the value of ζ , in terms of the M_a , and as shifting the X_a by $(km + 1)$ th roots of unity preserves the M_a , we see another $(km + 1)$ -fold ambiguity, for altogether a decomposition into $(km + 1)^2$ universes.

Finally, as a consistency check, let us take the $R \rightarrow 0$ limit and compare to quantum cohomology. We start from the Coulomb branch equations, repeated here

$$(-1)^{k-1} q_{3d} X_a^k = (1 - X_a (\det X)^m)^n \prod_{b=1}^k [X_b (1 - X_b (\det X)^m)^{nm}]. \tag{4.60}$$

To get quantum cohomology, we take a small R limit. Expanding, we have

$$X_a = \exp(-2\pi R \sigma_a) = 1 - 2\pi R \sigma_a + \dots, \tag{4.61}$$

$$\det X = \exp\left(-2\pi R \sum_a \sigma_a\right) = 1 - 2\pi R \sum_a \sigma_a + \dots, \tag{4.62}$$

$$q_{3d} = (2\pi R)^{n(mk+1)} q_{2d}. \tag{4.63}$$

Plugging these into the Coulomb branch equations and sending $R \rightarrow 0$, we obtain equation (4.40), as expected.

4.4 More general weighted Grassmannians

Next, for completeness, we consider a more general analogue of weighted projective spaces for Grassmannians, and their quantum K theory. Physically, these are described by a $U(k)$ gauge theory with n chiral superfields, where the i th is in the $U(k)$ representation with highest weight

$$[m_i + 1, m_i, \dots, m_i]. \tag{4.64}$$

In special cases, this will be a gerbe.

The twisted one-loop effective superpotential (for the three-dimensional theory) is given by

$$W = \frac{k}{2} \sum_a (\ln X_a)^2 - \frac{1}{2} \left(\sum_a \ln X_a \right)^2 + \ln(-1)^{k-1} q_{3d} \sum_a \ln X_a + \sum_i \sum_a \text{Li}_2(X_a (\det X)^{m_i}). \tag{4.65}$$

The Coulomb branch equation is given by

$$(-1)^{k-1} q_{3d} X_a^k = (\det X) \prod_i \left[(1 - X_a (\det X)^{m_i}) \prod_b (1 - X_b (\det X)^{m_i})^{m_i} \right]. \tag{4.66}$$

Then taking the 2d limit, we obtain

$$(-1)^{k-1} q_{2d} = \prod_i \left[(\sigma_a + m_i \sum_c \sigma_c) \prod_b (\sigma_b + m_i \sum_c \sigma_c)^{m_i} \right], \quad (4.67)$$

where

$$q_{3d} = (2\pi R)^{n+k(m_1+\dots+m_n)} q_{2d}. \quad (4.68)$$

Now, let S denote a vector bundle associated to the ordinary fundamental of $U(k)$, of highest weight $[1, 0, \dots, 0]$, then the σ_a couple to S , and $\sigma_a + m_i \sum_b \sigma_b$ couples to $S \otimes (\det S)^{m_i}$.

For the ordinary case, we symmetrized so that the expression is symmetric in σ_a 's. For the gerby case where all the m_i 's are equal to m , we defined $\tau_a = \sigma_a + m \sum_b \sigma_b$ and made symmetrization in terms of τ_a 's. Here, we define

$$\tau_a^i = \sigma_a + m_i \sum_b \sigma_b, \quad (4.69)$$

and we symmetrize over τ_a^i for each fixed $i = 1, \dots, n$. The Coulomb branch equations become

$$(-1)^{k-1} q_{2d} = \tau_a^1 \tau_a^2 \dots \tau_a^n \prod_b (\tau_b^1)^{m_1} (\tau_b^2)^{m_2} \dots (\tau_b^n)^{m_n}. \quad (4.70)$$

4.5 Gerbes on flag manifolds

In this section, we will outline

$$\prod_{i=1}^s \mathbb{Z}_{m_i k_i + 1} \quad (4.71)$$

gerbes on a flag manifold $Fl(k_1, \dots, k_s, n)$.

The GLSM for an ordinary flag manifold $Fl(k_1, k_2, \dots, k_s, n)$ is a $U(k_1) \times U(k_2) \times \dots \times U(k_s)$ gauge theory with matter fields which are bifundamentals in the $(\mathbf{k}_i, \overline{\mathbf{k}}_{i+1})$ representation of $U(k_i) \times U(k_{i+1})$ for $i = 1, 2, \dots, s-1$, and n fundamentals of $U(k_s)$ [84].

The GLSM for the desired gerbe on the flag manifold $Fl(k_1, k_2, \dots, k_s, n)$ is constructed as a generalization of the weighted Grassmannian. Specifically, it is a $U(k_1) \times U(k_2) \times \dots \times U(k_s)$ gauge theory, with chiral fields $\Phi^{(i)}$ transforming in the $U(k_i)$ representation with highest weight $[m_i + 1, m_i, \dots, m_i]$ and in the $U(k_{i+1})$ representation with highest weight $[-m_{i+1} - 1, -m_{i+1}, \dots, -m_{i+1}]$, for $i = 1, 2, \dots, s-1$. There are also n chiral fields $\Phi^{(s)}$ transforming in the $U(k_s)$ representation with highest weight $[m_s + 1, m_s, \dots, m_s]$.

Then the pertinent twisted superpotential for the i th ($i = 1, 2, 3, \dots, s$) step is

$$\begin{aligned} W_i = & \frac{k_i}{2} \sum_{a=1}^{k_i} (\ln X_a^{(i)})^2 - \frac{1}{2} \left(\sum_{a=1}^{k_i} \ln X_a^{(i)} \right)^2 + (\ln(-1)^{k_i-1} q_i) \sum_{a=1}^{k_i} \ln X_a^{(i)} \\ & + \sum_{a=1}^{k_i} \sum_{b=1}^{k_{i-1}} \text{Li}_2 \left(\frac{X_b^{(i-1)} (\det X^{(i-1)})^{m_{i-1}}}{X_a^{(i)} (\det X^{(i)})^{m_i}} \right) + \sum_{a=1}^{k_i} \sum_{b=1}^{k_{i+1}} \text{Li}_2 \left(\frac{X_a^{(i)} (\det X^{(i)})^{m_i}}{X_b^{(i+1)} (\det X^{(i+1)})^{m_{i+1}}} \right), \end{aligned}$$

with $k_0 = 0$ understood and $X_a^{(s+1)}$ being the equivariant parameters.

Let $Y_a^{(i)} = X_a^{(i)} (\det X^{(i)})^{m_i}$, the Coulomb branch equation is

$$\begin{aligned} & (-1)^{k_i-1} q_i \left(Y_a^{(i)} \right)^{k_i} \prod_{b=1}^{k_i-1} \left[\left(1 - \frac{Y_b^{(i-1)}}{Y_a^{(i)}} \right) \prod_{c=1}^{k_i} \left(1 - \frac{Y_b^{(i-1)}}{Y_c^{(i)}} \right)^{m_i} \right] \\ &= \left(\det Y^{(i)} \right) \prod_{b=1}^{k_i+1} \left[\left(1 - \frac{Y_a^{(i)}}{Y_b^{(i+1)}} \right) \prod_{c=1}^{k_i} \left(1 - \frac{Y_c^{(i)}}{Y_b^{(i+1)}} \right)^{m_i} \right]. \end{aligned} \quad (4.72)$$

For this, we obtain

$$q_i^{k_i} \prod_{a=1}^{k_i} \prod_{b=1}^{k_i-1} \left(1 - \frac{Y_b^{(i-1)}}{Y_a^{(i)}} \right)^{m_i k_i + 1} = \prod_{a=1}^{k_i} \prod_{b=1}^{k_i+1} \left(1 - \frac{Y_a^{(i)}}{Y_b^{(i+1)}} \right)^{m_i k_i + 1}. \quad (4.73)$$

Therefore, we obtain

$$q_i^{\frac{k_i}{m_i k_i + 1}} \zeta_i \prod_{a,b} \left(1 - \frac{Y_b^{(i-1)}}{Y_a^{(i)}} \right) = \prod_{a,b} \left(1 - \frac{Y_a^{(i)}}{Y_b^{(i+1)}} \right), \quad (4.74)$$

where ζ_i is $(m_i k_i + 1)$ th root of unity. Then, we can rewrite the Coulomb branch equation as

$$(-1)^{k_i-1} q_i^{\frac{1}{m_i k_i + 1}} \left(Y_a^{(i)} \right)^{k_i} \prod_{b=1}^{k_i-1} \left(1 - \frac{Y_b^{(i-1)}}{Y_a^{(i)}} \right) = \left(\det Y^{(i)} \right) \prod_{b=1}^{k_i+1} \left(1 - \frac{Y_a^{(i)}}{Y_b^{(i+1)}} \right) \zeta_i^{m_i}. \quad (4.75)$$

We see that, these relations describe $\prod_i (m_i k_i + 1)$ copies of the quantum K-theory ring relations of $Fl(k_1, k_2, \dots, k_s, n)$, indexed by $(\zeta_1, \zeta_2, \dots, \zeta_s)$, as generated by the $Y^{(i)}$'s, which are invariant under multiplication of the $X^{(i)}$ by $(k_i m_i + 1)$ th roots of unity, for altogether a decomposition into $\prod_i (m_i k_i + 1)^2$ universes.

We can take the $R \rightarrow 0$ limit, and obtain the quantum cohomology ring relations. We have

$$X_a^{(i)} = \exp(-2\pi R \sigma_a^{(i)}) = 1 - 2\pi R \sigma_a^{(i)} + \dots \quad (4.76)$$

$$\det X^{(i)} = \exp \left(-2\pi R \sum_a \sigma_a^{(i)} \right) = 1 - 2\pi R \sum_a \sigma_a^{(i)} + \dots \quad (4.77)$$

$$q_i^{3d} = (2\pi R)^{(k_{i+1} - k_{i-1})(m_i k_i + 1)} q_i^{2d}. \quad (4.78)$$

Let

$$\tau_a^{(i)} = \sigma_a^{(i)} + m_i \sum_b \sigma_b^{(i)}, \quad (4.79)$$

the quantum cohomology ring relations can be written as

$$\begin{aligned} & (-1)^{k_i-1} q_i^{2d} \prod_{b=1}^{k_i-1} \left[\left(\tau_b^{(i-1)} - \tau_a^{(i)} \right) \prod_{c=1}^{k_i} \left(\tau_b^{(i-1)} - \tau_c^{(i)} \right)^{m_i} \right] \\ &= \prod_{b=1}^{k_i+1} \left[\left(\tau_a^{(i)} - \tau_b^{(i+1)} \right) \prod_{c=1}^{k_i} \left(\tau_c^{(i)} - \tau_b^{(i+1)} \right)^{m_i} \right]. \end{aligned} \quad (4.80)$$

Finally, we have

$$(-1)^{k_i-1} q_i^{1/(m_i k_i + 1)} \prod_{b=1}^{k_i-1} \left(\sigma_b^{(i-1)} - \sigma_a^{(i)} \right) = \prod_{b=1}^{k_i+1} \left(\tau_a^{(i)} - \tau_b^{(i+1)} \right) \zeta_i^{m_i}, \quad (4.81)$$

for ζ_i a $(m_i k_i + 1)$ th root of unity. Again, these describe $\prod_i (m_i k_i + 1)$ copies of the quantum cohomology ring relation of $Fl(k_1, k_2, \dots, k_s, n)$, indexed by the value of ζ_i 's, as expected from decomposition for two-dimensional theories.

4.6 More general levels: projective spaces

So far, we have discussed three-dimensional GLSMs for gerbes with Chern-Simons terms chosen so as to get OPE rings matching quantum K theory in mathematics. In this section, we will briefly outline projective spaces with more general levels, to outline some of the complications that can ensue.

Consider a GLSM for a gerby projective space \mathbb{P}^n , meaning a $U(1)$ gauge theory with $n + 1$ chiral superfields of charge ℓ , and with Chern-Simons terms at level k . Following [55, equ'n (2.1)], the superpotential describing this theory is

$$W = \frac{1}{2} \left(k + \ell^2 \frac{n+1}{2} \right) (\ln X)^2 + (\ln q) (\ln X) + \sum_{i=1}^{n+1} \text{Li}_2 \left(X^\ell \right). \tag{4.82}$$

The equations of motion are

$$\left(1 - X^\ell \right)^{\ell(n+1)} = q X^{k + \ell^2(n+1)/2}. \tag{4.83}$$

If we wanted to recover quantum K theory specifically, we would determine the Chern-Simons level from $U(1)_{-1/2}$ quantization, which would stipulate

$$k = -\frac{1}{2} \sum_i (Q_i)^2 = -\ell^2 \frac{n+1}{2}, \tag{4.84}$$

where the Q_i 's are the gauge charges of the chiral superfields. It is easy to see that for this level, the equations of motion reduce to

$$\left(1 - X^\ell \right)^{\ell(n+1)} = q, \tag{4.85}$$

which we have discussed previously.

Now, suppose k is more general. Let us consider some cases.

- First, suppose that k is divisible by ℓ : $k = p\ell$ for some integer p . In this case, from the general discussion of section 2, there should be a $B\mathbb{Z}_\ell$ symmetry and a decomposition in the two-dimensional theory, and corresponding to that, we can take an ℓ th root of the equations of motion (4.83) to get

$$\left(1 - X^\ell \right)^{n+1} = \xi q^{1/\ell} X^{p + \ell(n+1)/2}, \tag{4.86}$$

where ξ is an ℓ th root of unity. For each choice of ξ , we get a different theory, and the equation above describes the classical solutions of that theory.

- If in addition, k is divisible by ℓ^2 , then the equations of motion above are a polynomial in X^ℓ . If we write $p = \ell r$, and define $Y = X^\ell$, then the equations of motion become

$$\left(1 - Y \right)^{n+1} = \xi q^{1/\ell} Y^{r + (n+1)/2}, \tag{4.87}$$

which are the vacua corresponding to the GLSM for an ordinary projective space \mathbb{P}^n with level r . Taking roots of $Y = X^\ell$ results in ℓ copies. In other words, if $k = \ell^2 r$, then the equations of motion are the same as ℓ^2 copies of those for the GLSM for \mathbb{P}^n with Chern-Simons term at level r . In short, a decomposition squared, as expected.

However, it is essential for this last point that k be divisible by ℓ^2 . If k is only divisible by ℓ , not ℓ^2 , then we do not get two orders of decomposition.

- For completeness, if k is not divisible by ℓ , but the $\gcd(k, \ell) > 1$, then we can repeat a similar argument, in which we get at least an order $\gcd(k, \ell)$ decomposition, and potentially more if the Chern-Simons level has further divisibility properties.

We leave a thorough classification of all possibilities for future work.

5 Conclusions

In this paper we have discussed how decomposition [1] plays a role in three-dimensional gauge theories with one-form symmetries. Although the three-dimensional theory itself does not decompose, effective two-dimensional theories of parallel one-dimensional objects, or for that matter dimensional reductions, do decompose, in two separate ways. As a result, if one starts with a theory with a $B\mathbb{Z}_k$ one-form symmetry, the effective two-dimensional theory will decompose into, locally, k^2 universes. This was initially proposed in [16, 17], and we have extended their analysis to more general cases (resulting in more complex decompositions). This structure also immediately makes a prediction for quantum K theory rings, which are realized as OPE rings of parallel Wilson lines in three-dimensional theories.

In principle, the same ideas should apply in higher dimensions. For example, parallel surfaces in four-dimensional gauge theories with one-form symmetries should also exhibit decomposition in their OPEs, in multiple ways, even though the theory as a whole does not decompose, as outlined in the introduction. We leave this for future work. Similar ideas should also apply in theories with gauged trivially-acting noninvertible symmetries, as discussed in e.g. [30].

Acknowledgments

We would like to thank W. Gu, A. Perez-Lona, I. Melnikov, L. Mihalcea, T. Pantev, H.-H. Tseng, W. Xu, and X. Yu for useful comments. We would also like to thank the referee for suggesting a generalization of the analysis in the original draft. E.S. was partially supported by NSF grant PHY-2310588.

A Bundles on stacks and gerbes

Deligne-Mumford stacks can typically be presented as $[X/G]$, where G is any group (not necessarily finite), with any action on X (not necessarily effective). The stack $[X/G]$ is said to be a K -gerbe if a subgroup $K \subset G$ acts trivially on X .

The cohomology of the stack $\mathfrak{X} = [X/G]$ is most naturally defined on the inertia stack $I_{\mathfrak{X}}$. Intuitively, the inertia stack is the zero-momentum part of the loop space of \mathfrak{X} , and as

such, has one component which is a copy of \mathfrak{X} , plus other components (due to the existence of automorphisms encoded in \mathfrak{X}). Each component is a copy of a substack of \mathfrak{X} . Those components are labelled by automorphisms α . The group generated by any automorphism α , call it $\langle \alpha \rangle$, is cyclic.

For one example, suppose $\mathfrak{X} = [\mathbb{C}^2/\mathbb{Z}_2]$, with the \mathbb{Z}_2 acting by sign flips. This has one fixed point, at the origin of the plane \mathbb{C}^2 . In this case,

$$I_{\mathfrak{X}} = [\mathbb{C}^2/\mathbb{Z}_2] \coprod [\text{point}/\mathbb{Z}_2]. \tag{A.1}$$

The second component is associated with an order-two automorphism.

For another example, suppose $\mathfrak{X} = [X/\mathbb{Z}_k]$ where all of \mathbb{Z}_k acts trivially. In that case,

$$I_{\mathfrak{X}} = \coprod_{m=0}^{k-1} [X/\mathbb{Z}_k]. \tag{A.2}$$

Let $\pi : I_{\mathfrak{X}} \rightarrow \mathfrak{X}$ denote the projector whose restriction to any component is the projection onto that component. We denote the restriction of π to the component λ by π_{λ} .

Let $E \rightarrow \mathfrak{X}$ be a vector bundle. A sheaf or bundle on the stack $\mathfrak{X} = [X/G]$, is precisely the same as a G -equivariant sheaf or bundle on X , the covering space, so E is the same as a G -equivariant bundle on X .

On each component of $I_{\mathfrak{X}}$, $\pi_{\lambda}^* E$ will decompose into eigenbundles of the action of the stabilizer $\alpha(\lambda)$:

$$\pi_{\lambda}^* E|_{\lambda} = \bigoplus_{\chi} E_{\lambda, \chi}, \tag{A.3}$$

where χ is a character of the stabilizer $\alpha(\lambda)$. One defines $\text{ch}^{\text{rep}}(E)$ over a component of $I_{\mathfrak{X}}$ to be

$$\text{ch}^{\text{rep}}(E)|_{\lambda} = \bigoplus_{\chi} \text{ch}(E_{\lambda, \chi}) \otimes \chi, \tag{A.4}$$

where ch denotes the ordinary Chern character in equivariant cohomology. The reader should note that, curiously, ch^{rep} is a complex-valued cohomology class.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] S. Hellerman et al., *Cluster decomposition, T-duality, and gerby CFT's*, *Adv. Theor. Math. Phys.* **11** (2007) 751 [[hep-th/0606034](#)] [[INSPIRE](#)].
- [2] T. Pantev and E. Sharpe, *Notes on gauging noneffective group actions*, [hep-th/0502027](#) [[INSPIRE](#)].
- [3] T. Pantev and E. Sharpe, *String compactifications on Calabi-Yau stacks*, *Nucl. Phys. B* **733** (2006) 233 [[hep-th/0502044](#)] [[INSPIRE](#)].
- [4] T. Pantev and E. Sharpe, *GLSM's for Gerbes (and other toric stacks)*, *Adv. Theor. Math. Phys.* **10** (2006) 77 [[hep-th/0502053](#)] [[INSPIRE](#)].

- [5] A. Caldararu et al., *Non-birational twisted derived equivalences in abelian GLSMs*, *Commun. Math. Phys.* **294** (2010) 605 [[arXiv:0709.3855](#)] [[INSPIRE](#)].
- [6] L.B. Anderson et al., *General aspects of heterotic string compactifications on stacks and gerbes*, *Adv. Theor. Math. Phys.* **19** (2015) 531 [[arXiv:1307.2269](#)] [[INSPIRE](#)].
- [7] E. Sharpe, *Decomposition in diverse dimensions*, *Phys. Rev. D* **90** (2014) 025030 [[arXiv:1404.3986](#)] [[INSPIRE](#)].
- [8] E. Sharpe, *Undoing decomposition*, *Int. J. Mod. Phys. A* **34** (2020) 1950233 [[arXiv:1911.05080](#)] [[INSPIRE](#)].
- [9] Y. Tanizaki and M. Ünsal, *Modified instanton sum in QCD and higher-groups*, *JHEP* **03** (2020) 123 [[arXiv:1912.01033](#)] [[INSPIRE](#)].
- [10] Z. Komargodski, K. Ohmori, K. Roumpedakis and S. Seifnashri, *Symmetries and strings of adjoint QCD₂*, *JHEP* **03** (2021) 103 [[arXiv:2008.07567](#)] [[INSPIRE](#)].
- [11] W. Gu, E. Sharpe and H. Zou, *Notes on two-dimensional pure supersymmetric gauge theories*, *JHEP* **04** (2021) 261 [[arXiv:2005.10845](#)] [[INSPIRE](#)].
- [12] R. Eager and E. Sharpe, *Elliptic Genera of Pure Gauge Theories in Two Dimensions with Semisimple Non-Simply-Connected Gauge Groups*, *Commun. Math. Phys.* **387** (2021) 267 [[arXiv:2009.03907](#)] [[INSPIRE](#)].
- [13] A. Cherman and T. Jacobson, *Lifetimes of near eternal false vacua*, *Phys. Rev. D* **103** (2021) 105012 [[arXiv:2012.10555](#)] [[INSPIRE](#)].
- [14] M. Nguyen, Y. Tanizaki and M. Ünsal, *Semi-Abelian gauge theories, non-invertible symmetries, and string tensions beyond N-ality*, *JHEP* **03** (2021) 238 [[arXiv:2101.02227](#)] [[INSPIRE](#)].
- [15] M. Nguyen, Y. Tanizaki and M. Ünsal, *Noninvertible 1-form symmetry and Casimir scaling in 2D Yang-Mills theory*, *Phys. Rev. D* **104** (2021) 065003 [[arXiv:2104.01824](#)] [[INSPIRE](#)].
- [16] W. Gu, D. Pei and M. Zhang, *On phases of 3d N=2 Chern-Simons-matter theories*, *Nucl. Phys. B* **973** (2021) 115604 [[arXiv:2105.02247](#)] [[INSPIRE](#)].
- [17] W. Gu, *Vacuum Structures Revisited*, [arXiv:2110.13156](#) [[INSPIRE](#)].
- [18] D.G. Robbins, E. Sharpe and T. Vandermeulen, *Anomalies, extensions, and orbifolds*, *Phys. Rev. D* **104** (2021) 085009 [[arXiv:2106.00693](#)] [[INSPIRE](#)].
- [19] D.G. Robbins, E. Sharpe and T. Vandermeulen, *Quantum symmetries in orbifolds and decomposition*, *JHEP* **02** (2022) 108 [[arXiv:2107.12386](#)] [[INSPIRE](#)].
- [20] D.G. Robbins, E. Sharpe and T. Vandermeulen, *Anomaly resolution via decomposition*, *Int. J. Mod. Phys. A* **36** (2021) 2150220 [[arXiv:2107.13552](#)] [[INSPIRE](#)].
- [21] E. Sharpe, *Topological operators, noninvertible symmetries and decomposition*, *Adv. Theor. Math. Phys.* **27** (2023) 2319 [[arXiv:2108.13423](#)] [[INSPIRE](#)].
- [22] M. Honda, E. Itou, Y. Kikuchi and Y. Tanizaki, *Negative string tension of a higher-charge Schwinger model via digital quantum simulation*, *PTEP* **2022** (2022) 033B01 [[arXiv:2110.14105](#)] [[INSPIRE](#)].
- [23] T. Pantev, D.G. Robbins, E. Sharpe and T. Vandermeulen, *Orbifolds by 2-groups and decomposition*, *JHEP* **09** (2022) 036 [[arXiv:2204.13708](#)] [[INSPIRE](#)].
- [24] T. Pantev and E. Sharpe, *Decomposition in Chern-Simons theories in three dimensions*, *Int. J. Mod. Phys. A* **37** (2022) 2250227 [[arXiv:2206.14824](#)] [[INSPIRE](#)].

- [25] L. Lin, D.G. Robbins and E. Sharpe, *Decomposition, Condensation Defects, and Fusion*, *Fortsch. Phys.* **70** (2022) 2200130 [[arXiv:2208.05982](#)] [[INSPIRE](#)].
- [26] S. Meynet and R. Moscrop, *McKay quivers and decomposition*, *Lett. Math. Phys.* **113** (2023) 63 [[arXiv:2208.07884](#)] [[INSPIRE](#)].
- [27] D.G. Robbins, E. Sharpe and T. Vandermeulen, *Decomposition, trivially-acting symmetries, and topological operators*, *Phys. Rev. D* **107** (2023) 085017 [[arXiv:2211.14332](#)] [[INSPIRE](#)].
- [28] A. Perez-Lona and E. Sharpe, *Three-dimensional orbifolds by 2-groups*, *JHEP* **08** (2023) 138 [[arXiv:2303.16220](#)] [[INSPIRE](#)].
- [29] T. Pantev and E. Sharpe, *Decomposition and the Gross–Taylor string theory*, *Int. J. Mod. Phys. A* **38** (2023) 2350156 [[arXiv:2307.08729](#)] [[INSPIRE](#)].
- [30] A. Perez-Lona et al., *Notes on gauging noninvertible symmetries. Part I. Multiplicity-free cases*, *JHEP* **02** (2024) 154 [[arXiv:2311.16230](#)] [[INSPIRE](#)].
- [31] E. Sharpe, *Dilaton shifts, probability measures, and decomposition*, [arXiv:2312.08438](#) [[DOI:10.1088/1751-8121/ad8196](#)] [[INSPIRE](#)].
- [32] L. Bhardwaj, D. Pajer, S. Schafer-Nameki and A. Warman, *Hasse Diagrams for Gapless SPT and SSB Phases with Non-Invertible Symmetries*, [arXiv:2403.00905](#) [[INSPIRE](#)].
- [33] E. Sharpe, *Landau-Ginzburg models, gerbes, and Kuznetsov’s homological projective duality*, *Proc. Symp. Pure Math.* **81** (2010) 237 [[INSPIRE](#)].
- [34] E. Sharpe, *GLSM’s, Gerbes, and Kuznetsov’s Homological Projective Duality*, *J. Phys. Conf. Ser.* **462** (2013) 012047 [[arXiv:1004.5388](#)] [[INSPIRE](#)].
- [35] E. Sharpe, *Categorical Equivalence and the Renormalization Group*, *Fortsch. Phys.* **67** (2019) 1910019 [[arXiv:1903.02880](#)] [[INSPIRE](#)].
- [36] E. Sharpe, *An introduction to decomposition*, [arXiv:2204.09117](#) [[INSPIRE](#)].
- [37] E. Andreini, Y. Jiang and H.-H. Tseng, *On Gromov-Witten theory of root gerbes*, [arXiv:0812.4477](#).
- [38] E. Andreini, Y. Jiang and H.-H. Tseng, *Gromov-Witten theory of product stacks*, [arXiv:0905.2258](#).
- [39] E. Andreini, Y. Jiang and H.-H. Tseng, *Gromov-Witten theory of root gerbes I: structure of genus 0 moduli spaces*, [arXiv:0907.2087](#).
- [40] H.-H. Tseng, *On degree zero elliptic orbifold Gromov-Witten invariants*, [arXiv:0912.3580](#).
- [41] A. Gholampour and H.-H. Tseng, *On Donaldson-Thomas invariants of threefold stacks and gerbes*, [arXiv:1001.0435](#).
- [42] X. Tang and H.-H. Tseng, *Duality theorems of etale gerbes on orbifolds*, [arXiv:1004.1376](#) [[INSPIRE](#)].
- [43] M. Bullimore, H.-C. Kim and P. Koroteev, *Defects and Quantum Seiberg-Witten Geometry*, *JHEP* **05** (2015) 095 [[arXiv:1412.6081](#)] [[INSPIRE](#)].
- [44] K. Ueda and Y. Yoshida, *3d $\mathcal{N} = 2$ Chern-Simons-matter theory, Bethe ansatz, and quantum K-theory of Grassmannians*, *JHEP* **08** (2020) 157 [[arXiv:1912.03792](#)] [[INSPIRE](#)].
- [45] N.A. Nekrasov and S.L. Shatashvili, *Supersymmetric vacua and Bethe ansatz*, *Nucl. Phys. B Proc. Suppl.* **192-193** (2009) 91 [[arXiv:0901.4744](#)] [[INSPIRE](#)].

- [46] C. Closset and H. Kim, *Comments on twisted indices in 3d supersymmetric gauge theories*, *JHEP* **08** (2016) 059 [[arXiv:1605.06531](#)] [[INSPIRE](#)].
- [47] H. Jockers and P. Mayr, *A 3d Gauge Theory/Quantum K-Theory Correspondence*, *Adv. Theor. Math. Phys.* **24** (2020) 327 [[arXiv:1808.02040](#)] [[INSPIRE](#)].
- [48] H. Jockers and P. Mayr, *Quantum K-Theory of Calabi-Yau Manifolds*, *JHEP* **11** (2019) 011 [[arXiv:1905.03548](#)] [[INSPIRE](#)].
- [49] H. Jockers, P. Mayr, U. Ninad and A. Tabler, *Wilson loop algebras and quantum K-theory for Grassmannians*, *JHEP* **10** (2020) 036 [[arXiv:1911.13286](#)] [[INSPIRE](#)].
- [50] C. Closset, H. Kim and B. Willett, *Supersymmetric partition functions and the three-dimensional A-twist*, *JHEP* **03** (2017) 074 [[arXiv:1701.03171](#)] [[INSPIRE](#)].
- [51] C. Closset, H. Kim and B. Willett, *Seifert fibering operators in 3d $\mathcal{N} = 2$ theories*, *JHEP* **11** (2018) 004 [[arXiv:1807.02328](#)] [[INSPIRE](#)].
- [52] C. Closset and H. Kim, *Three-dimensional $N=2$ supersymmetric gauge theories and partition functions on Seifert manifolds: A review*, *Int. J. Mod. Phys. A* **34** (2019) 1930011 [[arXiv:1908.08875](#)] [[INSPIRE](#)].
- [53] C. Closset and O. Khlaif, *Twisted indices, Bethe ideals and 3d $\mathcal{N} = 2$ infrared dualities*, *JHEP* **05** (2023) 148 [[arXiv:2301.10753](#)] [[INSPIRE](#)].
- [54] C. Closset and O. Khlaif, *Grothendieck lines in 3d $\mathcal{N} = 2$ SQCD and the quantum K-theory of the Grassmannian*, *JHEP* **12** (2023) 082 [[arXiv:2309.06980](#)] [[INSPIRE](#)].
- [55] W. Gu, L. Mihalcea, E. Sharpe and H. Zou, *Quantum K theory of symplectic Grassmannians*, *J. Geom. Phys.* **177** (2022) 104548 [[arXiv:2008.04909](#)] [[INSPIRE](#)].
- [56] W. Gu, L.C. Mihalcea, E. Sharpe and H. Zou, *Quantum K theory of Grassmannians, Wilson line operators, and Schur bundles*, [arXiv:2208.01091](#) [[INSPIRE](#)].
- [57] W. Gu et al., *Quantum K theory rings of partial flag manifolds*, *J. Geom. Phys.* **198** (2024) 105127 [[arXiv:2306.11094](#)] [[INSPIRE](#)].
- [58] W. Gu et al., *Quantum K Whitney relations for partial flag varieties*, [arXiv:2310.03826](#) [[INSPIRE](#)].
- [59] S. Chun, S. Gukov, S. Park and N. Sopenko, *3d-3d correspondence for mapping tori*, *JHEP* **09** (2020) 152 [[arXiv:1911.08456](#)] [[INSPIRE](#)].
- [60] O. Aharony, S.S. Razamat and B. Willett, *From 3d duality to 2d duality*, *JHEP* **11** (2017) 090 [[arXiv:1710.00926](#)] [[INSPIRE](#)].
- [61] N. Seiberg and E. Witten, *Gapped Boundary Phases of Topological Insulators via Weak Coupling*, *PTEP* **2016** (2016) 12C101 [[arXiv:1602.04251](#)] [[INSPIRE](#)].
- [62] N. Seiberg, T. Senthil, C. Wang and E. Witten, *A Duality Web in 2+1 Dimensions and Condensed Matter Physics*, *Annals Phys.* **374** (2016) 395 [[arXiv:1606.01989](#)] [[INSPIRE](#)].
- [63] P.-S. Hsin, H.T. Lam and N. Seiberg, *Comments on One-Form Global Symmetries and Their Gauging in 3d and 4d*, *SciPost Phys.* **6** (2019) 039 [[arXiv:1812.04716](#)] [[INSPIRE](#)].
- [64] F. Benini, C. Copetti and L. Di Pietro, *Factorization and global symmetries in holography*, *SciPost Phys.* **14** (2023) 019 [[arXiv:2203.09537](#)] [[INSPIRE](#)].
- [65] D. Belov and G.W. Moore, *Classification of Abelian spin Chern-Simons theories*, [hep-th/0505235](#) [[INSPIRE](#)].

- [66] D.S. Freed, *Classical Chern-Simons theory. Part 1*, *Adv. Math.* **113** (1995) 237 [[hep-th/9206021](#)] [[INSPIRE](#)].
- [67] D.S. Freed, *Classical Chern-Simons theory, part 2*, *Houston J. Math.* **28** (2002) 293, [https://www.math.uh.edu/~hjm/pdf28\(2\)/03freed.pdf](https://www.math.uh.edu/~hjm/pdf28(2)/03freed.pdf).
- [68] *Kinds of Wilson Loops in a $U(1)$ Chern-Simons Theory*, <https://physics.stackexchange.com/questions/521200/kinds-of-wilson-loops-in-a-u1-chern-simons-theory>.
- [69] C. Closset, E. Furrer and O. Khlaif, *One-form symmetries and the 3d $\mathcal{N} = 2A$ -model: Topologically twisted indices for any G* , [arXiv:2405.18141](#) [[INSPIRE](#)].
- [70] Y. Qiu and Z. Wang, *Representations of Motion Groups of Links via Dimension Reduction of TQFTs*, *Commun. Math. Phys.* **382** (2021) 2071 [[arXiv:2002.07642](#)] [[INSPIRE](#)].
- [71] L. Müller and L. Woike, *Dimensional reduction, extended topological field theories and orbifoldization*, *Bull. London Math. Soc.* **53** (2023) 392 [[arXiv:2004.04689](#)] [[INSPIRE](#)].
- [72] T. Lan, L. Kong and X.-G. Wen, *Classification of $(3+1)D$ Bosonic Topological Orders: The Case When Pointlike Excitations Are All Bosons*, *Phys. Rev. X* **8** (2018) 021074 [[INSPIRE](#)].
- [73] D.R. Morrison and M.R. Plesser, *Summing the instantons: Quantum cohomology and mirror symmetry in toric varieties*, *Nucl. Phys. B* **440** (1995) 279 [[hep-th/9412236](#)] [[INSPIRE](#)].
- [74] M. Zhang, *Quantum K-theory of toric stacks*, <https://mathweb.ucsd.edu/~miz017/quantumk.pdf>.
- [75] E. González and C. Woodward, *Quantum Kirwan for quantum K-theory*, [arXiv:1911.03520](#).
- [76] T.J. Jarvis, R. Kaufmann and T. Kimura, *Stringy K-theory and the Chern character*, [math/0502280](#) [[DOI:10.1007/s00222-006-0026-x](#)].
- [77] A. Adem, Y. Ruan and B. Zhang, *A Stringy Product on Twisted Orbifold K-theory*, [math/0605534](#).
- [78] E. Witten, *The Verlinde algebra and the cohomology of the Grassmannian*, [hep-th/9312104](#) [[INSPIRE](#)].
- [79] A. Givental and Y.-P. Lee, *Quantum K-theory on flag manifolds, finite-difference Toda lattices and quantum groups*, *Invent. Math.* **151** (2003) 193 [[math/0108105](#)] [[INSPIRE](#)].
- [80] P. Koroteev, P.P. Pushkar, A.V. Smirnov and A.M. Zeitlin, *Quantum K-theory of quiver varieties and many-body systems*, *Selecta Math.* **27** (2021) 87 [[arXiv:1705.10419](#)] [[INSPIRE](#)].
- [81] A. Givental and B.-S. Kim, *Quantum cohomology of flag manifolds and Toda lattices*, *Commun. Math. Phys.* **168** (1995) 609 [[hep-th/9312096](#)] [[INSPIRE](#)].
- [82] B. Kim, *Quantum cohomology of flag manifolds G/B and quantum Toda lattices*, [alg-geom/9607001](#).
- [83] W. Gu and E. Sharpe, *A proposal for nonabelian mirrors*, [arXiv:1806.04678](#) [[INSPIRE](#)].
- [84] R. Donagi and E. Sharpe, *GLSM's for partial flag manifolds*, *J. Geom. Phys.* **58** (2008) 1662 [[arXiv:0704.1761](#)] [[INSPIRE](#)].