

Table 2. Incidence of *Cylindrocladium parasiticum* in normal and speckled peanut seed from 19 commercial seed lots before and after storage at 15°C.

Cultivar	Seed lot	Date of initial assay	Seed assay (% positive) ^a		
			Normal	Speckled	
			Initial	Initial	Jun 00
VA 98R	314	23 Nov 99	2	90	32
	315	23 Nov 99	0	88	8
	316	23 Nov 99	0	94	18
	318	11 Dec 99	0	88	4
	319	11 Dec 99	0	66	12
	320	11 Dec 99	0	64	6
	321	12 Dec 99	0	44	6
NC-V 11	322	30 Dec 99	0	90	78
	323	30 Dec 99	0	88	72
	324	13 Jan 00	0	86	74
	325	19 Jan 00	0	96	68
VA-C 92R	330	22 Dec 99	0	82	64
	331	23 Dec 99	0	86	48
	337	13 Jan 00	0	80	6
NC 7	338	19 Jan 00	0	80	22
	339	21 Jan 00	0	76	20
	340	22 Jan 00	0	74	10
	341	12 Mar 00	0	40	18
	342	12 Mar 00	0	40	14

^a Data are recovery percentages of *C. parasiticum* from 50 seed. Seeds were assayed on a selective medium.