

## List of Figures

Figure		page #
6.B.1	True underlying mean function from Equation 6.B.1 for various values of $\gamma$ .	53
6.B.2	True underlying variance function from Equation 6.B.2.	53
6.B.3	Scatter plot of a random data set generated from the dual model given in (6.B.1) and (6.B.2) where $\gamma = 5.0$ .	54
6.B.4	Plot of true means function in Example 1 along with the raw data and EWLS means fit.	54
6.B.5	Plot of parametric, GLM variance estimate using squared EWLS from the means fit in Figure 6.B.4.	55
6.B.6	Plot of true means function in Example 1 along with the raw data and LLR means fit.	56
6.B.7	Plot of non-parametric, residual-based variance estimate using the squared LLR residuals from the means fit in Figure 6.B.6	56
6.B.8	Plot of non-parametric, difference-based variance estimate using the squared pseudo-residuals formed from the raw data.	57
6.B.9	Plot of true means function in Example 1 along with the raw data and MMRR means fit.	58
6.B.10	Plot of the VMRR variance estimate using the squared MMRR residuals from the fit shown in Figure 6.B.9.	59
6.B.11	EWLS fit to the raw data from Example 2 where the weights used in EWLS are the estimated variances from VMRR.	59
6.B.12	LLR fit to the residuals from the EWLS fit in Figure 6.B.11	60
6.B.13	Plot of the true mean function in Example 1 along with the EWLS, LLR, and MMRR estimates of the mean	60
6.B.14	Plot of the true variance function in Example 1 along with the GLM, Difference-Based, Residual-Based, and VMRR variance estimates.	61
6.B.15	Plot of $E(e_i^{2(\text{ewls})})$ and $E(e_i^{2(\text{mmrr})})$ for Example 1 when $\gamma = 5.0$ and $n = 41$ .	66
6.C.1	Scatter plot of a random data set generated from the dual model given in equations (6.C.1) and (6.C.2) where $\gamma = 0.0$ .	69
6.C.2	Plot showing generated data for Example 2 along with EWLS fit and the true means function.	69
6.C.3	Plot showing the EWLS squared residuals, the GLM variance estimate and the true underlying variance function for Example 2.	70

6.C.4	Plot showing generated data for Example 2 along with LLR fit and the true means function.	70
6.C.5	Plot showing the squared pseudo-residuals, the difference-based estimate and true underlying variance function for Example 2.	71
6.C.6	Plot showing the squared local linear residuals, the residual-based estimate and true underlying variance function for Example 2.	71
6.C.7	Plot showing generated data for Example 2 along with MMRR fit and the true means function.	72
6.C.8	Plot of the true means function in Example 2 along with the EWLS, LLR, and MMRR estimates of the mean.	72
6.C.9	Plot showing the MMRR squared residuals along with the VMRR estimate and true variance function for Example 2.	74
6.C.10	Parametric GLM fit to the squared MMRR residuals from Example 2.	74
6.C.11	LLR fit to residuals from parametric GLM estimate which was displayed in Figure 6.C.10.	75
6.C.12	Plot of the true variance function in Example 1 along with the GLM, Difference-Based, Residual-Based, and VMRR variance estimates ( $\gamma = 0.0$ ).	75
6.C.13	Scatter plot of random data set generated from the dual model given in equations (6.C.1) and (6.C.2) where $\gamma = 2.5$ .	78
6.C.14	Plot showing generated data for Example 2 ( $\gamma = 2.5$ ) along with the EWLS fit and true underlying means function.	78
6.C.15	Plot showing the EWLS squared residuals, the parametric variance estimate and the true underlying variance function for Example 2 ( $\gamma = 2.5$ ).	79
6.C.16	Plot showing generated data for Example 2 ( $\gamma = 2.5$ ) along with the LLR fit and true underlying means function.	79
6.C.17	Plot showing the squared pseudo-residuals, difference-based variance estimate, and true underlying variance function for Example 2 ( $\gamma = 2.5$ ).	80
6.C.18	Plot showing the squared LLR residuals, residual-based variance estimate, and true underlying variance function for Example 2 ( $\gamma = 2.5$ ).	80
6.C.19	Plot showing the generated data for data for Example 2 ( $\gamma = 2.5$ ) along with the MMRR fit and true underlying means function.	81
6.C.20	Plot showing the MMRR squared residuals along with the VMRR estimate and true variance function for Example 2 ( $\gamma = 2.5$ ).	81

6.C.21	Plot of the true means function in Example 2 ( $\gamma = 2.5$ ) along with the EWLS, LLR, and MMRR estimates of the mean.	82
6.C.22	Plot of the true variance function in Example 2 ( $\gamma = 2.5$ ) along with the GLM, Difference-Based, Residual-Based, and VMRR variance estimates.	82
7.C.1	Scatter plot of a random data set generated from the model in equation 7.C.1.	89
7.C.2	Plot of true underlying function and generated data set along with the EWLS fit.	89
7.C.3	Plot of true underlying function and generated data set along with the LLR fit.	90
7.C.4	Plot of true underlying function and generated data set along with the MMRR fit.	90
7.C.5	Plot of true underlying function and generated data set along with the EWLS, LLR, and MMRR fits.	91
7.C.6	Plot of $\ln(e_i^{2(\text{ewls})})$ vs. $z_i$ .	92
7.C.7	Plot of $\ln(e_i^{2(\text{mmrr})})$ vs. $z_i$ .	93
7.C.8	Plot of the parametric variance estimate using the squared EWLS from the means fit shown in Figure 7.C.2.	94
7.C.9	Plot of the nonparametric residual-based variance estimate using the squared LLR residuals from the fit shown in Figure 7.C.3.	95
7.C.10	Plot of the nonparametric difference-based variance estimate using squared pseudo-residuals formed from the raw data.	95
7.C.11	Plot of the VMRR variance estimate using the squared MMRR residuals from the fit shown in Figure 7.C.4.	96
7.C.12	Plot of the parametric, nonparametric residual-based, nonparametric difference-based, and VMRR estimates of variance.	96