Acknowledgements

First of all, I would like to thank my advisor, Dr. Madeline Schreiber, for all her support throughout my work at Virginia Tech. I would also like to thank Dr. Maurice Valett and Dr. J. Donald Rimstidt for valuable guidance. I would like to thank Athena Tilley for running the ICP-AES samples and Dr. Chang for running microwave digestion of soil samples. I appreciate Mary Harvey for helping me with various things in lab. I am also grateful to Brendan Brown, for helping me collecting soil samples, and Clayton Loehn for running microprobe analysis. I am also very grateful to every hydro group members. I would also like to thank my wife, Shrabanti Basu, and my family for their support. Without them, I would not be able to reach my goals in life.

Table of Contents

| Abstractii | |
|--|--|
| Acknowledgementsiii | |
| Table of Contentsiv | |
| List of Figuresvi | |
| List of Tablesvii | |
| 1.Introduction1 | |
| 2. Study site | |
| 3. Methods | |
| 3.1 Reagents and standards | |
| 3.2 Soil and rock sample collection and preparation | |
| 3.3 Sequential extraction method | |
| 3.4 Modifications of the Turpeinen method | |
| 3.4.1 Desorption experiment of As using phosphate solution | |
| 3.4.2 Extraction of arsenopyrite and scorodite | |
| 3.4.3 Quality control | |
| 3.5 Arsenic analysis | |
| 3.6 Imaging and elemental mapping: | |
| 4 Results | |
| 4.1 Results from sequential extraction | |
| 4.1.1 Total digestion results | |
| 4.1.2 Distribution of arsenic in different phases | |
| 4.1.3 Effect of grain size | |
| 4.1.4 Desorption of As by phosphate | |
| 4.1.5 Sequential extraction of arsenopyrite and scorodite on sequential extraction12 | |
| 4.2 Results from imaging, elemental mapping and electron microprobe analysis13 | |
| 4.2.1 SEM/ EMPA of sediments | |
| 4.3 SEM/EMPA of host rocks | |
| 4.3.1 SEM images | |
| 4.3.2 Elemental mapping of host rock minerals | |

| 4.3.3 Elemental distribution of arsenic in host rock | .22 |
|---|-----|
| 5. Discussion | 23 |
| 5.1 Arsenic partitioning in BAM sediments | 23 |
| 5.2 arsenic partitioning in host rock | 25 |
| 5.3 Reservoirs and Possible Release Mechanism for As at BAM | 28 |
| 6. Conclusions | 29 |
| References | 30 |
| Appendix | 32 |

List of Figures

| Figure 1: Topographic site map of Brinton Arsenic Mine | 3 |
|---|----|
| Figure 2: Average distribution of % of arsenic in different phases for mine tailing | |
| sediments and stream channel sediments | 10 |
| Figure 3: Weight % of As associated with oxides (Step 2+4) in mine tailing and stre | am |
| sediment samples | 11 |
| Figure 4: Release of As in sequential extraction from scorodite and arsenopyrite | 13 |
| Figure 5: SEM images of grains from sediment samples | 15 |
| Figure 6: % Fe in iron oxides determined by EMPA | 16 |
| Figure 7: Weight % As associated with iron oxides determined by EMPA | 17 |
| Figure 8: Weight % As associated with scorodite determined by EMPA | 18 |
| Fig 9: % As associated with arsenopyrite determined by EMPA | 19 |
| Figure 10: SEM images of scorodite/arsenopyrite | 20 |
| Figure 11: Elemental maps of arsenopyrite with scorodite and elemental S | 21 |
| Figure 12: Elemental maps of arsenopyrite grain with scorodite | 21 |
| Figure 13: Arsenopyrite grain with zones of scorodite | 22 |
| Figure 14: Arsenopyrite grain with outside intermediate zone of sulfur | 23 |
| Figure 15: Eh-pH diagram | 27 |

List of Tables:

| Table 1: The Turpeinen Extraction procedure | 5 |
|---|-----|
| Table 2: Total digestion data | 9 |
| Table 3: Effect of grain size | 12 |
| Table 4: Comparison of adsorbed As | .12 |
| Table 5: Mineral distribution in sediments | 15 |
| Table 6: % of As, Fe and S associated with Fe-oxide | 17 |
| Table 7: Weight % of Fe, As and S in scorodite | 18 |
| Table 8: Weight % of Fe, As and S in arsenopyrite | .19 |