

Near-term ecological forecasting for climate change action

Received: 23 September 2022

Accepted: 11 October 2024

Published online: 08 November 2024



Michael Dietze¹✉, Ethan P. White², Antoinette Abeyta³, Carl Boettiger⁴, Nievita Bueno Watts⁵, Cayelan C. Carey⁶, Rebecca Chaplin-Kramer^{7,8}, Ryan E. Emanuel⁹, S. K. Morgan Ernest¹⁰, Renato J. Figueiredo¹⁰, Michael D. Gerst¹¹, Leah R. Johnson¹², Melissa A. Kenney⁸, Jason S. McLachlan¹³, Ioannis Ch. Paschalidis¹⁴, Jody A. Peters¹³, Christine R. Rollinson¹⁵, Juniper Simonis¹⁶, Kira Sullivan-Wiley¹⁷, R. Quinn Thomas^{18,19}, Glenda M. Wardle¹⁹, Alyssa M. Willson¹³ & Jacob Zwart²⁰

A substantial increase in predictive capacity is needed to anticipate and mitigate the widespread change in ecosystems and their services in the face of climate and biodiversity crises. In this era of accelerating change, we cannot rely on historical patterns or focus primarily on long-term projections that extend decades into the future. In this Perspective, we discuss the potential of near-term (daily to decadal) iterative ecological forecasting to improve decision-making on actionable time frames. We summarize the current status of ecological forecasting and focus on how to scale up, build on lessons from weather forecasting, and take advantage of recent technological advances. We also highlight the need to focus on equity, workforce development, and broad cross-disciplinary and non-academic partnerships.

The dual climate and biodiversity crises¹ jeopardize our ability to manage and conserve natural resources and sustain socio-economic systems. Impacts are already being felt across all levels of society, from individuals to nations^{2–4}, and many of the world's ecosystems are at risk of collapse⁵. Indeed, when considering the most severe risks facing society over the next 10 years, the World Economic Forum ranked environmental changes as the top four most severe risks and they comprise six of the top ten⁶. In the face of accelerating change and increasingly

frequent extreme climate events, responses to these crises cannot continue to be focused primarily on projections that extend decades into the future. Similarly, historical patterns (for example, species ranges, fire/drought/flood frequency) can no longer be relied on as the primary basis for environmental decision-making⁷. Steady-state solutions do not work in a world dominated by non-equilibrium transient conditions; society is in uncharted territory. Moving forwards requires new approaches to research, management and decision-making.

¹Department of Earth & Environment, Boston University, Boston, MA, USA. ²Department of Wildlife Ecology & Conservation, University of Florida, Gainesville, FL, USA. ³Mathematics, Physical and Natural Sciences Division, University of New Mexico Gallup, Gallup, NM, USA. ⁴Department of Environmental Science, Policy and Management, University of California, Berkeley, Berkeley, CA, USA. ⁵Indian Natural Resource Science & Engineering Program, Cal Poly Humboldt, Arcata, CA, USA. ⁶Department of Biological Sciences, Virginia Tech, Blacksburg, VA, USA. ⁷Global Science, WWF, San Francisco, CA, USA. ⁸Institute on the Environment, University of Minnesota, St. Paul, MN, USA. ⁹Nicholas School of the Environment, Duke University, Durham, NC, USA. ¹⁰School of Electrical Engineering and Computer Science, Oregon State University, Corvallis, OR, USA. ¹¹Earth System Science Interdisciplinary Center, University of Maryland, College Park, MD, USA. ¹²Department of Statistics, Virginia Tech, Blacksburg, VA, USA. ¹³Department of Biological Sciences, University of Notre Dame, Notre Dame, IN, USA. ¹⁴Department of Electrical & Computer Engineering, Boston University, Boston, MA, USA. ¹⁵Center for Tree Science, The Morton Arboretum, Lisle, IL, USA. ¹⁶DAPPER Stats, Portland, OR, USA. ¹⁷The Pew Charitable Trusts, Washington DC, USA. ¹⁸Department of Forest Resources and Environmental Conservation, Virginia Tech, Blacksburg, VA, USA. ¹⁹School of Life and Environmental Sciences, University of Sydney, Sydney, New South Wales, Australia. ²⁰Water Mission Area, US Geological Survey, Madison, WI, USA. ✉e-mail: dietze@bu.edu




Fig. 1 | Near-term iterative ecological forecasting. Top: ecological forecasting provides quantitative predictions of how different management scenarios impact the environment on decision-relevant timescales. Bottom: iterative ecological forecasting involves continually updating predictions in light of new data. By establishing an iterative learning loop, ecological forecasting provides a win-win strategy: answering grand challenge questions about ecological predictability while improving environmental decisions. NEON, National Ecological Observatory Network.

Near-term iterative ecological forecasting has the potential to anticipate ecological change at the scale and speed needed by society⁸ (Fig. 1). Such forecasts provide predictions and scenario-based projections about the future state of ecosystems and their benefits to people, with fully specified uncertainties that are continually updated as new observations become available⁹. In contrast to long-term projections, near-term forecasts (on daily to decadal timescales) are more closely aligned with the timescales that are most relevant to environmental management, and thus allow society to anticipate challenges and improve decisions on actionable timeframes¹⁰. Actionable forecasts present decision-makers with a range of predictions (and in some cases projections under different decision alternatives or climate extremes) with clear statements of forecast confidence and uncertainty that can be propagated into decision analyses.

Climate change is happening now. As such, near-term iterative ecological forecasting is becoming more and more important for climate adaptation and mitigation. While many ecological forecasts have focused on projections to 2100 and beyond^{3,11–14}, the impacts of climate change on ecosystems and their services are increasingly urgent policy and management problems in the present. Recent advances in ecological forecasting are beginning to address this need. For example,

forecasts are being used in semi-arid systems to predict the impacts of real-time climate extremes on the likely success of ecological restoration efforts^{15,16} and the quantity and quality of grass that will be available for livestock to graze on^{17–19}. In marine systems, predictions about the impacts of climate variability on species migrations are fuelling real-time multispecies forecasts of fisheries bycatch risk^{20,21}. Ecological forecasts are starting to be used by freshwater managers to predict the impacts of climate change on a wide range of real-time water quality issues, including harmful algal blooms and anoxia^{22,23}. On the climate mitigation side, biogeochemical forecasts (for example, predictions of soil carbon storage and trace gas emissions that account for climate variability and alternative management scenarios) are already being used by industry as part of nature-based solutions²⁴. More broadly, the ability to forecast ecosystem responses to heatwaves, wildfires, droughts, land-use change, biological invasions and disease outbreaks will help us better understand and manage non-equilibrium conditions^{25,26} and their climate and biodiversity feedbacks.

While the development and application of near-term iterative ecological forecasts have grown rapidly in recent years, achieving the predictive capacity needed by decision-makers will require a substantial increase in the number of operationalized forecasts. Here we aim to summarize the current state of ecological forecasting research, how it is responding to societal needs (for example, climate change) and grand challenge scientific questions and where it is going, with a central focus on how to scale up ecological forecasting far beyond what is currently possible.

Global demand for ecological forecasts

While much past work on near-term ecological forecasting has focused on local- to regional-scale environmental issues, meaningfully contributing to international environmental goals and initiatives will require ecological forecasting to scale up to global issues. One place to start is forecasting the flows of mass and energy through terrestrial and marine ecosystems, as they are already present in most of the Earth system models that are part of the Coupled Model Intercomparison Project (CMIP), which in turn forms the basis for IPCC reports and projections²⁷. Next-generation Earth System models already include a wide range of ecological processes (for example, vegetation demography, microbial biogeochemistry, disturbance) and ecosystems (lakes, rivers and wetlands; urban; agricultural), as well as a more explicit representation of biodiversity^{28,29}, but these global models have primarily been applied to project long-term climate responses. So far, they have been under-used at shorter timescales where (if they are properly initialized and propagate uncertainty³⁰) there are opportunities to inform climate adaptation, mitigation, carbon monitoring and the UN Sustainable Development Goals (for example hunger, disease, water quality, sustainability, climate action and terrestrial and marine biodiversity). For example, ecological forecasts could be incorporated into the UN's Early Warnings for All climate adaptation initiative³¹, which focuses on global equity in forecasting, risk management, communication and preparedness activities from a weather and water perspective.

While long-term ecosystem forecasting is informing the UNFCCC through the IPCC process, biodiversity forecasting is just beginning to play a role in the UN Convention on Biological Diversity and the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (IPBES). For example, only one global scenario modelling effort has been integrated into IPBES so far and it was not a major focus of the assessment³². That said, the Group on Earth Observation's Biodiversity Observation Network (GEO BON) has proposed a global biodiversity observing system (GBIOS), which includes the goal of increasing the capacity to forecast biodiversity change and the loss of ecological and evolutionary resilience³³. GEO BON has also launched a new working group (EcoCode) with the aims of synthesizing biodiversity modelling tools, developing shared platforms and creating a biodiversity model intercomparison platform. These are exciting and

promising steps in the right direction. Even with initiatives like these, large-scale biodiversity forecasting will remain challenging because of the number of species involved and because the drivers can be both more complex and less well measured than those of ecosystem fluxes^{13,34}. However, there is a good understanding of how climate change impacts biodiversity in general (for example, most systems are expected to shift polewards or upslope, with many communities being compressed, eliminated or reorganized in new combinations) and this should be leveraged in scenario analyses and risk assessments related to climate change-based loss and damages^{13,35,36}. In addition, there are many areas where biodiversity forecasts can be developed to address specific international environmental goals. This need for forecasts is not limited to IPBES, but also extends to international efforts around sustainable use and biodiversity (Sustainable Development Goals 12, 14 and 15), threatened and endangered species (such as CITES (Convention on International Trade in Endangered Species) and the IUCN (International Union for Conservation of Nature)), international waters (for example, fisheries, whaling), neutral territories (such as Antarctica and CCAMLR (Commission for the Conservation of Antarctic Marine Living Resources)) and disease (for example, the World Health Organization). Overall, UN programmes and agencies have a unique opportunity to play a central leadership role in advancing ecological forecasting and its application in management decisions.

Efforts to reduce this gap in forecast development and use can be facilitated by combining what is known about how biodiversity responds in general with the available data from targeted systems and using iterative evaluation to refine models and prioritize data needs¹³. This iterative improvement will be most rapid when direct connections with management create learning feedback loops to improve outcomes. While the IPBES effort to explore plausible futures for biodiversity and ecosystem services used a large number of models driven by land-use change and climate change scenarios from integrated assessment models^{37,38}, a gap remains between such long-range projections and near-term forecasts driven by policy-relevant management scenarios³⁹. Scenario-based projections evaluating conservation and management options are particularly important for IPBES and other international conservation efforts because of the critical impacts of human choices on biodiversity preservation, as well as its resulting benefits to people.

One general area where ecological forecasting has the potential to actively inform international decision-making for climate adaptation and mitigation is predicting discrete events (for example, drought, wildfire and other disturbances, disease and pest outbreaks, coastal flooding, coral bleaching) and the time-lagged ecological consequences of event-driven change (such as biological invasions, land-use change, post-disturbance regeneration and restoration). Although ecologists have devoted substantial attention to some of these problems, such as trying to predict which species are likely to be invasive^{40–42} or understanding how ecosystems respond to drought^{43–45}, others are relatively underexplored, such as developing real-time forecasts of forest pests⁴⁶, forecasting the potential impacts of different ecosystem management strategies (for example, restoration, burning, grazing, herbicides, dam releases to rivers)^{15,16} or combining forecasts of multiple ecological variables to evaluate competing management objectives (for example, lake organic carbon storage versus methane production)²². Across this broad range of applications, taking a forecasting approach not only makes our science more robust and repeatable, but can force us to rethink old assumptions (both scientific and management-related) and open up new avenues for exploration and innovation⁴⁷.

Putting ecological forecasting into practice at the international level does not have to be limited to large-scale UN efforts. Numerous conservation organizations, non-governmental organizations (NGOs) and natural-resource industries (such as agriculture, forestry and fisheries) that operate at an international scale would benefit

from forecasts to assist with climate adaptation, the management of threatened, invasive, commercially important or bycatch species, ecosystem restoration, sustainability certification or natural climate solutions. Similarly, as COVID-19 made apparent, emerging infectious diseases are increasingly global in impact and frequently zoonotic in origin. A related emerging idea is forecast-based action, which proactively ties event-driven emergency management plans, including the release of emergency funds, to forecast-based thresholds (for example, climate extremes), rather than acting reactively to disaster events⁴⁸. Forecast-based action approaches have been adopted by international humanitarian organizations (such as the Red Cross) and by the UN itself to reduce the impacts of natural disasters, including many that are exacerbated by climate change and ecological in nature (such as plant drought stress). However, forecast-based action has not yet been adopted by international conservation organizations or applied explicitly to the problem of climate adaptation.

Finally, there are unique global opportunities to help build out ecological forecasting endeavours. Many ecological forecasts already rely heavily on datasets that are international in scope, such as meteorological forecasts, remote sensing data and large-scale community databases (for example, the Global Biodiversity Information Facility, FLUXNET, GLEON), which creates natural pathways for the spatial scaling of forecasts and for further international cooperation on monitoring and data sharing (for example, Group on Earth Observations). Similarly, efforts to build an ecological forecasting community, which includes community coordination on training, tools, standards and synthesis, have primarily been grass-roots operations (see the ‘State of ecological forecasting’), but such efforts are themselves increasingly international in scope. This international growth in focus and investment has not been distributed equally, however. Data volumes, investments in research and development, and existing ecological forecasts are all substantially biased towards the global north (with the important exception of satellite imagery), while the impacts of the dual biodiversity and climate crises are biased towards the global south^{49,50}. Investments in ecological forecasting (development, training, data collection and so on) are critically needed to deliberately address historic and current inequities, both within and across nations, and to support efforts led by Indigenous peoples.

Scientific acceleration and grand challenge questions

The cyclic nature of iterative forecasting aims to establish a learning loop in which predictions can be meaningfully compared with new observations and the models then updated to improve future predictions (Fig. 1). This approach is central to the idea of adaptive management, which emphasizes the importance of evaluating and learning from the outcomes of previous decisions to inform future decisions⁵¹. Similarly, results from numerous disciplines have demonstrated that a learning loop is essential to improve forecasting skill^{52,53}. As such, the ecological forecasting cycle can be a win–win—simultaneously improving decision-making while accelerating scientific understanding. This is possible because well-executed ecological forecasts are specific and quantitative (and thus falsifiable). Forecasts provide continuous real-time feedback, which facilitates rapid testing of the ecological hypotheses embedded in the forecasting models. In many cases the same forecasts can be used to address both decision-focused and basic science questions, helping to align the needs of these two fields of the ecological research community. For example, the Forecasting Lake And Reservoir Ecosystems system is used by drinking-water reservoir managers to improve water quality in the face of non-stationary climate variability, while scientists are using the same forecasts to improve understanding of freshwater ecosystem dynamics^{22,54,55}. More broadly, comparative analyses across forecasts for different systems have the potential to answer grand challenge questions about the predictability of nature^{56–58}: how far into the future can different aspects of nature be




Fig. 2 | Ecological predictability. The predictability of a forecast is measured by (1) the rate at which forecast uncertainty (blue shaded area) grows into the future and (2) the limit at which the forecast performs no better than a historical baseline. The growth in uncertainty is controlled by uncertainty in (and sensitivity to) the initial conditions (IC), exogenous drivers (X), parameter uncertainty (θ), parameter variability (α) and model structural (process) errors (σ). Credit: Kristina Davis.

successfully predicted? Why do some ecological systems respond to climate change more predictably than others? When are ecological forecasts limited by the quality of other forecasts (for example, weather) that are commonly used as model inputs? Is ecological understanding transferable across systems? What do the answers to these questions tell us about the overarching rules and patterns in ecology?

Existing theory about ecological predictability has focused on two key metrics (Fig. 2): (1) the rate at which forecast uncertainty grows with time; and (2) the limit at which forecast accuracy is no better than chance⁵⁹. Our previous work showed that the uncertainty growth rate is a function of uncertainties in five key inputs and the sensitivity of the forecast to each of these inputs⁵⁶. Atmospheric scientists used this type of uncertainty decomposition in the 1960s to determine that weather forecast uncertainty was dominated by uncertainty in the initial conditions⁶⁰, a theoretical advance that has guided large-scale investments (US\$ billions per year) in weather monitoring, modelling and data assimilation that aim to constrain the initial condition uncertainty in each new forecast. These investments have driven decades of continual improvements in the skill, understanding and utility of weather forecasts⁶¹. Ecological forecasting is now poised to make a similar leap in foundational understanding. While some ecological forecasts are (like weather) highly sensitive to initial conditions, others are dominated by sensitivity to the uncertainty in external drivers (for example, climate change), model structural (process) uncertainty, data limitations in constraining model parameters and the inherent variability of biological systems^{30,56,62–65}. Understanding which sources of uncertainty dominate which ecological forecasts is critical to deploying effective monitoring, modelling and model–data integration efforts. Furthermore, unlocking the grand challenge of understanding the patterns of predictability in nature will require a comparative approach across the diversity of ecological systems to quantify which ecological forecasts are limited by which uncertainties⁵⁶. Ultimately, being able to anticipate which ecological systems will be predictable, and what information will be needed to constrain new predictions, will drive improvements in our ability to make decision-relevant forecasts and decision-makers' abilities to make better and more confident decisions.

Learning from meteorology while addressing unique challenges

Ecological forecasting has much to learn from other forecasting communities that are deeply embedded in decision-making. For example, decades of progress in meteorology demonstrate that reliable predictions are possible for complex natural systems if iterative approaches are adopted^{61,66,67}. Meteorology also illustrates the potential for both the open provisioning of public goods (such as climate data,

weather forecasts, severe weather alerts) and private-sector innovation (for example, broadcast meteorology, weather apps, artificial intelligence-augmented forecasts)⁶⁸. Furthermore, weather forecasts are more than just the outputs from numerical models. Multiple times a day, at meteorological centres around the world, weather models are iteratively updated with new data from a globally connected network of satellite and ground sensors, and new predictions are generated from the updated models. These numerical predictions are then interpreted and adjusted by human forecasters and algorithms to produce the forecasts society depends on⁶⁹. Ecological forecasting should build on this framework by continually updating forecasts as new data become available to provide decision-makers with the best available scientific insight, accelerate scientific discovery and improve understanding of ecological predictability more broadly^{61,70}.

Ecological forecasting should also draw inspiration from the early days of numerical weather prediction, when forecast skill was low. Weather forecasters had a choice between stepping back from forecasting until the mechanics of the atmosphere were better understood or stepping forwards into an iterative forecast cycle of learning by doing⁶⁶. By stepping forwards, they achieved a critical win–win of relentless improvements in theory, skill and utility to society⁹. Ecology is now at a similar crossroads. Society is facing climate and biodiversity crises, but finally has the observations, models, theory and prototype iterative forecasts to improve decisions, both big and small⁸.

Compared with other forecasting fields, however, ecological forecasting presents unique challenges. Whereas a weather model has 10–20 state variables that follow the same physical equations throughout the atmosphere, there are millions of species on the planet and individual organisms vary (both within and between species) in the primary drivers of their dynamics and their responses to those drivers. Organisms are heterogeneous across space, time and phylogeny in how they respond to changes in climate and human activities, and those responses are continually adapting and evolving across a cascade of different timescales. Even if there is only interest in forecasting emergent ecosystem processes (such as carbon, water, and nutrient cycling), biodiversity causes the parameters in the equations, and sometimes the equations themselves, to continuously change across space and time. However, this variability is not unconstrained, nor wholly unpredictable. As the data necessary to make forecasts are only available for a fraction of the world's species, comparative analyses of predictability and forecast transferability (for example, how parameters and equations change across space, time and taxa) are critical to the advancement of ecological forecasting. Beyond being important to theory, sharing information across forecasts of different systems (through hierarchical models of across-species parameter variability, for example) lowers the amount of data necessary to make predictions, allowing the scope of ecological forecasting to be extended. This can be particularly important for rare and novel (for example, invasive) taxa.

The inherent complexity of biodiversity introduces further challenges in ecological forecasting related to both monitoring requirements and the infrastructure for producing forecasts. Weather forecasting represents a single large forecasting problem that nations have addressed by constructing large centres with dedicated staff, models and both physical infrastructure and cyberinfrastructure. Ecological forecasts represent a spectrum of forecasting challenges that vary in size (a single global forecast to many local forecasts), approach (process-based, statistical, machine learning), system (terrestrial, freshwater, marine) and biological scale (physiological, organismal, population, community, ecosystem/biogeochemistry). The diversity of forecasts has resulted in numerous unique data processing and forecasting workflows that require non-trivial costs and expertise to build and maintain^{22,71–74}. That said, these challenges also represent opportunities to use forecasts to optimize and iteratively adapt monitoring programmes, to develop new sensor technologies and to better integrate ecological monitoring and infrastructure^{75–77}. The expansion

of ecological forecasting has delivered progress and opportunities to leverage economies of scale through reusable community standards⁷⁸, workflows, models and cyberinfrastructure^{49,71} and to reduce the costs, time and learning curve involved in launching and maintaining forecasts. These advances will not be limited to near-term forecasts, but will simultaneously increase the capacity for, and confidence in, ecological forecasting on climate timescales.

Ecological forecasting is also generating opportunities to improve forecasting across disciplines. The challenges of forecasting ecological systems mean that researchers need to revisit theoretical assumptions around predictability⁵⁶ and the data assimilation tools used to integrate new observations into forecasts³⁰ while acknowledging that a wider range of uncertainties must also now be accounted for (Fig. 2). Hybrid approaches that combine traditional process-based models and data assimilation with newer machine learning methods are advancing rapidly in both Earth system and ecological forecasting^{79–83}. Ecological forecasters are important consumers of forecasts (atmosphere, oceans and so on), translating these physical forecasts into information about Earth's life support systems, on which humanity depends. The biosphere also generates feedbacks to other parts of the Earth system, such that improvements in ecological forecasting will improve Earth system predictions of the boundary conditions used in both weather and climate models^{29,84,85}.

State of the ecological forecasting community

Dramatic improvements in ecological forecasting are now emerging that were unimaginable even a decade ago. Advances in sensor technologies, satellites and genomics provide access to unprecedented volumes of environmental data that are born-digital and increasingly near real time. The urgency of climate change has driven shifts towards large-scale networked science, which in turn has made data access more standardized and equitable—both in terms of top-down international observatories⁸⁶ (such as NEON, Australia's Terrestrial Ecosystem Research Network (TERN), the South African Environmental Observation Network (SAEON) and the International Long Term Ecological Research network (ILTER)) and bottom-up globally distributed experiments with standardized protocols (NutNet^{87,88}, DroughtNet⁸⁹, Cellulose Decomposition Experiment (CELLDEX)⁹⁰ and so on). At the same time, there have been rapid advances in computational methods and cyberinfrastructure, including revolutions in artificial intelligence and machine learning^{91,92}, workflow containerization⁹³, distributed cloud computing and cloud-native data storage. Collectively, these advances have fuelled the recent growth of ecological forecasting and herald opportunities for further increases in scope and reach.

To leverage these technical advances, ecological forecasting needs to build coherent communities of practice to support its growth as a discipline⁹⁴. Successful ecological forecasts represent a convergence of expertise across a range of ecological subdisciplines, the social and decision sciences, physical environmental sciences, computational and data sciences and statistics⁸. However, each discipline operates under different norms and approaches, as do the different sectors (academia, agencies, industry, NGOs and so on) involved in forecast research, development and operationalization. Overcoming the resulting barriers will require: educational efforts designed to broaden disciplinary backgrounds and train researchers in interdisciplinary collaboration^{95,96}, transdisciplinary organizations and teams with the time and commitment to develop shared interests and vocabularies⁹⁷ and funding mechanisms to support social science and non-academic partner collaborations⁹⁸. Expanding the ecological forecasting community in a way that includes everyone influenced by climate change will also require actively addressing issues of equity—including who has access to ecological forecasting tools and data streams, who can interpret and use them for decision-making⁹⁹ and whose perspectives and priorities are incorporated into their development and dissemination¹⁰⁰.

This sort of community building has accelerated recently through efforts such as the Ecological Forecasting Initiative (EFI), an international grass roots consortium aimed at fostering a community of practice around ecological forecasting⁸. EFI has engaged thousands of academic, agency, NGO and industry scientists and partners through a broad mix of international chapters and conferences, working groups, webinars, articles, videos, standards⁷⁸, policy briefs, training opportunities (including minority-serving institution student mentoring and faculty partnerships) and contributions to government reports^{16,67,98,101,102}. Since 2021, EFI's Research Coordination Network has hosted an ongoing NEON Forecasting Challenge^{72,103,104} with the goal of predicting data before it is collected. Over 200 teams (including 11 university courses and 2 minority-serving institution mentoring programmes) have participated and the Challenge has developed new educational resources and community cyberinfrastructure^{105,106}, as well as comparative analyses that help tackle grand challenge questions^{103,104}. EFI's progress is indicative of a new and growing discipline in which the number and diversity of forecasts, forecasters, decision scientists and end users could be rapidly scaled up to help address the climate and biodiversity crises.

A path forwards

Responding to pressing global environmental challenges in an era of climate change will require substantial local and global development to scale up the ecological forecasting enterprise. Rapid progress has been made, particularly over the past 5 years as efforts to bring together the community have accelerated. Meeting this challenge requires an intentional and inclusive approach to build the human dimensions of our forecasting capability.

The translation of forecasts from research to societal impacts has required new cross-sector partnerships that span academia, governmental agencies, industry, NGOs and other interested parties to be established. Furthermore, the technical requirements for building operational ecological forecasts often exceed the capacity of governmental agencies (the traditional producers of forecasts). For example, the soil carbon monitoring being conducted by the nascent carbon credit industry (which serves as both a key input and validation of their carbon forecasts) to inform climate mitigation is already on track to surpass the data volumes in government agency soil maps. While efforts like EFI have laid the groundwork for building bridges across sectors (for example, through interagency and cross-sector workshops and trainings), in scaling up ecological forecasting there is an important need to foster even greater innovation and engagement across a broader spectrum of partners, end users and decision-makers. Key to these partnerships is the idea of co-production: groups that will use forecasts need to be engaged in the process of launching new ecological forecasts from the outset by informing and contributing to the goals, approaches and product design^{107–109}. These partnerships and the associated scaling activities must acknowledge historical biases in participation and perspectives (for example, towards the global north) and actively promote equity in participation and recognition of marginalized perspectives going forwards. Such co-production is necessary to ensure that forecasts are equitable, useful, usable and credible¹¹⁰.

One key part of community building is engaging groups that have been traditionally excluded from both science and decision-making. For example, scientific research funding in the United States goes disproportionately to white principal investigators¹¹¹ and racial minority groups earn a disproportionately low fraction of US Earth science doctorates¹¹². At the same time, many of the world's environmental problems disproportionately impact members of marginalized groups, including Indigenous communities and the urban poor^{6,113}. Furthermore, while ecological forecasting is global in scope, low- and middle-income countries are under-represented in both research and community participation despite absorbing a disproportionately large share of the world's biodiversity, carbon storage and climate impacts^{49,50}. Active investment by international bodies in the building of ecological forecasting

capabilities in the global south, much as the World Meteorological Organization has historically done for meteorology¹¹⁴, would help to address both these inequities and international climate, biodiversity and sustainability goals. It is essential to actively ensure that our work not only broadens participation among these groups, but also helps identify and address the underlying structural issues that create and sustain their under-representation in the first place.

Increasing forecasting capacity will also require changes in training across a wide range of sectors, from boosting the number of individuals with the technical expertise to produce usable forecasts¹¹⁵, to training the next generation of managers to better use forecasts¹¹⁶, to using questions of prediction to increase the scientific literacy of the broader public. Training in ecological forecasting also goes beyond technical topics such as modelling and data science. It requires interdisciplinary teams that are able to integrate expert knowledge about a specific ecological process, the social context of the decisions being informed by the forecast, the decision science frameworks for making these decisions and the legal and ethical questions about what should and should not be forecast¹¹⁷. Examples of such training efforts have grown rapidly in the past 5 years, ranging from introductory ecology courses taught from a predictive perspective, to minority-serving institution mentoring in environmental data science, to dedicated upper-level forecasting courses to academic and non-academic workshops. Moving forwards, our universities and professional societies (both research and management) can help by making such training more broadly and equitably available, including opportunities for learners without the ability to travel to in-person opportunities and those facing technological challenges, such as a lack of stable internet access¹⁰⁰.

Community building is also key to advancing theory and technology. Comparative analyses are central to understanding patterns of predictability and transferability, and to improving forecasts by sharing information across systems and species about model structure and parameters. Syntheses beyond the NEON Challenge are currently limited by the number and diversity of forecasts available and by the forecasts that do exist not being sufficiently catalogued, archived and interoperable for synthetic analysis^{56,57,78}. Achieving the economies of scale required to generate such a broad catalogue of forecasts, as well as to respond to global environmental challenges, depends on the development and adoption of community conventions surrounding shared tools and cyberinfrastructure and the development of community norms around using them. The expansion of such technologies should move beyond academia to a model where agencies and industry play a central role in their co-production and adoption.

In conclusion, ecological forecasting is at a critical point for future growth, similar to weather forecasting in the twentieth century, and cannot afford to step back. Stepping forwards requires a quantum leap in forecasting capacity, game-changing scientific breakthroughs and technological developments and a new twenty-first-century vision of data-driven environmental management. The nations of the world, along with UN bodies, major international corporations and NGOs, can help by integrating ecological forecasting into their climate adaptation and mitigation strategies. At the same time, the scientific community, spanning academia, agencies and industry, can help build capacity to respond to this urgent need. Ecological forecasting simultaneously offers a new set of tools to advance these efforts and a new frontier of discovery about how nature works. Making forecasts of nature mainstream, and developed by diverse sections of society, will generate the foresight to mitigate further degradation and enable us to proactively build a future that is climate resilient and nurturing for people and the planet.

References

- Pulgar-Vidal, M. Applying the lessons of climate change to halting biodiversity loss. *IUCN* <https://www.iucn.org/crossroads-blog/202207/applying-lessons-climate-change-halting-biodiversity-loss> (2022).
- Costanza, R. et al. Changes in the global value of ecosystem services. *Glob. Environ. Change* **26**, 152–158 (2014).
- IPCC Special Report on Climate Change and Land (eds Shukla, P. R. et al.) (WMO, 2019).
- Brondizio, E. S. et al. (eds) *Global Assessment Report on Biodiversity and Ecosystem Services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services* (IPBES, 2019).
- Bergstrom, D. M. et al. Combating ecosystem collapse from the tropics to the Antarctic. *Glob. Change Biol.* **27**, 1692–1703 (2021).
- The Global Risks Report 2023* (World Economic Forum, 2023).
- Milly, P. C. D. et al. Stationarity is dead: whither water management? *Science* **319**, 573–574 (2008).
- Dietze, M. C. et al. Iterative near-term ecological forecasting: needs, opportunities, and challenges. *Proc. Natl Acad. Sci. USA* **115**, 1424–1432 (2018).

This study proposes a community roadmap for the development of iterative ecological forecasts.

- Dietze, M. C. *Ecological Forecasting* (Princeton Univ. Press, 2017).
- Tulloch, A. I. T., Hagger, V. & Greenville, A. C. Ecological forecasts to inform near-term management of threats to biodiversity. *Glob. Change Biol.* **26**, 5816–5828 (2020).
- Malhi, Y. et al. Climate change and ecosystems: threats, opportunities and solutions. *Phil. Trans. R. Soc. B* **375**, 20190104 (2020).
- Trisos, C. H., Merow, C. & Pigot, A. L. The projected timing of abrupt ecological disruption from climate change. *Nature* **580**, 496–501 (2020).
- Urban, M. C. et al. Improving the forecast for biodiversity under climate change. *Science* **353**, aad8466 (2016).
- This paper highlights the challenges in forecasting biodiversity.**
- Arora, V. K. et al. Carbon–concentration and carbon–climate feedbacks in CMIP6 models and their comparison to CMIP5 models. *Biogeosciences* **17**, 4173–4222 (2020).
- Bradford, J. B., Betancourt, J. L., Butterfield, B. J., Munson, S. M. & Wood, T. E. Anticipatory natural resource science and management for a changing future. *Front. Ecol. Environ.* **16**, 295–303 (2018).
- Bradford, J. B. et al. *Ecological Forecasting—21st Century Science for 21st Century Management* Open-File Report 2020–1073 (US Geological Survey, 2020); <https://doi.org/10.3133/ofr20201073>
- Hartman, M. D. et al. Seasonal grassland productivity forecast for the U.S. Great Plains using Grass-Cast. *Ecosphere* **11**, e03280 (2020).
- Pringle, M. J. et al. Using remote sensing to forecast forage quality for cattle in the dry savannas of northeast Australia. *Ecol. Indic.* **133**, 108426 (2021).
- Zhang, B. & Carter, J. FORAGE – an online system for generating and delivering property-scale decision support information for grazing land and environmental management. *Comput. Electron. Agric.* **150**, 302–311 (2018).
- Welch, H. et al. Practical considerations for operationalizing dynamic management tools. *J. Appl. Ecol.* **56**, 459–469 (2019).
- Scales, K. L. et al. Fit to predict? Eco-informatics for predicting the catchability of a pelagic fish in near real time. *Ecol. Appl.* **27**, 2313–2329 (2017).
- Carey, C. C. et al. Advancing lake and reservoir water quality management with near-term, iterative ecological forecasting. *Inland Waters* **12**, 107–120 (2022).
- Lofton, M. E., Howard, D. W., Thomas, R. Q. & Carey, C. C. Progress and opportunities in advancing near-term forecasting of freshwater quality. *Glob. Change Biol.* **29**, 1691–1714 (2023).
- Soil Enrichment Protocol Version 1.1: Reducing Emissions and Enhancing Soil Carbon Sequestration on Agricultural Lands* (Climate Action Reserve, 2022); <https://www.climateactionreserve.org/how/protocols/soil-enrichment/>

25. Woolway, R. I. et al. Lake heatwaves under climate change. *Nature* **589**, 402–407 (2021).
26. Swain, D. L., Singh, D., Touma, D. & Diffenbaugh, N. S. Attributing extreme events to climate change: a new frontier in a warming world. *One Earth* **2**, 522–527 (2020).
27. IPCC *Climate Change 2014: Synthesis Report* (eds Core Writing Team, Pachauri, R. K. & Meyer, L. A.) (IPCC, 2014); <https://www.ipcc.ch/report/ar5/syr>
28. Fisher, R. A. & Koven, C. D. Perspectives on the future of land surface models and the challenges of representing complex terrestrial systems. *J. Adv. Model. Earth Syst.* **12**, e2018MS001453 (2020).
29. Fisher, R. A. et al. Vegetation demographics in Earth system models: a review of progress and priorities. *Glob. Change Biol.* **24**, 35–54 (2018).
30. Raiho, A. et al. Determinants of predictability in multi-decadal forest community and carbon dynamics. Preprint at *bioRxiv* <https://doi.org/10.1101/2020.05.05.079871> (2020).
31. Egerton, P. et al. *Early Warnings for All: The UN Global Early Warning Initiative for the Implementation of Climate Adaptation* (World Meteorological Organization, 2022); <https://library.wmo.int/idurl/4/58209>
32. Shin, Y.-J. et al. in *Global Assessment Report on Biodiversity and Ecosystem Services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services* (eds Brondizio, E. S. et al.) Ch. 4 (IPBES, 2019); <https://doi.org/10.5281/zenodo.5656910>
33. Gonzalez, A. et al. A global biodiversity observing system to unite monitoring and guide action. *Nat. Ecol. Evol.* **7**, 1947–1952 (2023).
34. Briscoe, N. J. et al. Forecasting species range dynamics with process-explicit models: matching methods to applications. *Ecol. Lett.* **22**, 1940–1956 (2019).
35. Briscoe, N. J. et al. Mechanistic forecasts of species responses to climate change: the promise of biophysical ecology. *Glob. Change Biol.* **28**, 1451–1470 (2023).
36. Botkin, D. B. et al. Forecasting the effects of global warming on biodiversity. *BioScience* **57**, 227–236 (2007).
37. Kim, H. et al. A protocol for an intercomparison of biodiversity and ecosystem services models using harmonized land-use and climate scenarios. *Geosci. Model Dev.* **11**, 4537–4562 (2018).
38. Pereira, H. M. et al. Global trends and scenarios for terrestrial biodiversity and ecosystem services from 1900 to 2050. *Science* **384**, 458–465 (2024).
39. Rosa, I. M. D. et al. Challenges in producing policy-relevant global scenarios of biodiversity and ecosystem services. *Glob. Ecol. Conserv.* **22**, e00886 (2020).
40. Moles, A. T., Gruber, M. A. M. & Bonser, S. P. A new framework for predicting invasive plant species. *J. Ecol.* **96**, 13–17 (2008).
41. Van Kleunen, M., Dawson, W., Schlaepfer, D., Jeschke, J. M. & Fischer, M. Are invaders different? A conceptual framework of comparative approaches for assessing determinants of invasiveness. *Ecol. Lett.* **13**, 947–958 (2010).
42. Ibáñez, I. et al. Integrated assessment of biological invasions. *Ecol. Appl.* **24**, 25–37 (2014).
43. Konings, A. G. et al. Detecting forest response to droughts with global observations of vegetation water content. *Glob. Change Biol.* **27**, 6005–6024 (2021).
44. Anderegg, W. R. L., Trugman, A. T., Badgley, G., Konings, A. G. & Shaw, J. Divergent forest sensitivity to repeated extreme droughts. *Nat. Clim. Change* **10**, 1091–1095 (2020).
45. Vose, J., Clark, J. S., Luce, C. & Patel-Weyand, T. *Effects of Drought on Forests and Rangelands in the United States: A Comprehensive Science Synthesis General Technical Report WO-93b* (Forest Service, US Department of Agriculture, 2016).
46. Jepsen, J. U., Vindstad, O. P. L. & Ims, R. A. Spatiotemporal dynamics of forest geometrid outbreaks. *Curr. Opin. Insect Sci.* **55**, 100990 (2023).
47. Wheeler, K. I. *Cold-Deciduous Broadleaf Phenology: Monitoring Using a Geostationary Satellite and Predicting Using Trigger-Less Dynamic Models* (Boston Univ., 2022).
48. Boulton, V. L. Forecast-based action for conservation. *Conserv. Biol.* <https://doi.org/10.1111/cobi.14054> (2023).
This paper highlights the potential ecological applications of forecast-based action (the proactive initiation of action plans based on forecast thresholds).
49. Slingsby, J. A., Wilson, A. M., Maitner, B. & Moncrieff, G. R. Regional ecological forecasting across scales: a manifesto for a biodiversity hotspot. *Methods Ecol. Evol.* <https://doi.org/10.1111/2041-210X.14046> (2023).
50. Schimel, D. et al. Observing terrestrial ecosystems and the carbon cycle from space. *Glob. Change Biol.* **21**, 1762–1776 (2015).
51. Walters, C. J. *Adaptive Management of Renewable Resources* (Macmillan, 1986).
52. Tetlock, P. E. & Gardner, D. *Superforecasting: The Art and Science of Prediction* (Crown, 2015).
53. Silver, N. *The Signal and the Noise* (Penguin, 2012).
54. Thomas, R. Q. et al. A near-term iterative forecasting system successfully predicts reservoir hydrodynamics and partitions uncertainty in real time. *Water Resour. Res.* **56**, e2019WR026138 (2020).
55. Thomas, R. Q. et al. Near-term forecasts of NEON lakes reveal gradients of environmental predictability across the U.S. *Front. Ecol. Environ.* **21**, 220–226 (2023).
56. Dietze, M. C. Prediction in ecology: a first-principles framework. *Ecol. Appl.* **27**, 2048–2060 (2017).
This paper derives theoretical expectations for how different uncertainties (model, parameters, initial conditions, inputs) affect ecological predictability, and how this differs from weather forecasting.
57. Lewis, A. et al. The power of forecasts to advance ecological theory. *Methods Ecol. Evol.* **14**, 746–756 (2022).
58. Lewis, A. S. L. et al. Increased adoption of best practices in ecological forecasting enables comparisons of forecastability. *Ecol. Appl.* <https://doi.org/10.1002/eap.2500> (2021).
A meta-analysis on best practice adoption that also provides some high-level across-forecast syntheses concerning patterns of predictability.
59. Petchey, O. L. et al. The ecological forecast horizon, and examples of its uses and determinants. *Ecol. Lett.* **18**, 597–611 (2015).
This study discusses how far into the future useful ecological forecasts can be made and which system properties may affect this.
60. Lorenz, E. N. Deterministic nonperiodic flow. *J. Atmos. Sci.* **20**, 130–141 (1963).
61. Bauer, P., Thorpe, A. & Brunet, G. The quiet revolution of numerical weather prediction. *Nature* **525**, 47–55 (2015).
62. Lovenduski, N. S., Bonan, G. B., Yeager, S. G., Lindsay, K. & Lombardo, D. L. High predictability of terrestrial carbon fluxes from an initialized decadal prediction system. *Environ. Res. Lett.* **14**, 124074 (2019).
63. Lovenduski, N. S. & Bonan, G. B. Reducing uncertainty in projections of terrestrial carbon uptake. *Environ. Res. Lett.* **12**, 044020 (2017).
64. Bonan, G. B. et al. Model structure and climate data uncertainty in historical simulations of the terrestrial carbon cycle (1850–2014). *Glob. Biogeochem. Cycles* **33**, 1310–1326 (2019).
65. Luo, Y., Keenan, T. F. & Smith, M. Predictability of the terrestrial carbon cycle. *Glob. Change Biol.* **21**, 1737–1751 (2015).

66. Shuman, F. G. History of numerical weather prediction at the National Meteorological Center. *Weather Forecast.* **4**, 286–296 (1989).
67. Droegeemeier, K. K. et al. *Earth System Predictability Research and Development Strategic Framework and Roadmap* (National Science & Technology Council, 2020).
68. Lewis, M. *The Coming Storm* (Audible Originals, LLC, 2018).
69. Hoffman, R. R., LaDue, D. S., Mogil, H. M., Roebber, P. J. & Trafton, J. G. (eds) *Minding the Weather: How Expert Forecasters Think* (MIT Press, 2017).
70. Liang, X.-Z. et al. DAWN: Dashboard for Agricultural Water use and Nutrient management—a predictive decision support system to improve crop production in a changing climate. *Bull. Am. Meteorol. Soc.* <https://doi.org/10.1175/BAMS-D-22-0221.1> (2024).
71. Fer, I. et al. Beyond ecosystem modeling: a roadmap to community cyberinfrastructure for ecological data-model integration. *Glob. Change Biol.* **27**, 13–26 (2021).
72. Thomas, R. Q. et al. The NEON Ecological Forecasting Challenge. *Front. Ecol. Environ.* **21**, 112–113 (2023).
A common set of forecasting challenges, as highlighted in this paper, can help build cohesion in the field of ecological forecasting, allowing the development and sharing of best practices and innovation.
73. White, E. P. et al. Developing an automated iterative near-term forecasting system for an ecological study. *Methods Ecol. Evol.* **10**, 332–344 (2019).
This study on ecological forecasting cyberinfrastructure provides an open-source automated workflow and discusses best practices.
74. Yenni, G. M. et al. Developing a modern data workflow for regularly updated data. *PLoS Biol.* **17**, e3000125 (2019).
75. McCord, S. E. & Pilliod, D. S. Adaptive monitoring in support of adaptive management in rangelands. *Rangelands* **44**, 1–7 (2022).
76. Lindenmayer, D. B. & Likens, G. E. Adaptive monitoring: a new paradigm for long-term research and monitoring. *Trends Ecol. Evol.* **24**, 482–486 (2009).
77. Zeng, X. et al. Use of observing system simulation experiments in the United States. *Bull. Am. Meteorol. Soc.* **101**, E1427–E1438 (2020).
78. Dietze, M. C. et al. A community convention for ecological forecasting: output files and metadata version 1.0. *Ecosphere* **14**, e4686 (2023).
79. Luo, Y. & Smith, B. (eds) *Land Carbon Cycle Modeling: Matrix Approach, Data Assimilation, & Ecological Forecasting* (CRC, 2022).
80. Zwart, J. A. et al. Near-term forecasts of stream temperature using deep learning and data assimilation in support of management decisions. *J. Am. Water Resour. Assoc.* **59**, 317–337 (2023).
81. Grover, A., Kapoor, A. & Horvitz, E. A deep hybrid model for weather forecasting. In *Proc. 21st ACM SIGKDD International Conference on Knowledge Discovery and Data Mining* 379–386 (Association for Computing Machinery, 2015).
82. Fathi, M., Haghi Kashani, M., Jameii, S. M. & Mahdipour, E. Big data analytics in weather forecasting: a systematic review. *Arch. Comput. Methods Eng.* **29**, 1247–1275 (2022).
83. Arcomano, T. et al. A machine learning-based global atmospheric forecast model. *Geophys. Res. Lett.* **47**, e2020GL087776 (2020).
84. Bonan, G. *Ecological Climatology: Concepts and Applications* (Cambridge Univ. Press, 2015).
85. Khanna, J., Medvigy, D., Fueglistaler, S. & Walko, R. Regional dry-season climate changes due to three decades of Amazonian deforestation. *Nat. Clim. Change* **7**, 200–204 (2017).
86. National Research Council *NEON: Addressing the Nation's Environmental Challenges* (National Academies, 2004).
87. Adler, P. B. et al. Productivity is a poor predictor of plant species richness. *Science* **333**, 1750–1753 (2011).
88. Fraser, L. H. et al. Coordinated distributed experiments: an emerging tool for testing global hypotheses in ecology and environmental science. *Front. Ecol. Environ.* **11**, 147–155 (2013).
89. Knapp, A. K. et al. Pushing precipitation to the extremes in distributed experiments: recommendations for simulating wet and dry years. *Glob. Change Biol.* **23**, 1774–1782 (2017).
90. Tiegs, S. D. et al. Global patterns and drivers of ecosystem functioning in rivers and riparian zones. *Sci. Adv.* **5**, eaav0486 (2019).
91. Goodfellow, I., Bengio, Y. & Courville, A. *Deep Learning* (MIT Press, 2016).
92. Brisimi, T. S. et al. Federated learning of predictive models from federated electronic health records. *Int. J. Med. Inform.* **112**, 59–67 (2018).
93. Park, S. et al. FaaSr: cross-platform function-as-a-service serverless scientific workflows in R. In *Proc. 20th IEEE International Conference on e-Science* 1–10 (IEEE, 2024).
94. Woelmer, W. M. et al. Ten simple rules for training yourself in an emerging field. *PLoS Comput. Biol.* **17**, e1009440 (2021).
95. Farrell, K. J. et al. Training macrosystems scientists requires both interpersonal and technical skills. *Front. Ecol. Environ.* **19**, 39–46 (2021).
96. Vogler, J. S. et al. The hard work of soft skills: augmenting the project-based learning experience with interdisciplinary teamwork. *Instr. Sci.* **46**, 457–488 (2018).
97. National Research Council *Enhancing the Effectiveness of Team Science* (National Academies, 2015).
98. *Next Generation Earth Systems Science at the National Science Foundation* (National Academies, 2021).
99. Kenney, M. A., Gerst, M. D. & Read, E. The usability gap in water resources open data and actionable science initiatives. *J. Am. Water Resour. Assoc.* **60**, 1–8 (2024).
This decision support paper describes a generalized approach to user testing and presents an expansion of FAIR standards to include human access and usability of decision support tools.
100. Willson, A. M. et al. Assessing opportunities and inequities in undergraduate ecological forecasting education. *Ecol. Evol.* **13**, e10001 (2023).
101. Geller, G. et al. *NASA Biological Diversity and Ecological Forecasting: Current State of Knowledge and Considerations for the Next Decade* (NASA, 2022).
102. Williams, C. et al. *2021 NACP Science Implementation Plan Report of the North American Carbon Program* (US Carbon Cycle Science Program, 2021); <https://doi.org/10.5065/kwe1-w815>
103. Wheeler, K. I. et al. Predicting spring phenology in deciduous broadleaf forests: NEON phenology forecasting community challenge. *Agric. For. Meteorol.* **345**, 109810 (2024).
104. Olsson, F. et al. What can we learn from 100,000 freshwater forecasts? A synthesis from the NEON Ecological Forecasting Challenge. *ESS Open Archive* <https://doi.org/10.22541/essoar.171458144.44104603/v1> (2024).
105. Thomas, R. Q. et al. Ecological forecasting initiative: NEON Ecological Forecasting Challenge documentation V1.0. *Zenodo* <https://doi.org/10.5281/zenodo.4780155> (2021).
106. Boettiger, C., Thomas, Q., Laney, C. & Lunch, C. neonstore: NEON Data Store. R package version 0.5.1 <https://CRAN.R-project.org/package=neonstore> (2024).
107. Meadow, A. M. et al. Moving toward the deliberate coproduction of climate science knowledge. *Weather Clim. Soc.* **7**, 179–191 (2015).

108. Lemos, M. C. & Morehouse, B. J. The co-production of science and policy in integrated climate assessments. *Glob. Environ. Change* **15**, 57–68 (2005).
109. Bremer, S. & Meisch, S. Co-production in climate change research: reviewing different perspectives. *WIREs Clim. Change* **8**, e482 (2017).
110. Kirchhoff, C. J., Carmen Lemos, M. & Dessai, S. Actionable knowledge for environmental decision making: broadening the usability of climate science. *Annu. Rev. Environ. Resour.* **38**, 393–414 (2013).
111. Chen, C. Y. et al. Systemic racial disparities in funding rates at the National Science Foundation. *eLife* **11**, e83071 (2022).
112. Bernard, R. E. & Cooperdock, E. H. G. No progress on diversity in 40 years. *Nat. Geosci.* **11**, 292–295 (2018).
This paper highlights the persistent under-representation of racial and minority ethnic groups in US geosciences and recommends actions to spur systemic change.
113. Howell, J. & Elliott, J. R. Damages done: the longitudinal impacts of natural hazards on wealth inequality in the United States. *Soc. Probl.* **66**, 448–467 (2019).
114. *Capacity Development Programme* (World Meteorological Organization, 2023); <https://wmo.int/capacity-development-programme>
115. Emery, N. C. et al. Data science in undergraduate life science education: a need for instructor skills training. *BioScience* <https://doi.org/10.1093/biosci/biab107> (2021).
116. Allum, N., Besley, J., Gomez, L. & Brunton-Smith, I. Disparities in science literacy. *Science* **360**, 861–862 (2018).
117. Hobday, A. J. et al. Ethical considerations and unanticipated consequences associated with ecological forecasting for marine resources. *ICES J. Mar. Sci.* **76**, 1244–1256 (2019).
Forecasting ecological dynamics has societal impacts that need to be considered at all stages of forecast development and dissemination.

Acknowledgements

This work was supported by the NSF Research Coordination Network under grant number 1926388 and an Alfred P. Sloan Foundation grant. We thank K. Davis at Notre Dame for her work on figure development. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the US Government.

Author contributions

M.C.D. organized and led the writing of the Perspective. All authors contributed to writing the original draft, reviewing and editing.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence should be addressed to Michael Dietze.

Peer review information *Nature Climate Change* thanks Vicky Boulton, Matteo Convertino, David Garcia-Callejas and the other, anonymous, reviewer(s) for their contribution to the peer review of this work.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

© Springer Nature Limited 2024