

VI. LITERATURE CITED

- Alliance for the Chesapeake Bay. 1993. Nutrients and the Chesapeake: Refining the bay cleanup effort.
- American Oyster, Chesapeake Bay Foundation, (2002), [Online], http://www.chesapeakebay.net/info/american_oyster.cfm, (verified 7 March 2003)
- American Oceans, Campaign Florida, Briefing Book – Shellfish and Coastal Polluted Runoff, (2002), [Online], <http://www.americoceans.org/florida/chapter2.htm>, (verified 21 March 2002)
- Bacterial Indicators, Canadian Shellfish Quality Resource, (2001), [Online], <http://www.shellfishquality.ca/indicators.htm>, (verified 16 January 2002)
- Bernhard, A. and Field, K.G. (2000a) Identification of nonpoint sources of fecal pollution in coastal waters by using host-specific 16S ribosomal DNA genetic markers from fecal anaerobes. *Applied and Environmental Microbiology* 66, 1587-1594.
- Bernhard, A. and Field, K.G. (2000b) A PCR assay to discriminate human and ruminant feces based on host differences in *Bacteroides-Prevotella* 16S ribosomal DNA. *Applied and Environmental Microbiology* 66, 4571-4574.
- Bower, R. J. (2001) Fecal source identification using antibiotic resistance analysis. *Puget Sound Notes* 45, 3-8.
- Cape Cod Commission, Coastal Embayment Project, (1998), [Online], <http://www.capecodcommission.org/water/overview.htm>, (verified 16 January 2002)
- Carson, A.C., Shear, B.L., Ellersieck, M.R. and Asfaw, A. (2001) Identification of fecal *Escherichia coli* from humans and animals by ribotyping. *Applied and Environmental Microbiology* 67, 1503-1507.
- Dombek, P.E., Johnson, L.K., Zimmerley, S.T. and Sadowsky, M.J. (2000) Use of Repetitive DNA Sequences and the PCR to differentiate *Escherichia coli* isolates from human and animal sources. *Applied and Environmental Microbiology* 66, 2572-2577.
- Friends of the Rivers of Virginia (FORVA). (2001) *State of Our Rivers Report for the Commonwealth of Virginia*. Roanoke, Va: FORVA.
- Food and Drug Administration, FDA, (2002), [Online], <http://um.cfsan.fda.gov/~mow/chap37.html>, (verified 31 January 2002)
- Formiga-Cruz, Tofino-Quesada, G., Bofill-Mas, S., Lees, D.N., Henshilwood, K., Allard A.K., Conden-Hansson, A.C., Hernroth, B.E., Vantarakis, A., Tsibouxi, A., Papapetropoulou, M., Furones, M.D., Girones, R., (2002), Distribution of Human Viruses in Shellfish from Different Growing Areas in Greece, Spain, Sweden, and the United Kingdom, *Applied and Environmental Microbiology* 68.12,5990-5998
- Graves, A.K., Hagedorn, C., Teetor, A., Mahal, M., Booth, A.M., and Reneau, R. B., Jr. (2002) Antibiotic resistance profiles to determine sources of fecal contamination in a rural Virginia watershed. *J. Environmental Quality* 31, 1300-1308.
- Hagedorn, C., Robinson, S.L., Filtz, J.R., Grubbs, S.M., Angier, T.A. and Reneau, R.B. Jr. (1999) Using antibiotic resistance patterns in the fecal streptococci to determine sources of fecal pollution in a rural Virginia watershed. *Applied and Environmental Microbiology* 65, 5522-5531.
- Hagedorn, C., Bacterial Source Tracking in Total Maximum Daily Load Process, (2002)

- Hager, M.C., Detecting Bacteria in Coastal Waters, Stormwater, The Journal for Surface Water Quality Professionals, November 2002
- Hartel, P.G., Segars, W.I., Stern, N.J., Steiner, J. and Buchan, A. (1999) Ribotyping to determine host origin of *Escherichia coli* isolates in different water samples. In *American Water Resources Association (AWRA) Technical Publication Series TPS-99-3p. Wildland Hydrology*. eds. Olsen D.S. and Potyondy J.P. pp. 377-382. Herndon, Va: AWRA.
- Harwood, V. J., Whitlock, J. and Withington, V.H., (2000) Classification of the antibiotic resistance patterns of indicator bacteria by discriminant analysis: Use in predicting the source of fecal contamination in subtropical Florida waters. *Applied and Environmental Microbiology* 66, 3698-3704.
- Hsu, Fu-Chih, Carol-Shieh, Y.S., van Duin, J., Beekwilder, M.J. and Sobsey, M.D. (1995) Genotyping male-specific RNA coliphages by hybridization with oligonucleotide probes. *Applied and Environmental Microbiology* 61, 3960-3966.
- Infoplease, Online Dictionary, Internet Encyclopedia, Atlas, and Almanac Reference, (2002), [Online], <http://www.infoplease.com/ce6/us/a08972022.htm>, (verified 21 March 2002)
- Jersey Shore, Seafood from the Garden State, (2002), [Online], <http://www.state.nj.us/agriculture/rural/seafood/>, (verified 21 March 2002)
- Marine Biotoxins Program, Coastal Research, (2002), [Online], <http://www.chbr.noaa.gov/coastalresearch/prodino>, (verified 23 February 2002)
- Maryland Department of the Environment (MDE), Clean Water – Safe Shellfish, (2003), [Online], <http://www.mde.state.md.us/reference/factsheets/cleanwater.html>, (verified 16 January 2003)
- McClellan, P., Shanholtz, V.O., Petrauskas, B. and Kern, J. (2000) Bacterial source tracking: a tool for total maximum daily load development. In *Virginia Water Research Symposium 2000*, VWRRC Special Report SR-19-2000. eds. Younos, T. and Poff, J. pp. 157-172. Blacksburg, Va: VWRRC.
- McKenzie, M. C. (1998) Total maximum daily load: a major step to restoring our waters. *Small Flows* 12, 9-10.
- National Oceanic and Atmospheric Administration (NOAA), *The 1995 national shellfish register of classified growing waters*, (1997), Silver Spring, MD: NOAA, Office of Ocean Resources Conservation and Assessment (ORCA), Strategic Environmental Assessments (SEA) Division. 398 pp.
- NOAA. *1990 shellfish register of classified estuarine Waters: Data supplement*, (1992), Rockville, MD: NOAA, ORCA, SEA Division. 199 pp.
- NOAA, Case Studies, State of the Coastal Environment, Classified Shellfish Growing Waters, National Marine Fisheries Service, (2002), [Online], http://state-of-coast.noaa.gov/bulletins/html/sg_04/case.html, (verified 19 February 2002)
- NOAA, National Picture, State of the Coastal Environment, Classified Shellfish Growing Waters, (2002), [Online], http://state-of-coast.noaa.gov/bulletins/html/sg_04.national.html, (verified 16 January 2002)

- NOAA, Regional Contrasts, State of the Coastal Environment, Classified Shellfish Growing Waters, (2002), [Online], http://state-of-coast.noaa.gov/bulletins/html/sg_04/regional.html, (verified 16 January 2002)
- NOAA, State of the Coast Report, (2002), [Online], <http://state-of-coast.noaa.gov>, (verified 19 February 2002)
- Northern Neck, Virginia, History of the Neck, (1997-2000), [Online], <http://www.nothernneck.com/history.htm>, (verified 19 February 2002)
- Northumberland County Chamber of Commerce, Virginia, (2003), [Online], <http://www.riv.net/chamber>, (verified 12 March 2003)
- Oceana, American Oceans Campaign, (2002), [Online], <http://www.oceana.org>, (verified 21 March 2002)
- Patterson, S., Once a Bounty Now Wasted, (1997), Times Union Staff Writer, Florida Times
- Parveen, S., Murphree, R.L., Edmiston, L., Kaspar, C.W., Portier, K.M. and Tamplin, M.L. (1997) Association of multiple-antibiotic-resistance profiles with point and nonpoint sources of *Escherichia coli* in Apalachicola Bay. *Applied and Environmental Microbiology* 63, 2607-2612.
- Parveen, S., K., Portier, M., Robinson, K., Edmiston, L. and Tamplin, M.L. (1999) Discriminant analysis of ribotype profiles of *Escherichia coli* for differentiating human and nonhuman sources of fecal pollution. *Applied and Environmental Microbiology* 65, 3142-3147.
- Rippey, S.R., Infectious Diseases Associated with Molluscan Shellfish Consumption, (1994), *Clinical Microbiology Review* 7(4):419-425
- Rhodes, M. W. and Kator, H. (1999) Sorbitol-fermenting bifidiobacteria as indicators of diffuse human faecal pollution in estuarine watersheds. *Applied and Environmental Microbiology* 67, 528-535
- Roanoke Times, Associated Press, (February 2002), Acting State Health Commissioner Closes Virginia's Potomac River and Tributaries for Shellfish Harvesting
- Rountree, H., (1989), University of Oklahoma Press, pg. 148
- Scott, T. M., Rose, J.B., Jenkins, T. M., Farrah, S. R., Lukasik, J., (2002), Microbial Source Tracking: Current Methodology and Future Directions. *Applied and Environmental Microbiology*, 68, 5796-5803
- Shellfish Protection, United States Environmental Protection Agency, (2003), [Online], <http://www.epa.gov/waterscience/shellfish>, (verified 13 February 2002)
- Simmons, G. M., Jr., Herbein, S.A., and James, C.M. (1995) Managing nonpoint fecal coliform sources to tidal inlets. *Water Resource Update* 100, 64-74.
- Simmons, G. M., Jr. and Herbein, S.A. (1998) Potential Sources of *Escherichia coli* to a Children's Pool in La Jolla, CA. Final Report for the City of San Diego and the County of San Diego Department of Environmental Health, San Diego, CA.
- Simmons, G. M., Jr., Wayne, D.F., and Herbein, S.A., (2002), Estimating Nonpoint Source Fecal Coliform Sources Using DNA Profile Analysis, *Advance in Water Monitoring Research*, pg. 143-167

- Simpson, J. M., Santo-Domingo, J. W., Reasoner, D. J., (2002), Critical Review – Microbial Source Tracking: State of the Science, U.S. Environmental Protection Agency, *Environmental Science and Technology*, 36(24) 5279-5288
- Sinton, L. W., Donnison, A. M., and Hastie C. M., (1993) Faecal streptococci as faecal pollution indicators: a review. II, Sanitary significance, survival, and use. *N. Z. J. Marine Freshwater Research* 27:117-137.
- U.S. Environmental Protection Agency, (1986), *Ambient Water Quality Criteria for Bacteria-1986*. Report EPA440/5-84-002. U.S. Environmental Protection Agency, Washington, D.C.
- U.S. Environmental Protection Agency, (1997), Recreational Water Quality. [Online] http://www.epa.nsw.gov.au/soe/97/ch3/18_1.htm (verified 28 April 2002).
- U.S. Environmental Protection Agency, (1999a), Total Maximum Daily Load (TMDL) Program. [Online] <http://www.epa.gov/owow/wtr1/tmdl/tmdlfs.html> (verified 28 April 2002).
- U.S. Environmental Protection Agency, (1999b), National Watershed Database 305b. [Online] <http://www.epa.gov/surf3/states/VA> (verified 28 April 2002).
- U.S. Environmental Protection Agency, (2002), Press Release, Headquarters
- U.S. Environmental Protection Agency, (2002), Wastewater Technology Fact Sheet, Bacterial Source Tracking, EPA Fact Sheet, Office of Water, Municipal Technology Branch
- Virginia Department of Environmental Quality, (2001), Virginia 1998 303(d) Impaired Waters Map. [Online] <http://www.deq.statEnt.va.us/tmdl> (verified 28 April 2002).
- Virginia Department of Game and Inland Fisheries, VADGIF, (2002), [Online], Fisheries Program and Funding, http://www.dgif.state.va.us/fishing/fisheries_program_and_funding.html, (verified 22 January 2003)
- Whitlock, J. E., Jones, D. T., and Harwood, V. J., (2002), Identification of the sources of fecal coliforms in an urban watershed using antibiotic resistance analysis, *Water Research* 36:4273-4282.
- Wiggins, B.A., (1996) Discriminant analysis of antibiotic resistance patterns in fecal streptococci, a method to differentiate human and animal sources of fecal pollution in natural waters. *Applied and Environmental Microbiology* 62, 3997-4002.
- Wiggins, B.A., Andrews, R.W., Conway, R.A., Corr, C.L., Dobratz, ENT.J., Dougherty, D.P., Eppard, J.R., Knupp, S.R., Limjoco, M.C., Mettenburg, J.M., Rinehardt, J.M., Sonsino, J., Torrijos, O.L. and Zimmerman. (1999) Identification of sources of fecal pollution using discriminant analysis: Supporting evidence from large data sets. *Applied and Environmental Microbiology* 65, 3483-3486.

