



US009228242B2

(12) **United States Patent**  
**Meng et al.**

(10) **Patent No.:** **US 9,228,242 B2**  
(45) **Date of Patent:** **Jan. 5, 2016**

(54) **PORCINE TORQUE TENO VIRUS VACCINES AND DIAGNOSIS**

(75) Inventors: **Xiang-Jin Meng**, Blacksburg, VA (US);  
**Yaowei Huang**, Blacksburg, VA (US)

(73) Assignee: **Virginia Tech Intellectual Properties, Inc.**, Blacksburg, VA (US)

(\* ) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 596 days.

(21) Appl. No.: **12/861,378**

(22) Filed: **Aug. 23, 2010**

(65) **Prior Publication Data**

US 2011/0045019 A1 Feb. 24, 2011

**Related U.S. Application Data**

(60) Provisional application No. 61/235,833, filed on Aug. 21, 2009, provisional application No. 61/316,519, filed on Mar. 23, 2010.

(51) **Int. Cl.**

**C12N 15/33** (2006.01)  
**A61K 39/12** (2006.01)  
**C12N 15/70** (2006.01)  
**C12N 1/21** (2006.01)  
**C12N 7/00** (2006.01)  
**C12Q 1/70** (2006.01)  
**C07K 14/005** (2006.01)  
**C07K 16/08** (2006.01)  
**G01N 33/569** (2006.01)  
**A61K 39/00** (2006.01)

(52) **U.S. Cl.**

CPC ..... **C12Q 1/701** (2013.01); **A61K 39/12** (2013.01); **C07K 14/005** (2013.01); **C07K 16/081** (2013.01); **C12N 7/00** (2013.01); **G01N 33/56983** (2013.01); **A61K 2039/552** (2013.01); **C12N 2750/00021** (2013.01); **C12N 2750/00022** (2013.01); **C12N 2750/00034** (2013.01); **C12N 2750/14034** (2013.01); **G01N 2333/085** (2013.01)

(58) **Field of Classification Search**

CPC ..... **C12N 7/00**; **C12N 2750/00**; **C12N 2750/00021**; **C12N 2750/14034**; **C07K 14/005**  
See application file for complete search history.

(56) **References Cited**

**FOREIGN PATENT DOCUMENTS**

WO 2008127279 A2 10/2008  
WO 2008150275 A2 12/2008  
WO WO 2010/044889 A2 \* 4/2010

**OTHER PUBLICATIONS**

Y.W. Huang et al., *Virology*, 2010, 396:289-297, available online Nov. 13, 2009.\*

Huang YW, Ni YY, Dryman BA, and Meng XJ. ORF1 protein [Torque teno sus virus 1 b]. GenBank Acc. No. ADD46854. Updated Nov. 17, 2010.\*

Huang YW, Ni YY, Dryman BA, Meng XJ. Multiple infection of porcine Torque teno virus in a single pig and characterization of the full-length genomic sequences of four U.S. prototype PTTV strains: implication for genotyping of PTTV. *Virology*. Jan. 20, 2010;396(2):289-97. Epub Nov. 13, 2009.\*

Huang YW and Meng XJ. ORF1 protein [Torque teno sus virus 1 b]. GenBank: ADD46854.1. Dep. Nov. 17, 2010.\*

Anderson, et al., "Failure to genotype herpes simplex virus by real-time PCR assay and melting curve analysis due to sequence variation within probe binding sites". *Journal of Clinical Microbiology*, 2003, pp. 2135-2137 vol. 41, American Society for Microbiology.

Bao, et al., "Virus Classification by Pairwise Sequence Comparison (PASC)", 2008, pp. 342-348, vol. 5, Elsevier Ltd. Oxford, U.K.

Biagini, et al., "Classification of TTV and related viruses (anelloviruses)". *Current Topics in Microbiology Immunology*, 2009, pp. 21-33, vol. No. 331, Springer-Verlag Berlin Heidelberg.

Biagini, et al., "Distribution and genetic analysis of TTV and TTMV major phylogenetic groups in French blood donors". *Journal of Medical Virology*, 2006, pp. 298-304, vol. No. 78, Issue No. 2, Journal of Medical Virology, Marseille, France.

Biagini, et al., "Circular genomes related to anelloviruses identified in human and animal samples by using a combined rolling-circle amplification/sequence-independent single primer amplification approach". *Journal of General Virology*, 2007, pp. 2696-2701, vol. 88, Pt 10, Marseille, France.

Brassard, et al., "Development of a real-time TaqMan PCR assay for the detection of porcine and bovine Torque teno virus", *Journal of Applied Microbiology, Agriculture and Agri-food Canada*, Nov. 2009, pp. 2191-2198, Food Research and Development Centre, Saint-Hyacinthe, QC, Canada.

Davidson, et al., "Unraveling the puzzle of human anellovirus infections by comparison with avian infections with the chicken anemia virus", *Virus Research*, 2008, pp. 1-15, vol. 137, Issue 1, Israel.

(Continued)

*Primary Examiner* — Benjamin P Blumel

*Assistant Examiner* — Rachel Gill

(74) *Attorney, Agent, or Firm* — Blank Rome LLP

(57) **ABSTRACT**

The present invention provides four purified preparation containing a polynucleic acid molecule encoding porcine Torque teno virus (PTTV) genotypes or subtypes PTTV1a-VA, PTTV1b-VA, PTTV2b-VA, and PTTV2c-VA. The present invention also provides infectious DNA clones, biologically functional plasmid or viral vector containing the infectious nucleic acid genome molecule of the same. The present invention further provides live, attenuated, vector-expressed and purified recombinant capsid subunit or killed viral vaccines for protection against PTTV infection. The present invention additionally provides subunit vaccines comprising PTTV specific gene products, especially ORF1 capsid gene product for protection against PTTV infection. Further, the present invention provides methods for diagnosing PTTV infection via polymerase chain reaction (PCR) using specific primer for PTTV1, PTTV2, and individual PTTV1 genotypes. Finally, the present invention provides methods for diagnosing PTTV infection via immunological methods, e.g., enzyme-linked immunoabsorbent assay (ELISA) and Western blot using PTTV specific antigens for detecting serum PTTV specific antibodies.

**5 Claims, 31 Drawing Sheets**

(56)

## References Cited

## OTHER PUBLICATIONS

- De Smit, et al., "Apoptosis-inducing proteins in chicken anemia virus and TT virus". *Current Topics in Microbiology and Immunology*, 2009, pp. 131-149, vol. 331.
- Ellis, et al., "Effect of coinfection with genogroup 1 porcine torque teno virus on porcine circovirus type 2-associated postweaning multisystemic wasting syndrome in gnotobiotic pigs". *American Journal of Veterinary Research*, Dec. 2008, pp. 1608-1614, vol. 69, Issue 12, Schaumburg, IL.
- Gallei, et al., "Porcine Torque teno virus: Determination of viral genomic loads by genogroup-specific multiplex rt-PCR, detection of frequent multiple infections with genogroups 1 or 2, and establishment of viral full-length sequences". *Veterinary Microbiology*, 2010, pp. 202-212, vol. 143, Veterinary Microbiology, Munster, Germany.
- Gibellini, et al., "Simultaneous detection of HCV and HIV-1 by SYBR Green real time multiplex RT-PCR technique in plasma samples". *Molecular and Cellular Probes*, Mar. 2006, pp. 223-229, vol. 20.
- Hino, et al., "Torque teno virus (TTV): current status". *Reviews in Medical Virology*, 2007, pp. 45-57, vol. 17, Wiley Interscience.
- Hino, et al., "Relationship of Torque teno virus to chicken anemia virus". *Current Topics in Microbiology and Immunology*, 2009, pp. 117-130, vol. 331, Springer Verlag Berlin Heidelberg.
- Ilyina, et al., "Conserved sequence motifs in the initiator proteins for rolling circle DNA replication encoded by diverse replicons from eubacteria, eucaryotes and archaeobacteria". *Nucleic Acids Research*, pp. 3279-3285, vol. 20, No. 13, NIH, Bethesda, MD.
- Inami, et al., "Full-length nucleotide sequence of a simian TT virus isolate obtained from a chimpanzee: evidence for a new TT virus-like species". *Virology*, 2000, pp. 330-335, vol. 277, No. 2, Academic Press.
- Jelcic, et al., "Isolation of multiple TT virus genotypes from spleen biopsy tissue from a Hodgkin's disease patient: genome reorganization and diversity in the hypervariable region". *Journal of Virology*, 2004, pp. 7498-7507, vol. 78, No. 14, American Society for Microbiology.
- Kakkola, et al., "Replication of and protein synthesis by TT viruses". *Current Topics in Microbiology and Immunology*, 2009, pp. 53-64, vol. 331, Springer Verlag Berlin Heidelberg.
- Kekarainen, et al., "Detection of swine Torque teno virus genogroups 1 and 2 in boar sera and semen". *Theriogenology*, 2007, pp. 966-971, vol. 68, No. 7.
- Kekarainen, et al., "Prevalence of swine Torque teno virus in post-weaning multisystemic wasting syndrome (PMWS)-affected and non-PMWS-affected pigs in Spain". *Journal of General Virology*, 2006, pp. 833-837, vol. 87, Part 4, UK.
- Krakowka, et al., "Evaluation of the effects of porcine genogroup 1 torque teno virus in gnotobiotic swine". *American Journal of Veterinary Research*, 2008, pp. 1623-1629, vol. 69.
- Krakowka, et al., "Evaluation of induction of porcine dermatitis and nephropathy syndrome in gnotobiotic pigs with negative results for porcine circovirus type 2". *American Journal of Veterinary Research*, 2008, pp. 1615-1622, vol. 69, Part 12.
- Maggi, et al., "Immunobiology of the Torque teno viruses and other anelloviruses". *Current Topics in Microbiology and Immunology*, 2009, pp. 65-90, vol. 331.
- Martinez, "Simultaneous detection and genotyping of porcine reproductive and respiratory syndrome virus (PRRSV) by real-time RT-PCR and amplicon melting curve analysis using SYBR Green". *Research in Veterinary Science*, 2008, pp. 184-193 vol. 85, Issue 1.
- McKeown, et al., "Molecular characterization of porcine TT virus, an orphan virus, in pigs from six different countries". *Veterinary Microbiology*, 2004, pp. 113-117, vol. 104, Issues 1-2.
- Mouillesseaux, et al., "Improvement in the specificity and sensitivity of detection for the Taura syndrome virus and yellow head virus of penaeid shrimp by increasing the amplicon size in SYBR Green real-time RT-PCR". *Journal of Virological Methods*, 2003, pp. 121-127, vol. 111, Issue 2.
- Mueller, et al., "Gene expression of the human Torque Teno Virus isolate P/1C1". *Virology*, 2008, pp. 36-45, vol. 381, Issue 1.
- Ng, et al., "Novel anellovirus discovered from a mortality event of captive California sea lions". *Journal of General Virology*, 2009, pp. 1256-1261, vol. 90, Pt 5.
- Niel, et al., "Rolling-circle amplification of Torque teno virus (TTV) complete genomes from human and swine sera and identification of a novel swine TTV genogroup". *Journal of General Virology*, 2005, pp. 1343-1347, vol. 86, Pt. 5.
- Niel, et al., "Coinfection with multiple TT virus strains belonging to different genotypes is a common event in healthy Brazilian adults". *Journal of Clinical Microbiology*, 2000, pp. 1926-1930, vol. 38, No. 5.
- Ninomiya, et al., "Analysis of the entire genomes of torque teno midi virus variants in chimpanzees: infrequent crossspecies infection between humans and chimpanzees". *Journal of General Virology*, 2009, pp. 347-358, vol. 90, Pt 2.
- Nishizawa, et al., "A novel DNA virus (TTV) associated with elevated transaminase levels in posttransfusion hepatitis of unknown etiology". *Biochemical Biophysical Research Communications*, 1997, pp. 92-97, vol. 241, No. 1.
- Okamoto, et al., "History of discoveries and pathogenicity of TT viruses". *Current Topics in Microbiology and Immunology*, 2009, pp. 1-20, vol. 331.
- Okamoto, et al., "TT viruses in animals". *Current Topics in Microbiology and Immunology*, 2009, pp. 35-52, vol. 331.
- Okamoto, et al., "Genomic and evolutionary characterization of TT virus (TTV) in tupaia and comparison with species-specific TTVs in humans and non-human primates". *Journal of General Virology*, 2001, pp. 2041-2050, vol. 82, Pt 9.
- Okamoto, et al., "Species-specific TT viruses in humans and nonhuman primates and their phylogenetic relatedness". *Virology*, 2000, pp. 368-378, vol. 277, No. 2.
- Okamoto, et al., "TT virus mRNAs detected in the bone marrow cells from an infected individual". *Biochemical and Biophysical Research Communications*, 2000, pp. 700-707, vol. 279, No. 2.
- Okamoto, et al., "Genomic characterization of TT viruses (TTVs) in pigs, cats and dogs and their relatedness with species-specific TTVs in primates and tupaia". *Journal of General Virology*, 2002, pp. 700-707, vol. 83, Pt 6.
- Opriessnig, et al., "Porcine circovirus type 2 associated disease: update on current terminology, clinical manifestations, pathogenesis, diagnosis, and intervention strategies". *Journal of Veterinary Diagnostic Investigation*, 2007, pp. 591-615, vol. 19.
- Pal, et al., "Development and validation of a duplex real-time PCR assay for the simultaneous detection and quantification of porcine circovirus type 2 and an internal control on porcine semen samples". *Journal of Virological Methods*, 2008, pp. 217-225, vol. 149.
- Peters, et al., "Attenuation of chicken anemia virus by site-directed mutagenesis of VP2". *Journal of General Virology*, 2007, pp. 2168-2175, vol. 88, Pt. 8.
- Peters, et al., "Site-directed mutagenesis of the VP2 gene of Chicken anemia virus affects virus replication, cytopathology and host-cell MHC class I expression". *Journal of General Virology*, 2006, pp. 823-831, vol. 87, Pt. 4.
- Peters, et al., "Chicken anemia virus VP2 is a novel dual specificity protein phosphatase". *Journal of Biological Chemistry*, 2002, pp. 39566-39573, vol. 277, No. 42.
- Pozzuto, et al., "In utero transmission of porcine torque teno viruses". *Veterinary Microbiology*, 2009, pp. 375-379, vol. 137.
- Prasetyo, et al., "Replication of chicken anemia virus (CAV) requires apoptin and is complemented by VP3 of human torque teno virus (TTV)". *Virology*, 2009, pp. 85-92, vol. 385, No. 1.
- Qiu, et al., "Human circovirus TT virus genotype 6 expresses six proteins following transfection of a full-length clone". *Journal of Virology*, 2005, pp. 6505-6510, vol. 79, No. 10.
- Ririe, et al., "Product differentiation by analysis of DNA melting curves during the polymerase chain reaction". *Analytical Biochemistry*, 1997, pp. 154-160, vol. 245.
- Sibila, et al., "Swine torque teno virus (TTV) infection and excretion dynamics in conventional pig farms". *Veterinary Microbiology*, 2009, pp. 213-228, vol. 139.
- Takayama, et al., "Prevalence and persistence of a novel DNA TT virus (TTV) infection in Japanese haemophiliacs". *British Journal of Haematology*, 1999, vol. 104, No. 3, pp. 626-629.

(56)

**References Cited**

OTHER PUBLICATIONS

Wilhelm, et al., "Real-time PCR protocol for the detection of porcine parvovirus in field samples". *Journal of Virological Methods*, 2006, pp. 257-260, vol. 134.

Okamoto, et al. "Genomic characterization of TT viruses (TTVs) in pigs, cats and dogs and their relatedness with species-specific TTVs in primates and tupias," *Journal of General Virology*, 2002, pp. 1291-1297, The Society for General Microbiology, Reading, UK.

Y.W. Huang, et al., Multiple infection of porcine Torque teno virus in a single pig and characterization of the full-length genomic

sequences of four U.S. prototype PTTV strains: Implication for genotyping of PTTV, Nov. 2009, p. 289-297, *Virology*, vol. 396.

Genbank; GU456383.1.

Genbank; GU456384.1.

Genbank; GU456385.1.

Genbank; GU456386.1.

Okamoto, H. et al., "Genomic characterization of TT viruses (TTVs) in pigs, cats and dogs and their relatedness with species-specific TTVs in primates and tupaias"; website: <http://vir.sgmjournals.org/content/83/6/1291.full.pdf+html>; *Journal of General Virology*; vol. 83; *Paginas*; Jun. 30, 2002; pp. 1291-1297.

\* cited by examiner




FIG. 1A




FIG. 1B




FIG. 2




FIG. 3A




FIG. 3B




FIG. 3C




FIG. 3D




FIG. 3E




FIG. 4A

	210	220	230	240	250	260	270	280	290	300
PTTV1a-VA	VMLQAKNSRLIVCKQKMP	IRRRVK	--SIFIPPPAQLTTQWKFOQELCQFPLFNWACICIDMDTPFDYNG	--ZWRNANWLMRRLQNG	-----	NMEYIERW	286			
Sd-ITV31	VMLQAKNSRLIVCKQKMP	IRRRVK	--SIFIPPPAQLTTQWKFOQELCQFPLFNWACICIDMDTPFDYNG	--ZWRNANWLMRRLQNG	-----	NMEYIERW	286			
PTTV1b-VA	VMMAKRTRLVIARDRAPHRRVR	--KIFIPPPSRDITIQFQTD	FCRRLFTWAAGLIDMOKPFDANG	--AFRNAWMLETRNDQG	-----	EMKYIELW	286			
TTV-1p	VMMAKRTRLVIARERAPHRRKVR	--KIFIPPPSRDITIQFQTD	FCRRLFTWAAGLIDMOKPFDANG	--AFRNAWMLEQRNDQG	-----	EMKYIELW	284			
TTV-2p	IMLLKQHKIVLSQONCNPNRQKPV	T--KFKPPKLT	SQWRLSRELAKMPLIRLGV	SFIDLTEPWE	CGNAFYSLGYEAVK	DQGHMSNW	QIKYWIY	263		
PTV-2b-VA	IMLLKQHKIVLSQKDCNPNRQKPV	T--KFRPPKLT	SQWRLSRELSKIPIRLGIS	LIDLSEPWLECGNAFYSLGYE	ASKHGRWSNW	QMKYFWIY	263			
PTV-2c-VA	IMLLKQHKIVLSQKDCNPNRQKPV	T--KIRPPKLT	SQWRLSRELAKMPLVRLGV	SLIDLSEPWLECGNAFYSLGYE	ASKHGRWSNW	QIKYFWIY	263			
	310	320	330	340	PTTV1-HVR2	360	370	380	390	400
PTTV1a-VA	GRIPMTGDTE	_PPADDFKAGGVNKNFK	P	TG--IQRIY	P	AVAVC	_VEGNKR	VVKWATVHNGP	L	P
Sd-ITV31	GRIPMTGDTE	_PPADDFKAGGVNKNFK	P	TG--IQRIY	P	AVAVC	_VEGNKR	VVKWATVHNGP	L	P
PTTV1b-VA	GRVPPQGDTE	_PKQSEFKKGDNNPNY	IT	GHEKN	I	Y	P	E	L	_YVDQ
TTV-1p	GRVPPQGDSE	_PKKKEF	STGTDNPNY	VQDNEEK	N	I	Y	P	E	_YVDQ
TTV-2p	DTGVGNNAVYVILLK	KDVIDNPGNMAITFKAS	--GGQHPDA	_DHIELINQ	GPW	L	Y	F	G	K
PTTV2b-VA	DTGVGNNAVYVILLK	KDVS	DNPGDMATQ	F	Y	T	G--SGQHPDA	_DHIEMVNEG	P	W
PTTV2c-VA	DTGVGNNAVYVILLK	QEVDDNPGAMATK	F	Y	T	G--PGQHPDA	_DRIEQINEG	P	W	L

FIG. 4B




FIG. 4C

PTV1a-VA	MKEKYWEEAWLTSCTSIHDDHHCNCGSWRDHLWTICALDDADLAAAAD_IIEEEADGGEDFGFVDG--DPGDAGG	73
Sd-TTV31	MKEKYWEEAWLTSCTSIHDDHHCNCGSWRDHLWTICALDDADLAAAAD_IIEEEADGGEDFGFVDG--DPGDAGG	73
PTV1b-VA	MP--EHWEEAWLEATKGWHDLDRCRCGNWQDHLWLLADGDAALAAAVDAIER-DAMGGEDVTTATDRVTIGDDGW	72
TTV-1p	MP--EHWEEAWLEATKGWHDLDRCRCGNWQDHLWLLADGDAALAAAVDAIER-DAMAGDDATTATGRVTIGDDGW	72
TTV-2p	M-----EERWLTVA <sup>^</sup> YCAHGLFC <sup>^</sup> CKDPKKHLEKCLTD---AIADAEEDRHGDCGTGGGDA <sup>^</sup> TFDIGIDALLAAAAQR	68
PTV1b-VA	M-----EERWLTVA <sup>^</sup> YCAHGLFC <sup>^</sup> CKNPKKHLEKCLTD---AIADAEGRHGDG <sup>^</sup> GTGGGDA <sup>^</sup> TFDIGIDALLAAAAQR	68
PTV1c-VA	M-----EERWLTVA <sup>^</sup> YCAHGLFC <sup>^</sup> GCKDPKKHLEKCLTD---AIADAEGRHGDG <sup>^</sup> GTGGGDA <sup>^</sup> SFDIGIDALLAAAAQR	68

FIG. 5




FIG. 6A




FIG. 6B




FIG. 7A




FIG. 7B




FIG. 7C




FIG. 7D




FIG. 7E




FIG. 8




FIG. 9A

DOMAIN I 322 349  
 | SEQDIKKLAHDQXIAREYARDPKSKKLIK  
 CONSENSUS  
 PTTV2c-VA E.....NN.....  
 PTTV2b-VA .....D.V.....  
 TTV2#472142 .....D.....  
 TTV-2p .....E.SAE.S.T.....

DOMAIN II 536 625  
 | YDCDGIVRTDILKRLLLELPTETEETEKAYPLLGQKTEKEPLSDSDEESVISSTSSGSSQEEFTQRRRXHKPSKRRLLKHLQRVVVKRMKTL  
 CONSENSUS  
 PTTV2c-VA .....O.....  
 PTTV2b-VA .....KH.....  
 TTV2#472142 .....Q.....  
 TTV-2p .....E.....H.....

FIG. 9B




FIG. 10A




FIG. 10B




FIG. 10C




FIG. 11A




FIG. 11B




FIG. 11C




FIG. 12




FIG. 13




FIG. 14A




FIG. 14B




FIG. 15A




FIG. 15B





FIG. 15C



CONVENTIONAL PIGS FROM  
WISCONSIN

FIG. 16




FIG. 17A




FIG. 17B




FIG. 17C




FIG. 17D




FIG. 17E




FIG. 17F




FIG. 18A




FIG. 18B




FIG. 19A




FIG. 19B




FIG. 20A




FIG. 20B




FIG. 21






FIG. 22A






FIG. 22B




FIG. 23A




FIG. 23B

1

## PORCINE TORQUE TENO VIRUS VACCINES AND DIAGNOSIS

### REFERENCE TO RELATED APPLICATION

This patent application claims the benefit of U.S. Provisional Patent Application No. 61/235,833, filed on Aug. 21, 2009, and U.S. Provisional Patent Application 61/316,519, filed on Mar. 23, 2010, whose disclosures are hereby incorporated by reference in their entirety into the present disclosure.

### FIELD OF INVENTION

The present invention relates to vaccines for protecting against porcine Torque teno virus (TTV) infection, and infectious DNA clones of porcine TTV (PTTV) and their uses thereof. The present invention also relates to diagnosis of porcine Torque teno virus (PTTV) infection, particularly diagnosis of species- or type-specific PTTV infection, and simultaneous infection of multiple strains from different genotypes.

### BACKGROUND OF THE INVENTION

Torque teno virus (TTV) was first discovered in a Japanese patient with post-transfusion non-A-E hepatitis in 1997 (Nishizawa, T., Okamoto, H., Konishi, K., Yoshizawa, H., Miyakawa, Y., and Mayumi, M. (1997). A novel DNA virus (TTV) associated with elevated transaminase levels in post-transfusion hepatitis of unknown etiology. *Biochem Biophys Res Commun* 241(1), 92-7.). Since then, a large number of human TTV strains and two groups of TTV-related viruses, designated subsequently as Torque teno mini virus (TTMV) and Torque teno midi virus (TTMDV), have been identified with high prevalence in serum and other tissues from healthy humans (Hino, S., and Miyata, H. (2007). Torque teno virus (TTV): current status. *Rev Med Virol* 17(1), 45-57; Okamoto, H. (2009a). History of discoveries and pathogenicity of TT viruses. *Curr Top Microbiol Immunol* 331, 1-20). Human TTV, TTMV and TTMDV are non-enveloped spherical viruses with circular single-stranded DNA (ssDNA) genomes of 3.6-3.9, 2.8-2.9 and 3.2 kb in length, respectively, and they are currently classified into a newly-established family Anelloviridae by the International Committee on Taxonomy of Viruses (ICTV (Biagini, P. (2009). Classification of TTV and related viruses (anelloviruses). *Curr Top Microbiol Immunol* 331, 21-33). These three groups of TTV-related viruses exhibit a high degree of genetic heterogeneity, each consisting of many genogroups and genotypes (Biagini, P., Gallian, P., Cantaloube, J. F., Attoui, H., de Micco, P., and de Lamballerie, X. (2006). Distribution and genetic analysis of TTV and TTMV major phylogenetic groups in French blood donors. *J Med Virol* 78(2), 298-304; Jelcic, I., Hotz-Wagenblatt, A., Hunziker, A., Zur Hausen, H., and de Villiers, E. M. (2004). Isolation of multiple TT virus genotypes from spleen biopsy tissue from a Hodgkin's disease patient: genome reorganization and diversity in the hypervariable region. *J Virol* 78(14), 7498-507). The prevalence of multiple infections of TTV with different genotypes as well as dual or triple infections of TTV, TTMV and TTMDV have been documented in humans, and are considered to be a common event in healthy human adults (Niel, C., Saback, F. L., and Lampe, E. (2000). Coinfection with multiple TT virus strains belonging to different genotypes is a common event in healthy Brazilian adults. *J Clin Microbiol* 38(5), 1926-30; Ninomiya, M., Takahashi, M., Hoshino, Y., Ichiyama, K., Simmonds, P., and Okamoto,

2

H. (2009). Analysis of the entire genomes of torque teno midi virus variants in chimpanzees: infrequent cross-species infection between humans and chimpanzees. *J Gen Virol* 90(Pt 2), 347-58; Okamoto, H. (2009a). History of discoveries and pathogenicity of TT viruses. *Curr Top Microbiol Immunol* 331, 1-20; Takayama, S., Miura, T., Matsuo, S., Taki, M., and Sugii, S. (1999). Prevalence and persistence of a novel DNA TT virus (TTV) infection in Japanese haemophiliacs. *Br J Haematol* 104(3), 626-9).

TTV infects not only humans but also various other animal species as well including non-human primates, tupaia, pigs, cattle, cats, dogs and sea lions (Biagini, P., Uch, R., Belhouchet, M., Attoui, H., Cantaloube, J. F., Brisbane, N., and de Micco, P. (2007). Circular genomes related to anelloviruses identified in human and animal samples by using a combined rolling-circle amplification/sequence-independent single primer amplification approach. *J Gen Virol* 88(Pt 10), 2696-701; Inami, T., Obara, T., Moriyama, M., Arakawa, Y., and Abe, K. (2000). Full-length nucleotide sequence of a simian TT virus isolate obtained from a chimpanzee: evidence for a new TT virus-like species. *Virology* 277(2), 330-5; Ng, T. F., Suedmeyer, W. K., Wheeler, E., Gulland, F., and Breitbart, M. (2009). Novel anellovirus discovered from a mortality event of captive California sea lions. *J Gen Virol* 90(Pt 5), 1256-61; Okamoto, H. (2009b). TT viruses in animals. *Curr Top Microbiol Immunol* 331, 35-52; Okamoto, H., Nishizawa, T., Takahashi, M., Tawara, A., Peng, Y., Kishimoto, J., and Wang, Y. (2001). Genomic and evolutionary characterization of TT virus (TTV) in tupaia and comparison with species-specific TTVs in humans and non-human primates. *J Gen Virol* 82(Pt 9), 2041-50; Okamoto, H., Nishizawa, T., Tawara, A., Peng, Y., Takahashi, M., Kishimoto, J., Tanaka, T., Miyakawa, Y., and Mayumi, M. (2000a). Species-specific TT viruses in humans and nonhuman primates and their phylogenetic relatedness. *Virology* 277(2), 368-78; Okamoto, H., Takahashi, M., Nishizawa, T., Tawara, A., Fukai, K., Muramatsu, U., Naito, Y., and Yoshikawa, A. (2002). Genomic characterization of TT viruses (TTVs) in pigs, cats and dogs and their relatedness with species-specific TTVs in primates' and tupaia. *J Gen Virol* 83(Pt 6), 1291-7). In addition, chimpanzees are also infected with TTMV and TTMDV (Ninomiya, M., Takahashi, M., Hoshino, Y., Ichiyama, K., Simmonds, P., and Okamoto, H. (2009). Analysis of the entire genomes of torque teno midi virus variants in chimpanzees: infrequent cross-species infection between humans and chimpanzees. *J Gen Virol* 90(Pt 2), 347-58; Okamoto et al., 2000a, supra). Although the genomic sizes of the identified animal TTV strains, especially non-primate animal TTV, are relatively smaller than that of human TTV, they share the same genomic structure with a minimum of two partially overlapping open reading frames (ORF1 and ORF2) translated from the negative ssDNA as well as a short stretch of untranslated region (UTR) with high GC content (~90%) (Okamoto, 2009b, supra). The arrangement of TTV ORFs is quite similar to that of chicken anemia virus (CAV) belonging to the genus Gyrovirus in the family Circoviridae but is different from porcine circovirus (PCV) types 1 (PCV1) and 2 (PCV2), which are also classified into the same family (Davidson, I., and Shulman, L. M. (2008). Unraveling the puzzle of human anellovirus infections by comparison with avian infections with the chicken anemia virus. *Virus Res* 137(1), 1-15; Hino, S., and Prasetyo, A. A. (2009). Relationship of Torque teno virus to chicken anemia virus. *Curr Top Microbiol Immunol* 331, 117-30). The genomes of PCV1 and PCV2 are ambisense, in which the ORF1 is coded for by the genomic strand and the ORF2 is coded for by the antigenomic strand (Hino and Miyata, 2007, supra). Recently, the tran-

scription pattern and translated products of both human TTV genotypes 1 and 6 have been identified by transfection of the respective TTV infectious DNA clones into cultured cells (Mueller, B., Maerz, A., Doberstein, K., Finsterbusch, T., and Mankertz, A. (2008). Gene expression of the human Torque Teno Virus isolate P/1C1. *Virology* 381(1), 36-45; Qiu, J., Kakkola, L., Cheng, F., Ye, C., Soderlund-Venermo, M., Hedman, K., and Pintel, D. J. (2005). Human circovirus TT virus genotype 6 expresses six proteins following transfection of a full-length clone. *J Virol* 79(10), 6505-10. Expression of at least six proteins, designated ORF1, ORF2, ORF1/I, ORF2/2, ORF1/2 and ORF2/3, from three or more spliced mRNAs, have been reported (Kakkola, L., Hedman, K., Qiu, J., Pintel, D., and Soderlund-Venermo, M. (2009). Replication of and protein synthesis by TT viruses. *Curr Top Microbiol Immunol* 331, 53-64; Mueller et al., 2008, supra; Qiu et al., 2005, supra). Accordingly, it is likely that, when more data regarding the animal TTV become available, the presumed genome structure of animal TTV will need to be modified.

Although TTV was first identified in a cryptogenic hepatitis patient, subsequent studies were not able to produce evidence of a significant role of TTV in the pathogenesis of hepatitis or other diseases (Hino and Miyata, 2007, supra; Maggi, F., and Bendinelli, M. (2009). Immunobiology of the Torque teno viruses and other anelloviruses. *Curr Top Microbiol Immunol* 331, 65-90; Okamoto, 2009a, supra). While human TTV is not considered to be directly associated with a disease, porcine TTV (PTTV) was recently shown to partially contribute to the experimental induction of porcine dermatitis and nephropathy syndrome (PDNS) combined with porcine reproductive and respiratory syndrome virus (PRRSV) infection (Krakowka, S., Hartunian, C., Hamberg, A., Shoup, D., Rings, M., Zhang, Y., Allan, G., and Ellis, J. A. (2008). Evaluation of induction of porcine dermatitis and nephropathy syndrome in gnotobiotic pigs with negative results for porcine circovirus type 2. *Am J Vet Res* 69(12), 1615-22), and also to the experimental induction of postweaning multisystemic wasting syndrome (PMWS) combined with PCV2 infection in a gnotobiotic pig model (Ellis, J. A., Allan, G., and Krakowka, S. (2008). Effect of coinfection with genogroup 1 porcine torque teno virus on porcine circovirus type 2-associated postweaning multisystemic wasting syndrome in gnotobiotic pigs. *Am J Vet Res* 69(12), 1608-14). The data suggested that porcine TTV is pathogenic in pigs. However, further in-depth studies with a biologically pure form of PTTV virus to definitively characterize the diseases and lesions associated with PTTV infection are needed.

Compared to human TTV, the genomic information of PTTV is very limited. Currently, only one full-length and two near full-length genomic sequences of PTTV are reported from pigs in Japan and Brazil, respectively (Niel, C., Diniz-Mendes, L., and Devalle, S. (2005). Rolling-circle amplification of Torque teno virus (TTV) complete genomes from human and swine sera and identification of a novel swine TTV genogroup. *J Gen Virol* 86(Pt 5), 1343-7; Okamoto et al., 2002, supra). Among the three known PTTV strains, the Sd-TTV31 and TTV-1p stains were clustered together into the genogroup 1 (PTTV1), whereas TTV-2p was the sole strain classified into the genogroup 2 (PTTV2) (Niel et al., 2005, supra). However, genogroup classification is a vague concept in the taxonomy of virology, and further and more accurate classification of PTTV is needed but can only be performed when more full-length genomic sequences of new PTTV strains representing multiple genotypes become available.

It was previously showed that PTTV infections were widespread in pigs from six different countries including the

United States, Canada, Spain, China, Korea and Thailand (McKeown, N. E., Fenaux, M., Halbur, P. G., and Meng, X. J. (2004). Molecular characterization of porcine TT virus, an orphan virus, in pigs from six different countries. *Vet Microbiol* 104(1-2), 113-7).

Whether porcine TTVs play a significant role in pathogenesis of specific swine diseases is still debatable. In a gnotobiotic pig model, it was shown that PTTV1 infection alone did not develop any clinical diseases but induced mild histological lesions (Krakowka, S. and Ellis, J. A., 2008. Evaluation of the effects of porcine genogroup 1 torque teno virus in gnotobiotic swine. *Am J Vet Res* 69, 1623-9). Gnotobiotic pigs that were experimentally inoculated with both PTTV1 and porcine reproductive and respiratory syndrome virus (PRRSV) developed clinical porcine dermatitis and nephropathy syndrome (PDNS) (Krakowka, S., Hartunian, C., Hamberg, A., Shoup, D., Rings, M., Zhang, Y., Allan, G. and Ellis, J. A., 2008. Evaluation of induction of porcine dermatitis and nephropathy syndrome in gnotobiotic pigs with negative results for porcine circovirus type 2. *Am J Vet Res* 69, 1615-22), whereas pigs inoculated with both PTTV1 and porcine circovirus type 2 (PCV2) developed acute postweaning multisystemic wasting syndrome (PMWS) (Ellis et al., 2008, supra). Although PCV2 is considered as the primary causative agent for clinical PMWS or PCV-associated diseases (PCVAD), a higher prevalence of PTTV2 infection in PMWS-affected pigs with low or no PCV2 than that in non-PMWS-affected pigs was observed in Spain (Kekarainen et al., 2006, supra). The data collectively suggest that porcine TTVs may serve as co-factors involved in triggering or exacerbating diseases in pigs.

Porcine TTV has been detected in porcine serum, fecal, saliva, semen and tissue samples of infected pigs, indicating its diverse transmission routes including both horizontal and vertical transmissions (Kekarainen et al., 2007, supra; Pozzuto, T., Mueller, B., Meehan, B., Ringler, S. S., McIntosh, K. A., Ellis, J. A., Mankertz, A. and Krakowka, S., 2009. In utero transmission of porcine torque teno viruses. *Vet Microbiol* 137, 375-9; Sibila, M., Martinez-Guino, L., Huerta, E., Llorens, A., Mora, M., Grau-Roma, L., Kekarainen, T. and Segales, J., 2009. Swine torque teno virus (TTV) infection and excretion dynamics in conventional pig farms. *Vet Microbiol* 139, 213-8). However, current detection of porcine TTV infection was mainly based upon conventional PCR assays. Thus far, neither serological assay nor viral culture system has been established. In particular, nested PCR amplifications of the conserved regions in the UTR, of PTTV1 and PTTV2, respectively, developed by a Spanish group, have become widely used (Kekarainen et al., 2006, supra). Since the amount of virus is likely associated with the severity of clinical diseases, as demonstrated for PCV2-induced PCVAD (Opriessnig, T., Meng, X. J. and Halbur, P. G., 2007. Porcine circovirus type 2 associated disease: update on current terminology, clinical manifestations, pathogenesis, diagnosis, and intervention strategies. *J Vet Diagn Invest* 19, 591-615), it will be important to determine the viral load of porcine TTV by quantitative real-time PCR than the presence of TTV DNA by conventional PCR. In addition, real-time PCR is more reliable, rapid and less expensive than conventional PCR. Recently, two TaqMan probe-based real-time PCR assays were described for detection and quantification of two porcine TTV species (Brassard, J., Gagne, M. J., Houde, A., Poitras, E. and Ward, P., 2009. Development of a real-time TaqMan PCR assay for the detection of porcine and bovine Torque teno virus. *J Appl Microbiol*. Nov. 14, 2009, Epub ahead of print; Gallei, A., Pesch, S., Eskiing, W. S., Keller, C. and Ohlinger, V. F., 2009. Porcine Torque teno virus: Deter-

mination of viral genomic loads by genogroup-specific multiplex rt-PCR, detection of frequent multiple infections with genogroups 1 or 2, and establishment of viral full-length sequences. *Vet Microbiol.* Dec. 21, 2009, Epub ahead of print). A main drawback of probe-based assays is that the false-negative results may be obtained if the probe-binding sequences contain mutations (Anderson, T. P., Werno, A. M., Beynon, K. A. and Murdoch, D. R., 2003. Failure to genotype herpes simplex virus by real-time PCR assay and melting curve analysis due to sequence variation within probe binding sites. *J Clin Microbiol* 41, 2135-7). Considering the high degree of heterogeneity among the sequences of known porcine TTV strains, variations in the probe-binding sequences are expected for field strains of PTTVs. The SYBR green-based real-time PCR is an alternative method avoiding this potential problem, in spite of its relatively lower specificity, which provides a universal way to detect and quantify the potential porcine TTV variants. Moreover, melting curve analysis (MCA) following SYBR green real-time PCR ensures reaction specificity and also allows multiplex detection of distinct types of virus (Ririe, K. M., Rasmussen, R. P. and Wittwer, C. T., 1997. Product differentiation by analysis of DNA melting curves during the polymerase chain reaction. *Anal Biochem* 245, 154-60). MCA-based SYBR green real-time PCR methods have been successfully applied to various human and veterinary viruses (Gibellini, D., Gardini, F., Vitone, F., Schiavone, P., Furlini, G. and Re, M. C., 2006. Simultaneous detection of HCV and HIV-1 by SYBR Green real time multiplex RT-PCR technique in plasma samples. *Mol Cell Probes* 20, 223-9; Martinez, E., Rieira, P., Sitja, M., Fang, Y., Oliveira, S., and Maldonado, J., 2008. Simultaneous detection and genotyping of porcine reproductive and respiratory syndrome virus (PRRSV) by real-time RT-PCR and amplicon melting curve analysis using SYBR Green. *Res Vet Sci* 85, 184-93; Mouillesseaux, K. P., Klimpel, K. R. and Dhar, A. K., 2003. Improvement in the specificity and sensitivity of detection for the Taura syndrome virus and yellow head virus of penaeid shrimp by increasing the amplicon size in SYBR Green real-time RT-PCR. *J Virol Methods* 111, 121-7; Wilhelm, S., Zimmermann, P., Selbitz, H. J. and Truyen, U., 2006. Real-time PCR protocol for the detection of porcine parvovirus in field samples. *J Virol Methods* 134, 257-60).

Currently, little is known about PTTV-specific humoral response. Since PCR-based assays do not reflect the course of PTTV infection in pigs, an efficient enzyme-linked immunosorbent assay (ELISA) for detection of PTTV serum antibody is necessary to evaluate seroprevalence of PTTV and help characterize the role of PTTV in porcine diseases.

Thus far, no subunit, killed and live vaccines for porcine TTVs are available. It will be desirable and advantageous to express recombinant PTTV capsid proteins from different genotypes for development of PTTV subunit vaccines, and to construct infectious PTTV molecular DNA clones from different genotypes for propagating biological pure form of PTTVs in cell culture system that are used for killed and live vaccines development.

#### SUMMARY OF THE INVENTION

The present invention provides an infectious nucleic acid molecule ("infectious DNA clone") of porcine Torque teno virus (PTTV) comprising a nucleic acid molecule encoding an infectious PTTV which contains at least one copy of genomic sequence having at least 80% homology to a

genomic sequence selected from the group consisting of genotypes of PTTV1a-VA, PTTV1b-VA, PTTV2b-VA, and PTTV2c-VA.

According to one aspect of the present invention, the infectious DNA clones of PTTV of set forth in claim 1, wherein the genomic sequence is selected from sequences set forth in SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, and SEQ ID NO:12.

The present invention provides a biologically functional plasmid or viral vector containing the infectious PTTV genomes.

The present invention provides a suitable host cell transfected with the infectious clone DNA plasmid or viral vector.

The present invention provides an infectious PTTV produced by cells transfected with the PTTV infectious DNA clones.

The present invention also provides a viral vaccine comprising a nontoxic, physiologically acceptable carrier and an immunogenic amount of a member selected from the group consisting of (a) a nucleic acid molecule containing at least one copy of genomic sequence having at least 80% homology to a genomic sequence selected from the group consisting of PTTV genotypes or subtypes PTTV1a-VA, PTTV1b-VA, PTTV2b-VA, and PTTV2c-VA, or its complementary strand, (b) a biologically functional plasmid or viral vector containing a nucleic acid molecule containing at least one copy of genomic sequence having at least 80% homology to a genomic sequence selected from the group consisting of PTTV genotypes or subtypes PTTV1a-VA, PTTV1b-VA, PTTV2b-VA, and PTTV2c-VA, or its complementary strand, and (c) an avirulent, infectious nonpathogenic PTTV which contains at least one copy of genomic sequence having at least 80% homology to a genomic sequence selected from the group consisting of PTTV genotypes or subtypes PTTV1a-VA, PTTV1b-VA, PTTV2b-VA, and PTTV2c-VA.

According to one aspect of the present invention, the vaccine contains live PTTV virus derived from the PTTV infectious clones. According to another aspect of the present invention, the vaccine contains killed PTTV virus derived from the PTTV infectious clones.

The present invention provides purified recombinant proteins expressed from the ORF1 capsid genes of PTTV genotypes or subtypes PTTV1a-VA, PTTV1b-VA, and PTTV2c-VA in bacterial expression system, and the use of these recombinant capsid proteins as subunit vaccines against PTTV infections. In one embodiment of the present invention, the recombinant capsid proteins for the use as subunit vaccines are expressed in baculovirus expression system and other expression vector systems.

According to a further aspect of the present invention, further contains an adjuvant.

The present invention further provides a method of immunizing a pig against PTTV viral infection, comprising administering to a pig an immunologically effective amount of the viral vaccine.

According to one aspect of the present invention, the method comprising administering the recombinant subunit capsid protein, the infectious nucleic acid molecule or live PTTV virus to the pig.

According to another aspect of the present invention, the method comprising administering the vaccine parenterally, intranasally, intradermally, or transdermally to the pig. According a further aspect of the present invention, the method comprising administering the vaccine intralymphoidly or intramuscularly to the pig.

The present invention also provides an isolated polynucleotide consisting of the sequence of the nucleotide sequence of PTTV1a-VA set forth in SEQ ID NO:9.

The present invention also provides an isolated polynucleotide consisting of the sequence of the nucleotide sequence of PTTV1b-VA set forth in SEQ ID No:10.

The present invention also provides an isolated polynucleotide consisting of the sequence of the nucleotide sequence of PTTV2b-VA set forth in SEQ ID No:11.

The present invention also provides an isolated polynucleotide consisting of the sequence of the nucleotide sequence of PTTV2c-VA set forth in SEQ ID No:12.

The present invention further provides a subunit vaccine comprising an immunogenic fragment of a polypeptide sequence or a complete protein translated according to a polynucleotide sequence selected from the group consisting of ORF1, ORF2, ORF1/1, and ORF2/2 of PTTV genotypes or subtypes PTTV1a-VA, PTTV1b-VA, PTTV2b-VA, and PTTV2c-VA, particularly the ORF1 encoding the capsid protein.

According to one aspect of the present invention, the polynucleotide sequence is selected from the group consisting of ORF1 of PTTV genotypes or subtypes PTTV1a-VA, PTTV1b-VA, PTTV2b-VA, and PTTV2c-VA.

According to another aspect of the present invention, the polynucleotide sequence is ORF1 of PTTV genotype PTTV1a-VA. According to a further aspect of the present invention, the polynucleotide sequence is ORF1 of PTTV genotype PTTV1b-VA. According to yet another aspect of the present invention, the polynucleotide sequence is ORF1 of PTTV subtype PTTV2c-VA.

According to one aspect of the present invention, the polypeptide sequence is selected from the group consisting of sequence set forth in SEQ ID No:13, SEQ ID No:14, SEQ ID No:15, SEQ ID No:16, SEQ ID No:17, SEQ ID No:18, SEQ ID No:19, SEQ ID No:20, SEQ ID No:21, SEQ ID No:22, SEQ ID No:23, SEQ ID No:24, SEQ ID No:25, SEQ ID No:26, SEQ ID No:27, and SEQ ID No:28.

According to another aspect of the present invention, the polypeptide sequence is set forth in SEQ ID No:13. According to another aspect of the present invention, the polypeptide sequence is set forth in SEQ ID No:14. According to a further aspect of the present invention, the polypeptide sequence is set forth in SEQ ID No:16. In one specific embodiment of the present invention, the polypeptide sequence is C-terminal region (aa 310-625) of SEQ ID No:16. According to yet another aspect of the present invention, the polypeptide sequence is set forth in SEQ ID No:20.

According to an additional aspect of the present invention, the vaccine further contains an adjuvant.

The present invention further provides method of immunizing a pig against PTTV viral infection, comprising administering to a pig an immunologically effective amount of the vaccine comprising an immunogenic fragment of a polypeptide sequence or a complete protein translated according to a polynucleotide sequence selected from the group consisting of ORF1, ORF2, ORF1/1, and ORF2/2 of PTTV genotypes or subtypes PTTV1a-VA, PTTV1b-VA, PTTV2b-VA, and PTTV2c-VA.

According to one aspect of the present invention, the method comprises administering the immunogenic fragment or recombinant capsid protein to the pig.

According to another aspect of the present invention, the method comprises administering the vaccine parenterally, intranasally, intradermally, or transdermally to the pig.

According to a further aspect of the present invention, the method comprises administering the vaccine intralymphoidly or intramuscularly to the pig.

The present invention additionally provides a method for diagnosing PTTV1 infection and quantification of PTTV1 load, comprising extracting DNA from a sample suspected of PTTV1 infection, performing polymerase chain reaction (PCR) using primers comprising the sequences set forth in SEQ ID NO:29 and SEQ ID NO:30, and detecting PTTV1 specific amplification. According to one aspect of the present invention, the polymerase chain reaction is a SYBR green real-time PCR.

The present invention further provides a method for diagnosing PTTV2 infection and quantification of PTTV2 load, comprising extracting DNA from a sample suspected of PTTV2 infection, performing polymerase chain reaction (PCR) using primers comprising the sequences set forth in SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31 and SEQ ID NO:32, and detecting PTTV2 specific amplification. According to one aspect of the present invention, the polymerase chain reaction is a SYBR green real-time PCR.

The present invention also provides a method for simultaneously detecting and diagnosing PTTV1 and PTTV2 infection, comprising extracting DNA from a sample suspected of PTTV infection, performing polymerase chain reaction (PCR) using primers comprising the sequences set forth in SEQ ID NO:31 and SEQ ID NO:32, and detecting PTTV1 and PTTV2 specific amplification. According to one aspect of the present invention, the polymerase chain reaction is a SYBR green real-time PCR.

The present invention, in addition, provides a method for simultaneously detecting and diagnosing PTTV1a and PTTV1b infection, comprising extracting DNA from a sample suspected of PTTV1 infection, performing a first polymerase chain reaction (PCR) using primers comprising the sequences set forth in SEQ ID NO:33 and SEQ ID NO:34, performing a second PCR using primers comprising the sequences set forth in SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, and SEQ ID NO:38, and detecting PTTV1a and PTTV1b specific amplification.

The present invention provides a method for diagnosing PTTV infection, comprising immobilizing an immunogenic fragment of a polypeptide sequence translated according to a polynucleotide sequence selected from the group consisting of ORF1, ORF2, ORF1/1, and ORF2/2 of PTTV genotypes or subtypes PTTV1a-VA, PTTV1b-VA, PTTV2b-VA, and PTTV2c-VA; contacting a serum sample from a pig suspected of PTTV infection with the immobilized immunogenic fragment, and detecting captured antibody specific to the immunogenic fragment.

According to one aspect of the present invention, the polynucleotide sequence is selected from the group consisting of ORF1 of PTTV genotypes or subtypes PTTV1a-VA, PTTV2b-VA, and PTTV2c-VA.

According to one embodiment of the present invention, the polynucleotide sequence is ORF1 of PTTV genotype PTTV1a-VA. According to another embodiment of the present invention, the polynucleotide sequence is ORF1 of PTTV genotype PTTV1b-VA. According to a further embodiment of the present invention, the polynucleotide sequence is ORF1 of PTTV subtype PTTV2c-VA.

According to another aspect of the present invention, the polypeptide sequence is selected from the group consisting of sequence set forth in SEQ ID No:13, SEQ ID No:14, SEQ ID No:15, SEQ ID No:16, SEQ ID No:17, SEQ ID No:18, SEQ ID No:19, SEQ ID No:20, SEQ ID No:21, SEQ ID No:22,

SEQ ID No:23, SEQ ID No:24, SEQ ID No:25, SEQ ID No:26, SEQ ID No:27, and SEQ ID No:28.

According to one embodiment of the present invention, the polypeptide sequence is set forth in SEQ ID No:13. According to another aspect of the present invention, the polypeptide sequence is set forth in SEQ ID No:14. According to another embodiment of the present invention, the polypeptide sequence is set forth in SEQ ID No:16. According to a further embodiment of the present invention, the immunogenic fragment is C-terminal region (aa 310-625) of SEQ ID No:16. According to yet another embodiment of the present invention, the polypeptide sequence is set forth in SEQ ID No:20.

The present invention provides three standardized enzyme-linked immunosorbent assays (ELISA) to diagnose PTTV infections and detect antibodies in serum of pigs infected by PTTV genotypes PTTV1a-VA, PTTV1b-VA, and all known subtypes in PTTV species 2.

The ELISA diagnostic tests are based on the bacterial-expressed or baculovirus-expressed recombinant ORF1 capsid protein of PTTV genotypes PTTV1a-VA, PTTV1b-VA, and PTTV2c-VA.

According to another aspect of the present invention, the detecting captured antibody is via Western blot. According to yet another aspect of the present invention, the detecting captured antibody is via enzyme-linked immunosorbent assay (ELISA).

#### BRIEF DESCRIPTION OF THE DRAWINGS

The above-mentioned features of the invention will become more clearly understood from the following detailed description of the invention read together with the drawings in which:

FIGS. 1A and 1B (top panel and bottom panel) represent the schematic diagram of genomic structures, strategies for genomic cloning and assemblies of four prototype U.S. strains of porcine TTV virus group 1 (top panel: species 1) and group 2 (bottom panel: species 2) strains; (PTTV1a-VA=SEQ ID NO: 9, Sd-TTV31=SEQ ID NO: 53, PTTV1bVA=SEQ ID NO: 10, TTV-1p=SEQ ID: 56, TTV-2p=SEQ ID NO: 59, PTTV2b-VA=SEQ ID NO: 11, and PTTV2c-VA=SEQ ID NO: 12);

FIG. 2 represents PASC (pairwise sequence comparisons) distribution of nucleotide sequence comparisons of 121 TTV strains available in GenBank database. The genus, species, types, subtypes and variants and their corresponding percentage of nucleotide sequence identities are displayed;

FIG. 3A illustrates a phylogenetic tree constructed by the neighbor-joining method based upon the full-length genomic nucleotide sequences;

FIG. 3B illustrates a phylogenetic trees constructed based upon deduced amino acid sequences of ORF1 among seven porcine TTV strains;

FIG. 3C illustrates a phylogenetic trees constructed based upon deduced amino acid sequences of ORF1/1 among seven porcine TTV strains;

FIG. 3D illustrates a phylogenetic trees constructed based upon deduced amino acid sequences of ORF2 among seven porcine TTV strains;

FIG. 3E illustrates a phylogenetic trees constructed based upon deduced amino acid sequences of ORF2/2 among seven porcine TTV strains;

FIG. 4 represents an alignment of the full-length amino acid sequences of ORF1 among seven PTTV strains; (PTTV1a-VA=SEQ ID NO: 13, Sd-TTV31=SEQ ID NO: 54,

PTTV1b-VA=SEQ ID NO: 14, TTV-1p=SEQ ID NO: 57, TTV-2p=SEQ ID NO: 60, PTTV2b-VA=SEQ ID NO: 15, and PPT2c-VA=SEQ ID NO: 16);

FIG. 5 represents an alignment of the full-length amino acid sequences of ORF2 among seven PTTV strains (PTTV1a-VA=SEQ ID NO: 17, Sd-TTV31=SEQ ID NO: 55, PTTV1b-VA=SEQ ID NO: 18, TTV-1p=SEQ ID NO: 58, TTV-2p=SEQ ID NO: 61, PTTV2b-VA=SEQ ID NO: 19, and PPT2c-VA=SEQ ID NO: 20);

FIG. 6A illustrates melting curves of PTTV1 real-time PCR products after 40 cycles of amplifications of respective standard template (indicated in blue) and 20 porcine serum samples;

FIG. 6B illustrates melting curves of PTTV2 real-time PCR products after 40 cycles of amplifications of respective standard template and 20 porcine serum samples;

FIGS. 7A-7E illustrate melting curve analysis (MCA) of PTTV1/PTTV2 SYBR green-based duplex real-time PCR;

FIG. 8 represents an alignment of nucleotide sequences located at the N-terminal part of the putative ORF1 among seven PTTV strains (PTTV1a-VA=SEQ ID NO: 9, Sd-TTV31=SEQ ID NO: 53, PTTV1bVA=SEQ ID NO: 10, TTV-1p=SEQ ID NO: 56, TTV-2p=SEQ ID NO: 59, PTTV2b-VA=SEQ ID NO: 11, and PTTV2c-VA=SEQ ID NO: 12);

FIGS. 9A and 9B represent hydrophilicity profiles and conserved regions of the four known porcine TTV2 (TTV-2p=SEQ ID NO: 60, TTV2#472142=SEQ ID NO: 62, PTTV2b-VA=SEQ ID NO: 15, and PPT2c-VA=SEQ ID NO: 16);

FIGS. 10A-10C illustrate the expression and purification of recombinant PTTV2c ORF1 capsid protein;

FIGS. 11A-11C show representative results of Western blot analyses of selected porcine serum samples;

FIG. 12 illustrates the consistency of PTTV2c-ORF1-based Western blot and ELISA;

FIG. 13 shows Box-and-Whisker-plots of PTTV2 serum antibody level by viral load in 138 pigs from different sources;

FIG. 14A illustrates a retrospective evaluation of the viral load of PTTV2;

FIG. 14B illustrates antibody level to PTTV2 ORF1 capsid protein in 10 pigs growing from arrival to two months after arrival;

FIGS. 15A-15C illustrate the expression and purification of PTTV1a and PTTV1b recombinant ORF1 capsid protein; and

FIG. 16 shows examples of PTTV1a-ORF1-based Western blot analyses of selected porcine serum samples from a farm of Wisconsin.

FIGS. 17A-17F represent the schematic diagrams of construction of full-length genomic DNA clones of porcine TTVs. 17A: pSC-PTTV1a (from the US PTTV isolate PTTV1a-VA; GenBank accession no. GU456383). 17B: pSC-PTTV1b (from the US PTTV isolate PTTV1b-VA; GenBank accession no. GU456384). 17C: pSC-PTTV2c (from the US PTTV isolate PTTV2c-VA; GenBank accession no. GU456386). 17D: pSC-2PTTV2c-RR (tandem-dimerized genomes). 17E: TTV2- #471942-full (from the Germany PTTV isolate TTV2-#471942; a gift from Dr. Andreas Gallei, not generated by the applicants). 17F: pSC-2PTTV2b-RR (tandem-dimerized genomes; generated by the applicants based on the clone TTV2-#471942-full). The plasmid backbone used for the cloning of (A)-(D), and (F) was the pSC-B-amp/kan vector (indicated in black). Grey arrows indicated the PTTV genomic copies;

FIGS. 18A and 18B represent the identification of porcine TTV full-length DNA clones by restriction digestion patterns. 18A: BamH I single digestion of pSC-PTTV1a, pSC-PTTV1b and pSC-PTTV2c clones and the backbone vector pSC-B-amp/kan (pSC-B). The 4.3-Kb fragments indicated the size of the backbone vector whereas the 2.8-Kb fragments indicated the inserted PTTV genomes (black arrowheads). 18B: Comparisons of the Hind III single digestion between pSC-PTTV2b and pSC-2PTTV2b-RR (left; derived from the clone TTV2-#471942-full) and Afl II single digestion between pSC-PTTV2c and pSC-2PTTV2c-RR (right). M: DNA markers;

FIGS. 19A and 19B represent the immunofluorescence assay (IFA) results of transfection (19A) or transfected cell passaging (19B) of the concatemered TTV2-#471942-full DNA in PK-15 cells using a PTTV2-specific anti-ORF1 polyclonal antibody (Ab). 19A: Results observed at 5 days post-transfection. 19B: Cells transfected with DNA clones were passaged and used for the IFA detection at 2 days post-passaging. Magnification=200x. DAPI was used to stain the cell nucleus;

FIGS. 20A and 20B represent the IFA results of transfection (20A) or transfected cell passaging (20B) of the concatemered PTTV2c DNA in PK-15 cells using a PTTV2-specific anti-ORF1 Ab. 20A: Results observed at 5 days post-transfection. 20B: Cells transfected with the DNA clones were passaged and used for the IFA detection at 2 days post-passaging. Magnification=200x. DAPI was used to stain the cell nucleus;

FIG. 21 represents the IFA results of transfection of the concatemered PTTV1a DNA in PK-15 cells at 3 days post-transfection using a PTTV1a-specific anti-ORF1 Ab. Magnification=200x. DAPI was used to stain the cell nucleus.

FIGS. 22A and 22B represent the IFA results of transfection of the pSC-2PTTV2b-RR plasmid (22A) or pSC-2PTTV2c-RR plasmid (22B) in PK-15 cells at 3 days post-transfection. Magnification=200x. DAPI was used to stain the cell nucleus.

FIGS. 23A and 23B represent the determination of the in vivo infectivity of the two porcine TTV2 DNA clones, pSC-2PTTV2b-RR and pSC-2PTTV2c-RR, in conventional pigs, respectively. 23A: Changes of viremia or virus titers (copies/ml) as determined by PTTV2-specific real-time PCR. 23B: Seroconversion to IgG anti-porcine TTV2 ORF1 antibodies in pigs. Anti-PTTV2 antibody is plotted as the ELISA optical density (A405). The ELISA cutoff value, indicated by a dashed line in each panel, is 0.4.

## DETAILED DESCRIPTION OF THE INVENTION

In accordance with the present invention, in one specific example, the aforementioned four novel porcine TTV subtypes are isolated from a single boar in Virginia.

In FIG. 1A, both the PTTV1 and PTTV2 genomes are shown in bold and the sizes and directions of the four putative ORFs (ORF1, ORF2, ORF1/1 and ORF2/2) are indicated by arrows. The GC-rich regions are also shown. Dashed-line arcs A and D represent the regions used for detection of PTTV1 and PTTV2 from serum and semen samples by nested PCR, respectively. Dashed-line arcs B and C represent the two overlapping PCR fragments for genomic cloning of PTTV1 whereas dashed-line arcs E and F represent the two overlapping PCR fragments for genomic cloning of PTTV2. The locations of the primers used in the study (see Table 1) are also shown in the corresponding positions.

One boar serum sample (SR#5) that was shown to be positive for both PTTV1 and PTTV2 in the first-round PCR, thus indicative of higher virus load, was used for subsequent full-length genomic cloning of PTTV. Surprisingly, initial attempts to utilize two primer sets (NG372/NG373 and NG384/NG385) of an inverse PCR (Okamoto et al., 2002, supra) designed for cloning of the first PTTV strain Sd-TTV31 to amplify the virus genomic DNA were not successful. No PCR product, was obtained after several trials. Based upon the initial sequence of the region A of PTTV1 and the region D of PTTV2, two new pairs of primers (TTV1-If (SEQ ID NO:1)/TTV1-2340R (SEQ ID NO:2) and TTV1-2311F (SEQ ID NO:3)/TTV1-IR (SEQ ID NO:4)) were subsequently designed to amplify regions B and C spanning the assumed PTTV1 genome, and two additional pairs of primers (TTV2-IF (SEQ ID NO:5)/TTV2-2316R (SEQ ID NO:6) and TTV2-GCF (SEQ ID NO:7)/TTV2-IR (SEQ ID NO:8)) to amplify regions E and F spanning the assumed PTTV2 genome, respectively (FIG. 1A and Table 1). Primers TTV1-2340R (SEQ ID NO:2) and TTV1-2311F (SEQ ID NO:3) were deduced from a common sequence in PTTV1 stains Sd-TTV31 (Okamoto et al., 2002, supra) and TTV-1p (Niel et al., 2005) that is absent in PTTV2 strain TTV-2p (Niel et al., 2005, supra), whereas primers TTV2-2316R (SEQ ID NO:6) and TTV2-GCF (SEQ ID NO:7) were deduced from a sequence of strain TTV-2p that is absent in the two PTTV1 strains. The resulting four different PCR products with expected sizes were each inserted into a blunt-end cloning vector, and the resulting recombinant plasmids were transformed into *Escherichia coli*. Eight to fifteen positive (with white color) bacterial clones for each construct representing fragments B, C, E and F were identified and subsequently sequenced.

TABLE 1

Oligonucleotide primers used for nested PCR and genomic PCR amplifications of porcine TT viruses		
Primer ID	Sequence (5' to 3')	Used for:
TTV1-mF (SEQ ID NO: 45)	TACACTTCCGGGTTTCAGGAGGCT	Detection of porcine TTV1
TTV1-mR (SEQ ID NO: 46)	ACTCAGCCATTTCGGAACCTCAC	Detection of porcine TTV1
TTV1-nF (SEQ ID NO: 47)	CAATTTGGCTCGCTTCGCTCGC	Detection of porcine TTV1
TTV1-nR (SEQ ID NO: 48)	TACTTATATTCGCTTTCGTGGGAAC	Detection of porcine TTV1

TABLE 1-continued

Oligonucleotide primers used for nested PCR and genomic PCR amplifications of porcine TT viruses		
Primer ID	Sequence (5' to 3')	Used for:
TTV2-mF (SEQ ID NO: 49)	AGTTACACATAACCACCAAACC	Detection of porcine TTV2
TTV2-mR (SEQ ID NO: 50)	ATTACCGCCTGCCCGATAGGC	Detection of porcine TTV2
TTV2-nF (SEQ ID NO: 51)	CCAAACCACAGGAAACTGTGC	Detection of porcine TTV2
TTV2-nR (SEQ ID NO: 52)	CTTGACTCCGCTCTCAGGAG	Detection of porcine TTV2
TTV1-IF (SEQ ID NO: 1)	CATAGGGTGTAACCAATCAGATTTAAGGCGTT	Genomic cloning (fragment B)
TTV1-2340R (SEQ ID NO: 2)	GGTCATCAGACGATCCATCTCCCTCAG	Genomic cloning (fragment B)
TTV1-2311F (SEQ ID NO: 3)	CTTCTGAGGGAGATGGATCGTCTGATGA	Genomic cloning (fragment C)
TTV1-IR (SEQ ID NO: 4)	TTGAGCTCCCGACCAATCAGAATTGACT	Genomic cloning (fragment C)
TTV2-IF (SEQ ID NO: 5)	TTGTGCCGGAGCTCCTGAGAGC	Genomic cloning (fragment E)
TTV2-2316R (SEQ ID NO: 6)	AGGTGCTTGAGGAGTCGTCTGCTTG	Genomic cloning (fragment E)
TTV2-GCF (SEQ ID NO: 7)	CTCAAGCACGAGCAGTGGATCCTCTCA	Genomic cloning (fragment F)
TTV2-IR (SEQ ID NO: 8)	TACCCAGGCGGTTAGACACTCAGCTCT	Genomic cloning (fragment F)

Unexpectedly, two groups of sequence data from each construct were identified, indicating that there exist two types of PTTVs in genogroup 1 and genogroup 2 from the same pig. In order to differentiate and assemble the four PTTV strains, sequence comparisons were performed together with the three known PTTV strains, Sd-TTV31, TTV-1p and TTV-2p (FIGS. 1B and 1C).

FIG. 1B illustrates differentiation and assembly of full-length genomic sequences of PTTV1 strains PTTV1a-VA and PTTV1b-VA with PCR fragments B and C that were subsequently cloned. The initiation codons of ORF1 and ORF2 in the fragment B as well as the termination codons of ORF1 in the fragment C are marked by “\*” or “\*”. The corresponding sequences of two known PTTV1 strains, Sd-TTV31 and TTV-1p, are also shown. Conserved sequences are shaded, and dashes indicate nucleotide deletions.

For PTTV1, the initiation codon ATG and the termination codon TGA of the putative ORF1 were located in fragments B and C, respectively (FIG. 1B). The positions of the codons were differed in two PTTV1 groups, the first one identical to Sd-TTV31 and the second one identical to TTV-1p (FIG. 1B). In addition, the ORF2 initiation codons in the two groups were also located at different positions consistent with that of ORF1. Moreover, phylogenetic analyses using four different sequences of the region B (two from the sequencing data and two from strains Sd-TTV31 and TTV-1p) and four different sequences of the region C supported that the first sequence was clustered with Sd-TTV31 and the second was clustered with TTV-1p (data not shown). Therefore, we were able to differentiate and assemble two groups of sequence data from

both fragments B and C into two full-length PTTV1 genomes that were designated as strains PTTV1a-VA (SEQ ID NO:9) and PTTV1b-VA (SEQ ID NO:10), respectively (FIG. 1B).

FIG. 1C illustrates differentiation and assembly of full-length genomic sequences of PTTV2 strains PTTV2b-VA and PTTV2c-VA with PCR fragments E and F that were subsequently cloned. The corresponding sequence of TTV-2p strain is included and the conserved sequences are shaded. Dashes indicate nucleotide deletions. The unique nucleotides within the overlapping region (boxed with dashed-line) for each strain (a continuous “AG” nucleotides for PTTV2b-VA (SEQ ID NO:11) and two single “A” and “G” nucleotides for PTTV2c-VA (SEQ ID NO:12)) are shown, respectively.

Differentiation of the two PTTV2 strains was easier. A unique continuous “AG” nucleotides located in the overlapping region of two PCR fragments was shared by two groups of sequence data from fragments E and F, respectively (FIG. 1C). The assembled full-length genomic sequence represented a PTTV2 strain and was designated as PTTV2b-VA (SEQ ID. NO:11). Similarly, the complete genomic sequence of a second strain designated as PTTV2c-VA (SEQ ID NO:12) was assembled based upon two unique single “A” and “G” nucleotides shared in the overlapping region by another set of sequence data from fragments E and F, respectively (FIG. 1C). Phylogenetic analyses using four sequences from fragments E and F together with the two corresponding sequences from TTV-2p also supported this assignment (data not shown).

The present invention provides four isolated porcine TTV virus genotypes or subtypes that are associated with viral

infections in pigs. This invention includes, but is not limited to, porcine TTV virus genotypes or subtypes PTTV1a-VA, PTTV1b-VA, PTTV2b-VA, and PTTV2c-VA, the virus genotypes or subtypes which have nucleotide sequences set forth in SEQ ID NO:9 (PTTV1a-VA), SEQ ID NO:10 (PTTV1b-VA), SEQ ID NO:11 (PTTV2b-VA), and SEQ ID NO:12 (PTTV2c-VA), their functional equivalent or complementary strand. It will be understood that the specific nucleotide sequence derived from any porcine TTV will have slight

closely-related to PTTV2b-VA (SEQ ID NO:11) in, genomic length, respectively (data not shown).

The assembled genomic sequences of porcine TTV virus genotypes or subtypes PTTV1a-VA (SEQ ID NO:9), PTTV1b-VA (SEQ ID NO:10), PTTV2b-VA (SEQ ID NO:11), and PTTV2c-VA (SEQ ID NO:12) are submitted to Genbank® (*Nucleic Acids Research*, 2008 January; 36(Data-base issue):D25-30) with accession numbers GU456383, GU456384, GU456385, and GU456386, respectively.

TABLE 2

Virus	Porcine TTV species 1				Porcine TTV species 2		
	Type 1a		Type 1b		Subtype 2a	Subtype 2b	Subtype 2c
Strain	PTTV1a-VA	Sd-TTV31	PTTV1b-VA	TTV-1p	TTV-2p	PTTV2b-VA	PTTV2c-VA
Country	USA	Japan	USA	Brazil	Brazil	USA	USA
Full-length (nt)	2878	2878	2875	Uncompleted	Uncompleted	2750	2803
GenBank accession #	GU456383	AB076001	GU456384	AY823990	AY823991	GU456385	GU456386
TATA box	288-291	288-291	288-291	288-291	233-236	233-236	285-288
Putative mRNA 5'-end	316	316	316	316	261	261	313
ORF1							
Size (aa)	635	635	639	637	624	625	625
Exon #	1	1	1	1	1	1	1
Initiation	534	534	517	517	476	476	528
Termination	2441	2441	2436	2430	2350	2353	2405
ORF2							
Size (aa)	73	73	72	72	68	68	68
Exon #	1	1	1	1	1	1	1
Initiation	430	430	428	428	393	393	445
Termination	651	651	646	646	599	599	651
ORF1/1							
Size (aa)	174	174	182	182	178	178	178
Exon #	2	2	2	2	2	2	2
Initiation	534	534	517	517	476	476	528
Splicing	647/648	647/648	642/643	642/643	595/596	595/596	647/648
	2030/2031	2030/2031	2013/2014	2007/2008	1933/1934	1936/1937	1988/1989
Termination	2441	2441	2436	2430	2350	2353	2405
ORF2/2 (ORF3)							
Size (aa)	224	224	228	228	199	199	199
Exon #	2	2	2	2	2	2	2
Initiation	430	430	428	428	393	395	445
Splicing	647/648	647/648	642/643	642/643	595/596	595/596	647/648
	2030/2031	2030/2031	2013/2014	2007/2008	1933/1934	1936/1937	1988/1989
Termination	2487	2487	2485	2479	2330	2333	2385
Polyadenylation signal (AATAAA)	2458-2463	2458-2463	2462-2467	2456-2461	2473-2478	2476-2481	2528-2533

variations that exist naturally between individual viruses. These variations in sequences may be seen in deletions, substitutions, insertions and the like.

The proposed genomic structure for each of the four PTTV strains was analyzed in detail and summarized in Table 2, together with the three known PTTV strains, Sd-TTV31, TTV-1p and TTV-2p. All the four U.S. strains of PTTV have a similar genomic size of 2,878 by (PTTV1a-VA SEQ ID NO:9), 2,875 by (PTTV1b-VA SEQ ID NO:10), 2,750 by (PTTV2b-VA SEQ ID NO:11), and 2,803 by (PTTV2c-VA SEQ ID NO:12), respectively. Both PTTV1a-VA (SEQ ID NO:9) and Sd-TTV31 have the same genomic length. The published sequences of the strains TTV-1p and TTV-2p all have many undetermined nucleotides in the GC-rich region of the UTR. After artificial filling of these nucleotides with the consensus sequences corresponding to PTTV1 and PTTV2, it was shown that the TTV-1p is more closely-related to PTTV1b-VA (SEQ ID NO:10), and that TTV-2p is more

The numbers (except sizes of the full-length genome, ORFs and the exon numbers) indicate the nucleotide (nt) positions on the genome of respective MTV strains.

Two recent studies have identified the transcribed viral mRNAs and the expression of at least six viral proteins during human TTV replication (Mueller et al., 2008, supra; Qiu et al., 2005, supra), which is more than the predicted number of ORFs encoded by human TTV (Okamoto, H., Nishizawa, T., Tawara, A., Takahashi, M., Kishimoto, J., Sai, T., and Sugai, Y. (2000b). TT virus mRNAs detected in the bone marrow cells from an infected individual. *Biochem Biophys Res Commun* 279(2), 700-7), therefore we included the new human TTV genomic information for comparison with the PTTV sequences. The 5'-ends of the mRNA transcripts of human TTV strain P/1C1 were mapped to an "A" that is 25 nt downstream of the TATA-box (Mueller et al., 2008, supra). This starting point, its adjacent sequence (CGAATGGCTG AGTTTATGCCGC (SEQ ID NO:39); the starting point was

underlined) and the distance to the upstream TATA-box (24 nt; Table 2) are very conserved in all seven PTTV strains, suggesting that PTTV and human TTV may utilize a common 5'-end of mRNA for translation.

Five additional completely-conserved regions were identified in the vicinity of the TATA-box among all seven PTTV strains. Two regions of 11 nt each (AGTCCTCATTT (SEQ ID NO:40) and AACCAATCAGA (SEQ ID NO:41)) are located in the upstream of the TATA-box, whereas the remaining three regions (CTGGGCGGGTGCCGGAG of 17 nt (SEQ ID NO:42); CGGAGTCAAGGGGC of 14 nt (SEQ ID NO:43); TATCGGGCAGG of 11 nt (SEQ ID NO:44)) are located between the proposed 5'-end of mRNA and the initiation codon of ORF2. These conserved PTTV-specific sequences may contain the common elements regulating the viral gene expression.

Previously, three ORFs (ORFs 1-3) were proposed in the genome of the three known PTTV strains, respectively (Niel et al., 2005, supra; Okamoto et al., 2002, supra). The four prototype U.S. strains of PTTV identified in this study possess this structure. The corresponding ORF3 in human TTV has been renamed as ORF2/2 since it initiates at the same ATG in ORF2 and remains in the same ORF (extending ORF2) after the splicing (FIG. 1A) (Mueller et al., 2008, supra; Qiu et al., 2005, supra). We follow the nomenclature of human TTV for revising PTTV classification in this study. Human TTV ORF1/1 is a newly identified viral protein that is encoded by two exons in ORF1 (Qiu et al., 2005, supra). ORF1/1 share the identical N- and C-terminal part with ORF1. The PTTV ORF1/1 counterpart was readily identified in all seven PTTV strains (FIG. 1A and Table 2).

The ORF1 and ORF2 are encoded by a ~2.8 kb viral mRNA whereas the ORF1/1 and ORF2/2 are encoded by a spliced viral mRNA with ~1.2 kb in human TTV (Mueller et al., 2008, supra; Qiu et al., 2005, supra). Since these four ORFs were also deduced in PTTV genomes, and since the sequences and positions of the putative splice donor and acceptor sites in the seven PTTV strains are very conserved (Table 2), it is speculated that porcine TTV probably also encodes the two corresponding mRNAs.

Most of the human TTV strains share a genetic similarity with the CAV, encoding a TTV apoptosis-inducing protein (TAIP) in which its CAV counterpart was named apoptin (de Smit, M. H., and Noteborn, M. H. (2009). Apoptosis-inducing proteins in chicken anemia virus and TT virus. *Curr Top Microbiol Immunol* 331, 131-49). The ORF of TAIP is embedded within the ORF2. However, the corresponding TAIP does not exist in porcine TTV. A recent study showed that the expression of apoptin or TAIP was required for CAV replication in cultured cells (Prasetyo, A. A., Kamahora, T., Kuroishi, A., Murakami, K., and Hino, S. (2009). Replication of chicken anemia virus (CAV) requires apoptin and is complemented by VP3 of human torque teno virus (TTV). *Virology* 385(1), 85-92).

Pairwise sequence comparisons (PASC) is a useful method that plots the frequency distribution of pairwise nucleotide sequence identity percentages from all available genomic sequence of viruses in the same family (Bao, Y., Kapustin, Y.,

and Tatusova, T. (2008). Virus Classification by Pairwise Sequence Comparison (PASC). In "Encyclopedia of Virology, 5 vols." (B. W. J. Mahy, and M. H. V. Van Regenmortel, Eds.), Vol. 5, pp. 342-8. Elsevier, Oxford). The different peaks generated by the PASC program usually represent groups of virus genera, species, types, subtypes and strains (FIG. 2). In this study, we performed PASC analyses of TTV using 121 full-length genomic sequences of human and animal TTV-related strains available in GenBank database (FIG. 2). Assuming that TTV members are classified into a separate family, Anelloviridae, the two major peaks, at 36-55% and 55-67% nucleotide sequence identities, represent groups of genera and species, respectively (FIG. 2). Accordingly, a TTV type is defined as a group of TTV having 67-85% nucleotide sequence identity whereas a TTV subtype may be defined as a group of TTV sequences sharing 85-95% nucleotide sequence identity. TTV strains sharing more than 95% nucleotide sequence identity may be further classified into variants (FIG. 2). A similar classification has been proposed using sequences of 103 TTV isolates by Jelcic et al (Jelcic, I., Hotz-Wagenblatt, A., Hunziker, A., Zur Hausen, H., and de Villiers, E. M. (2004). Isolation of multiple TT virus genotypes from spleen biopsy tissue from a Hodgkin's disease patient: genome reorganization and diversity in the hypervariable region. *J Virol* 78(14), 7498-507).

This proposed criteria of TTV classification were applied to phylogenetic analyses of the genomic sequences of the 4 prototype U.S. strains of PTTV and the 3 other known PTTV strains. Pairwise comparison of full-length nucleotide sequences among these strains showed that the four PTTV1 strains have 54.0-56.4% nucleotide sequence identity compared to the three PTTV2 strains (Table 3). Therefore, the previously designated "genogroup" of PTTV in the literature will probably be more appropriate to designate as "species", and PTTV1 and PTTV2 probably should represent porcine TTV species 1 and species 2, respectively. PTTV species 1 consists of two types of viruses designated as type 1a (including Sd-TTV31 and PTTV1a-VA (SEQ ID NO:9)) and type 1b (including TTV-1p and PTTV1b-VA (SEQ ID NO:10)), respectively, since the nucleotide sequence identity between these two types of viruses is between 69.8-70.7% (Table 3). Sd-TTV31 and TTV1a-VA (SEQ ID NO:9) are recognized as variant strains of the same species due to their higher sequence identity (95.1%). However, the two type 1b strains, TTV-1p and PTTV1b-VA (SEQ ID NO:10), may belong to two different subtypes (nucleotide sequence identity=86.4%). For PTTV species 2, three strains are likely to be classified into separate subtypes (TTV-2p for subtype 2a, PTTV2b-VA (SEQ ID NO:11) for subtype 2b, and PTTV2c-VA (SEQ ID NO:12) for subtype 2c, respectively) based upon their 86.5-90.9% nucleotide sequence identity. This proposed new classification system for PTTV was clearly evident in the phylogenetic tree (FIG. 3A). Phylogenetic trees constructed based upon the deduced amino acid sequences of ORF1, ORF1/1, ORF2 and ORF2/2 of PTTV were also consistent with this proposed classification (FIGS. 3B to 3E).

TABLE 3

Pairwise sequence comparison of the full-length genomic sequence of the seven porcine TTV strains							
	Porcine TTV species 1				Porcine TTV species 2		
	Type 1a		Type 1b		Subtype 2a	Subtype 2b	Subtype 2c
	PTTV1a-VA	Sd-TTV31	PTTV1b-VA	TTV-1p	TTV-2p	PTTV2b-VA	PTTV2c-VA
Type 1a							
PTTV1a-VA	—	95.1	70.5	69.8	55.7	55.1	56.2
Sd-TTV31		—	70.7	70.1	55.9	56.0	56.4
Type 1b							
PTTV1b-VA			—	86.4	54.0	54.7	55.2
TTV-1p				—	55.2	54.7	55.4
Subtype 2a							
TTV-2p					—	86.5	86.8
Subtype 2b							
PTTV2b-VA						—	90.9
Subtype 2c							
PTTV2c-VA							—

The data were generated by using the PASC program, and the values indicate % nucleotide sequence identities.

Unique mutations and deletions and/or insertions are scattered throughout the genomes between PTTV species, types and subtypes. For example, the location of ORF1 initiation and termination codons and the ORF2 initiation codons between PTTV type 1a and 1b, which was shown in FIG. 1B as mentioned above, are different. The two PTTV1b strains also have a 2-codon deletion after the ORF2 initiation compared to PTTV1a (FIG. 1B).

Remarkably, both TTV-2p and PTTV2b-VA have a large 52-nt deletion, which is 39 nt upstream of the first 11-nt conserved sequence (AGTCCTCATT (SEQ ID NO:40)) in the UTR, compared to PTTV2c-VA. Due to this deletion, the genomic size of PTTV2b-VA (probably TTV-2p as well) was significantly smaller than that of PTTV2c-VA (Table 2). A number of “subviral” human TTV clones have been isolated from serum samples that are considered as full-length TTV genomes since the ORFs in a majority of these subviral molecules usually remain intact (de Villiers et al., 2009; Leppik et al., 2007). They have variable lengths in the UTR that are completely or partially deleted. The situation of TTV-2p and PTTV2b-VA appears to resemble that of the human TTV subviral molecules, implying that subtypes PTTV2a and PTTV2b might be the subviral molecules derived from subtype PTTV2c. Of note, the 3'-terminal sequence of a nested-PCR primer TTV2-nF (Table 1) that is commonly used for detection of the PTTV2 from field samples by other groups (Ellis et al., 2008, supra; Kekarainen et al., 2007, supra; Kekarainen et al., 2006, supra; Krakowka et al., 2008, supra) is located at both sides of the deletion. Therefore, the current nested-PCR assay for PTTV2 detection is likely not sufficient to identify the genetically diverse strains of PTTV2c subtype.

The source of the isolated virus strain is serum, fecal, saliva, semen and tissue samples of pigs having the porcine TTV viral infection. However, it is contemplated that recombinant DNA technology can be used to duplicate and chemically synthesize the nucleotide sequence. Therefore, the scope of the present invention encompasses the isolated polynucleotide which comprises, but is not limited to, a nucleotide sequence set forth in SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, and SEQ ID NO:12, or its complementary strand; a

25

polynucleotide which hybridizes to and which is at least 67% complementary to the nucleotide sequence set forth in SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, and SEQ ID NO:12, preferably 85% complementary, or more preferably 95% complementary; or an immunogenic fragment selected from the group consisting of an amino acid sequence of ORF1 protein set forth in SEQ ID NO:13 (PTTV1a-VA), SEQ ID NO:14 (PTTV1b-VA), SEQ ID NO:15 (PTTV2b-VA), SEQ ID NO:16 (PTTV2c-VA), an amino acid sequence of ORF2 protein set forth in SEQ ID NO:17 (PTTV1a-VA), SEQ ID NO:18 (PTTV1b-VA), SEQ ID NO:19 (PTTV2b-VA), SEQ ID NO:20 (PTTV2c-VA), an amino acid sequence of ORF1/1 protein set forth in SEQ ID NO:21 (PTTV1a-VA), SEQ ID NO:22 (PTTV1b-VA), SEQ ID NO:23 (PTTV2b-VA), SEQ ID NO:24 (PTTV2c-VA), an amino acid sequence of ORF2/2 protein set forth in SEQ ID NO:25 (PTTV1a-VA), SEQ ID NO:26 (PTTV1b-VA), SEQ ID NO:27 (PTTV2b-VA), SEQ ID NO:28 (PTTV2c-VA). The immunogenic or antigenic coding regions or fragments can be determined by techniques known in the art and then used to make monoclonal or polyclonal antibodies for immunoreactivity screening or other diagnostic purposes. The invention further encompasses the purified, immunogenic protein encoded by the isolated polynucleotides. Desirably, the protein may be an isolated or recombinant ORF1 protein or an ORF2 protein of at least one of the above isolated porcine TTV subtypes, more desirably ORF1 protein.

The ORF1 of porcine TTV is believed to encode a structural and replication-associated protein (Maggi, F., and Bendinelli, M. (2009). Immunobiology of the Torque teno viruses and other anelloviruses. *Curr Top Microbiol Immunol* 331, 65-90). The ORF1-encoding products of seven PTTV strains have 624-635 aa in length and possess a high number of arginine residues at the N-terminus that are thought to have the DNA-binding activity (FIG. 4). In FIG. 4, conserved sequences are shaded. Dashes indicate amino acid deletions. The RCR motifs are boxed with solid lines. Three HVRs (PTTV1-HVRs 1, 2 and 3) of PTTV1 strains and two HVRs (PTTV2-HVRs 1 and 2) of PTTV2 strains are boxed with dashed lines. The connection boundaries of ORF1/1 are indicated by arrows. The predicted rolling-circle replication (RCR) motifs (Ilyina, T. V., and Koonin, E. V. (1992). Con-

served sequence motifs in the initiator proteins for rolling circle DNA replication encoded by diverse replicons from eubacteria, eucaryotes and archaeobacteria. *Nucleic Acids Res* 20(13), 3279-85) are presented at different positions in different PTTV types and subtypes that may be type- or subtype-specific. RCR motif-III (YxxK) is conserved in the PTTV type 1a (aa position 14-17 of PTTV1a-VA SEQ ID NO:13) and type 1b strains (aa position 379-382 of PTTV1b-VA SEQ ID NO:14), respectively, whereas the same conserved motif identified in all three PTTV2 strains is located at aa position 482-485 of PTTV2b-VA SEQ ID NO:15 (FIG. 4). Both PTTV2b-VA SEQ ID NO:15 and PTTV2c-VA SEQ ID NO:16 also have a conserved RCR motif-II (HxQ) at aa position 331-333 of PTTV2b-VA that is absent in TTV-2p (FIG. 4).

The ORF1 proteins of PTTV strains between species 1 and species 2 share very low aa sequence identity with only 22.4 to 25.8%, which makes it difficult to identify significantly conserved aa sequences between the two species (FIG. 4). In PTTV species 1, the aa identity of ORF1 between type 1a and 1b strains are 50.3-52.7%. Three major hypervariable regions (HVR), PTTV1-HVRs 1 to 3, with a relatively high number of aa substitutions, were identified among the four PTTV1 strains, whereas two HVRs (PTTV2-HVRs 1 and 2) were observed among the three PTTV2 strains (FIG. 4): The three PTTV2 strains have an approximately 20-aa deletion in the corresponding PTTV1-HVR1 region. Moreover, the two HVRs of PTTV2 are within the corresponding PTTV1-HVR3 region (FIG. 4). These HVRs are located only in the ORF1 but not in the truncated ORF1/1. They likely play a role in evading the host immune surveillance and helping PTTV to establish a persistent infection, as suggested by studies of human TTV.

The aa sequences of ORF2 differed considerably between the four PTTV1 (PTTV1a-VA SEQ ID NO:17; PTTV1b-VA SEQ ID NO:18) and three PTTV2 (PTTV2b-VA SEQ ID NO:19; PTTV2c-VA SEQ ID NO:20) strains (FIG. 5). However, they share a conserved protein-tyrosine phosphatase (PTPase)-like motif (Wx<sub>7</sub>Hx<sub>3</sub>CxCx<sub>5</sub>H) at the N-terminus (FIG. 4). This motif is also conserved among all human TTV, TTMV and TTMDV strains as well as CAV. The TTMV or CAV ORF2 protein also exhibited a serine/threonine phosphatase (S/T PTPase) activity (Peters, M. A., Jackson, D. C., Crabb, B. S., and Browning, G. F. (2002). Chicken anemia virus VP2 is a novel dual specificity protein phosphatase. *J Biol Chem* 277(42), 39566-73). The dual specificity of the ORF2 protein is thought to regulate host gene transcription, signal transduction and cytokine responses during viral replication. Recently, mutagenesis analyses of two conserved basic aa residues before the last histidine residue of the motif in CAV revealed that the two residues affect virus replication, cytopathology in vitro and attenuation in vivo (Peters, M. A., Crabb, B. S., Washington, E. A., and Browning, G. F. (2006). Site-directed mutagenesis of the VP2 gene of Chicken anemia virus affects virus replication, cytopathology and host-cell MHC class I expression. *J Gen Virol* 87(Pt 4), 823-31; Peters, M. A., Crabb, B. S., Tivendale, K. A., and Browning, G. F. (2007). Attenuation of chicken anemia virus by site-directed mutagenesis of VP2. *J Gen Virol* 88(Pt 8), 2168-75). The two basic aa residues ("KK") are conserved in the three PTTV2 strains. However, only the first basic residue ("R") is retained in the two PTTV1a strains whereas both basic residues are substituted in the PTTV1b strains (FIG. 5). In FIG. 5, dashes indicate amino acid deletions. The five conserved amino acids within the common motif Wx<sub>7</sub>Hx<sub>3</sub>CxCx<sub>5</sub>H (underlined) identified in TTV, TTMV and CAV are shaded. The positions of the two basic aa residues before the last histidine

of the motif, which have been shown to affect virus replication, cytopathology or in vivo attenuation in CAV, are indicated by "".

In summary, the present invention has determined the full-length genomic sequences of four porcine TTV strains representing different genotypes or subtypes in a serum sample of a single boar in Virginia. The finding from this study clearly indicates that, similar to human TTV, multiple PTTV infections with distinct genotypes or subtypes exist and probably are common in pigs. We have also provided new information regarding the genomic organization, the degree of variability and the characteristics of conserved nucleotide and amino acid motifs of PTTV, which will improve the current PCR detection assay, aid in developing reagents for serological diagnostics and help initiate the structural and functional study of PTTV. A new classification of PTTV is also proposed in this study based upon the phylogenetic and genetic analyses of the genomic sequences of seven known PTTV strains.

The present invention also provides methods for diagnostics of porcine TTV infection by detecting viral DNA in samples of porcine TTV infected pigs or other mammals. One preferred embodiment of the present invention involves methods for detecting porcine TTV nucleic acid sequences in a porcine or other mammalian species using oligonucleotide primers for polymerase chain reaction (PCR) to further aid in the diagnosis of viral infection or disease. The diagnostic tests, which are useful in detecting the presence or absence of the porcine TTV viral nucleic acid sequence in the porcine or other mammalian species, comprise isolating viral DNA from samples of porcine TTV infected pigs, or pigs suspected of infection of TTV, and performing SYBR green real-time quantitative PCR using PTTV1-specific (SEQ ID NO:29/SEQ ID NO:30) or PTTV2-specific (SEQ ID NO:31/SEQ ID NO:32) primers.

In another embodiment of the present invention, the diagnostic method may be adapted to simultaneously detect PTTV1 and PTTV2 by using PTTV1/PTTV2 duplex real-time PCR. More specifically, the method comprises isolating viral DNA from samples of porcine TTV infected pigs or pigs suspected of infection of TTV, performing real-time PCR using both PTTV1-specific (SEQ ID NO:29/SEQ ID NO:30) or PVV2-specific (SEQ ID NO:31/SEQ ID NO:32) primers in the same real-time PCR reaction. Since the  $T_m$  value between PTTV1 and PTTV2 can be distinguished by MCA, the presence of PTTV1 and PTTV2 DNA can be simultaneously detected.

In a further embodiment of the present invention, the diagnostic method may employ duplex nested PCR. The method comprises isolating viral DNA from samples of porcine TTV infected pigs or pigs suspected of infection of TTV, performing a first round of PCR using one pair of primers P1ab-mF (SEQ ID NO:33)/P1ab-mR (SEQ ID NO:34), and performing a second round of PCR using a mixture of two pairs of primers, P1a-nF (SEQ ID NO:35)/P1a-nR (SEQ ID NO:36) for detection of PTTV1a, and P1b-nF (SEQ ID NO:37)/P1b-nR (SEQ ID NO:38) for detection of PTTV1b, and visualizing the PCR products.

The above diagnostics methods maybe optimized by one skilled in the art according to well known methods in the art.

Accordingly, an embodiment of the present invention develops two novel singleplex SYBR green real-time PCR assays to quantify the viral loads of two porcine TTV species, respectively. PTTV1- and PTTV2-specific primers were designed to target the extremely conserved regions across six PTTV1 and four PTTV2 full-length genomes available to date, respectively. Another embodiment of the present invention combines the two singleplex assays into a duplex real-

time PCR assay followed by MCA of the viral amplicons that can be identified by their distinct melting temperatures for simultaneous detection of the two porcine TTV species, PTTV1a and PTTV1b. In a third embodiment, a duplex nested PCR assay for simultaneous amplification of the viral DNAs from two types of PTTV1 in the first round PCR and differential detection of types 1a and 1b in the second round PCR was developed for the identification of two types of porcine TTV species, PTTV1a and PTTV1b, in a single sample. These assays represent simple and practical tools for diagnoses of species- or type-specific porcine TTVs.

Potential primers sequences were identified by multiple sequence alignments of 10 available porcine TTV full-length genomes. PTTV1-specific primers TTV1F (SEQ ID NO:29) and TTV1R (SEQ ID NO:30) were designed based upon two conserved genomic regions immediately before the putative ORF2 across six PTTV1 genomes, whereas PTTV2-specific primers TTV2F4 (SEQ ID NO:31) and TTV2R4 (SEQ ID NO:32) were designed based upon two conserved genomic regions immediately after the putative ORF2/2 across four PTTV2 genomes (Table 4). Primers showed no potentials for self- and cross-dimerization. The expected amplicon sizes were a 118-bp fragment from the PTTV1 primers corresponding to the PTTV1b-VA genome and a 200-bp fragment from the PTTV2 primers corresponding to the PTTV2c-VA genome, respectively.

TABLE 4

Oligonucleotide primers used for real-time PCR and duplex nested PCR detections of porcine TTVs.		
Primer ID	Sequence (5' to 3')	Purpose
TTV1F SEQ ID NO: 29	TCCGAATGGCTGAGTTTATGC	PTTV1-specific real-time PCR
TTV1R SEQ ID NO: 30	TCCGCTCAGCTGCTCCT	PTTV1-specific real-time PCR
TTV2F4 SEQ ID NO: 31	GGTGGTAAAGAGGATGAA	PTTV2-specific real-time PCR
TTV2R4 SEQ ID NO: 32	AATAGATTGGACACAGGAG	PTTV2-specific real-time PCR
P1ab-mF SEQ ID NO: 33	TATCGGGCAGGAGCAGCT	Duplex nested PCR
P1ab-mR SEQ ID NO: 34	TAGGGGCGCGCTCTACGT	Duplex nested PCR
P1a-nF SEQ ID NO: 35	CCTACATGAAGGAGAAAGACT	Duplex nested PCR
P1a-nR SEQ ID NO: 36	CCAGCGTCTCCAGGGTC	Duplex nested PCR
P1b-nF SEQ ID NO: 37	AAGCTACCAAGGGCTGG	Duplex nested PCR
P1b-nR SEQ ID NO: 38	GCGGTCT (T/G)GTAGCGGTAGT	Duplex nested PCR

According to one specific embodiment of the present invention, SYBR green simplex real-time PCR using PTTV1- and PTTV2-specific primers can be used specifically to detect porcine TTV1 and TTV2 DNA, respectively. For PTTV1, a standard curve was established over a range of target DNA concentrations per 25  $\mu$ l. The linear range was shown to span  $4.4 \times 10^1$  to  $4.4 \times 10^8$  copies. The minimum detection limit (44 copies) corresponded to a threshold cycle ( $C_t$ ) of 37.57. For PTTV2, standard curve was also generated

and used to detect DNA concentration ranging from  $8.6 \times 10^0$  to  $8.6 \times 10^8$  copies per 25  $\mu$ l reaction. The corresponding  $C_t$  of minimum detection limit (8.6 copies) was 36.53.

According to another specific embodiment of the present invention, SYBR green duplex real-time PCR is utilized for the simultaneous detection of porcine TTV1 and TTV2 DNA. The 7-degree difference of  $T_m$  value between PTTV1 (87.0° C.) and PTTV2 (80.0° C.) made it feasible to distinguish them from one another by the MCA. Therefore, two singleplex assays can be coupled into a duplex real-time PCR assay for the simultaneous detection of PTTV1 and PTTV2. A positive sample was one that had a symmetrical melt peak within the known  $T_m$ , for that product. This new assay was first validated by using a 10-fold dilution of PTTV1 and PTTV2 standards mixture. The non-template negative control using sterile water as the template showed a non-specific amplification caused by cross-dimerization between the PTTV1 and PTTV2 primers not seen in the singleplex assays (FIG. 7a). This produced a distinct melt peak between 72.0° C. and 76.0° C. FIG. 7A shows melt peaks of PTTV1 standard (red;  $T_m=87.0^\circ$  C.), PTTV2 standard (green;  $T_m=80.0^\circ$  C.) and non-template negative control (caused by primer cross-dimerization; black). FIGS. 7B-7E show melt peaks of representative serum samples with distinct viral loads of PTTV1 and PTTV2. FIG. 7B shows boar serum sample no. 5: relatively high viral loads of both PTTV1 and PTTV2, but

PTTV2>PTTV1; FIG. 7C shows boar serum sample no. 12: relatively high viral loads of both PTTV1 and PTTV2, but PTTV1>PTTV2; FIG. 7D shows boar serum sample no. 14: low viral loads of both PTTV1 and PTTV2; FIG. 7E shows boar serum sample no. 10: PTTV1 positive, but PTTV2 negative. The viral loads (unit: genomic copies/ml) of PTTV1 and PTTV2 in each sample that were determined by singleplex real-time PCR were indicated at the top of the corresponding melt peak.

In one example, when the duplex real-time assay was applied to the 20 serum samples of adult boars, samples with relatively high viral loads of both PTTV1 and PTTV2 displayed two distinct melt curves corresponding to PTTV1 and PTTV2 without a non-specific melt peak (FIGS. 7B & 7C), whereas samples with low viral load of either PTTV1 or PTTV2 showed virus-specific as well as non-specific melt curves (FIGS. 7D & 7E). Although the two melt peaks in sample #14 were very small, they were considered positive since they displayed a visually distinct and symmetrical rise and fall at the appropriate  $T_m$  of PTTV1 and PTTV2 (FIG. 7D). In contrast, sample #10 was considered only PTTV1 positive because a symmetrical PTTV2 melt peak was not evidently present (FIG. 7E). These results were consistent with that of the two singleplex assays (Table 5). Moreover, the size and shape of melt peaks qualitatively reflected the corresponding viral load in the detected sample.

According to another aspect of the present invention, duplex nested PCR is used for differential detection of two porcine TTV types, PTTV1a and PTTV1b.

The inventor of the present invention demonstrated the existence of two distinct genotypes, tentatively named PTTV1a and PTTV1b, in porcine TTV species 1. To further determine whether the co-infection of PTTV1a and PTTV1b is common in pigs, a novel duplex nested PCR assay to quickly distinguish between the two was developed. Alignment of porcine TTV genomic DNA sequences identified a conserved genomic region located at the N-terminal part of the putative ORF1 encoding the viral capsid protein (FIG. 8). This region also contains the entire ORF2 and the partial UTR in the upstream. Primers P1ab-mF (SEQ ID NO:33)/P1ab-mR (SEQ ID NO:34) were designed to simultaneously amplify both PTTV1a and PTTV1b DNAs in the first-round PCR. A mixture of PTTV1a-specific primers P1a-nF (SEQ ID NO:35)/P1a-nR (SEQ ID NO:36) and PTTV1b-specific primers P1b-nF (SEQ ID NO:37)/P1b-nR (SEQ ID NO:38) was used to differentially amplify each genotype in the second-round PCR. The final PCR products of PTTV1a and PTTV1b were 162 by and 96 by in sizes, respectively, which could be easily distinguished by gel electrophoresis on a 1% agarose gel stained with ethidium bromide. This assay was not expected to detect PTTV2 DNA due to the specificity of primers (FIG. 8). In FIG. 8, conserved sequences were indicated by dots and shaded. Dashes indicated nucleotide deletions. The locations and directions of three pairs of primers used for duplex nested PCR were marked by arrows.

In one example, the 20 serum samples from adult boars that were subjected to the duplex nested PCR assay were all found to be positive for both PTTV1a and PTTV1b, as determined by visualizing two bands of the expected sizes and subsequent sequencing confirmation of PCR products (data not shown). No PCR products were amplified in the 19 semen samples, which was consistent with the results of PTTV1 conventional nested PCR and real-time PCR assays described above.

Infection of pigs with the two species of porcine TTV has been found back to 1985 in Spanish pig farms according to a retrospective investigation (Segales et al., 2009, supra). However, whether porcine TTVs are associated with any particular pig diseases remains elusive. Since both of porcine TTV species have a high prevalence in domestic pigs, determination of TTV viral loads is presumably more important than assessing the presence of TTV DNA. The level of viral loads in serum and semen samples has been indicated as an important marker for PCVAD in PCV2 infection (Opriessnig et al., 2007, supra). Therefore, establishment of quantitative PTTV-specific real-time PCR assays would help identify potential disease conditions associated with porcine TTVs.

Two TaqMan probe-based real-time PCR assays have recently been described. The singleplex assay developed by a Canadian group was not species-specific and was only designed to quantify the total viral loads of two PTTV species (Brassard et al., 2009, supra). The duplex assay established by a Germany group allowed the specific and simultaneous detection of both species (Gallei et al., 2009, supra). The target sequences of primers used in those two assays were determined by alignment of the three porcine TTV genomic sequences (Sd-TTV31, TTV-1p and TTV-2p) and were located in the UTR. In the present study, with 7 additional complete PTTV genomic sequences available (4 PTTV1 and 3 PTTV2 sequences), we analyzed and re-determined the conserved regions across the 10 full-length PTTV genomes. Based upon the updated alignment result from this study, two species-specific singleplex SYBR green-based real-time PCR assays were developed to quantify the viral loads of PTTV1 and PTTV2, respectively. The primers used in our assays were designed to bind to conserved genomic regions distinct from the previous studies, which may increase the accuracy of quantification. Our assays showed a considerable species-specificity and sensitivity of detection with 44 genomic copies for PTTV1 and 8.8 genomic copies for PTTV2 per 25- $\mu$ l reaction, whereas the detection limit of 10 genomic copies per reaction was reported in the TaqMan probe-based duplex real-time PCR (Gallei et al., 2009, supra). In addition, the SYBR green-based real-time PCR assay is a flexible and inexpensive approach that can be directly carried out without the need to use fluorescently labeled probes. Finally, considering porcine TTVs exhibit a high degree of genetic diversity, the results from SYBR green-based assays are unlikely affected by the different genetic background of porcine TTV variants that likely contain mutations in the probe-binding sequences in the TaqMan probe-based assays.

In spite of the presence of TTV DNA, all serum samples from healthy pigs tested in this study had low amounts of PTTV1 and PTTV2 that were less than  $2 \times 10^6$  copies/ml. Moreover, only an extremely low titer of PTTV2 DNA was detected in three semen samples. Most of the tested serum samples were also positive for PCV2 DNA as determined by conventional nested PCR (data not shown). Many PCV2-positive pigs with low viral load do not develop clinical PCVAD. A proposed threshold for developing PCVAD is  $10^7$  or greater PCV2 genomic copies/ml of serum (Opriessnig et al., 2007, supra). In addition, semen PCV2 DNA-positivity is also a notable marker of diseased status (Opriessnig et al., 2007, supra; Pal, N., Huang, Y. W., Madson, D. M., Kuster, C., Meng, X. J., Halbur, P. G. and Opriessnig, T., 2008. Development and validation of a duplex real-time PCR assay for the simultaneous detection and quantification of porcine circovirus type 2 and an internal control on porcine semen samples. *J Virol Methods* 149, 217-25). The situation of species-specific PTTV may be analogous to that of PCV2 and a high PTTV titer greater than  $10^7$  copies/ml may be required for the induction of porcine diseases. The species-specific real-time PCR assays developed in this study will offer simple and practical tools for future investigations of PTTV association with diseases using a large number of clinical samples from various disease conditions.

Furthermore, by coupling the two species-specific singleplex assays, we developed and validated a quick, inexpensive and reliable screening for the simultaneous detection and differentiation of the two porcine TTV species, PTTV1 and PTTV2, in a MCA-based duplex real-time PCR assay. Although this assay is not intended for accurate quantification of both PTTV species, it is a more convenient approach that

could replace the conventional nested PCR for detection purpose. In comparison with real-time PCR, the conventional nested PCR assay for porcine TTVs detection is time-consuming (requiring total 4 rounds of PCR), laborious and prone to sample contamination occurring during multiple rounds of PCR processing. Due to the difference of  $T_m$  value between PTTV1 and PTTV2 species, an MCA following duplex PCR amplification is able to ensure distinct reaction specificity. Another advantage of this duplex real-time assay is that inclusion of PTTV1 and PTTV2 standards is dispensable when performing the described protocol, which makes it easier for much wider use in any diagnostic labs equipped with an automated real-time PCR instrument.

Multiple infection of porcine TTVs with distinct genotypes or subtypes of the same species has been demonstrated (Gallegi et al., 2009, supra). In particular, our previous study showed that porcine TTV species 1 consists of two distinct types, PTTV1a (including strains Sd-TTV31 and PTTV1a-VA) and PTTV1b (including strains TTV-1p and PTTV1b-VA). The two newly published PTTV1 isolates with full-length genomes, swSTHY-TT27 (GQ120664) from Canada and FTV1 #471819 (GU188045) from Germany, were both classified into type 1b based upon the phylogenetic analysis (data not shown). The duplex nested PCR described in this study confirmed that dual infection of two PTTV1 genotypes frequently occurred in pigs. This novel assay is the first diagnostic PCR approach developed to distinguish between PTTV1a and 1b so far. Since it is currently not known whether one or both of PTTV1a and PTTV1b infection represents a relevant factor associated with diseases, our differential PCR assay should be of great value for future potential disease associations of these two PTTV types.

According to another aspect of the invention, porcine TTV ORF proteins were expressed and used in immunodetection assays to detect the presence of porcine TTV specific antibodies. In one embodiment of the present invention, three truncated and Histidine-tagged ORF1 proteins of PTTV1a, PTTV1b and PTTV2, were expressed and purified in *Escherichia coli* (*E. coli*), respectively. Furthermore, both serum Western blot and ELISA assays based on these recombinant antigens were developed and validated using porcine serum samples from different sources. In particular, serological testing using the PTTV1a-, PTTV1b- and PTTV2-specific ELISA provides an accurate and simple tool for revealing the association of porcine TTV infection with diseases.

According to a further aspect of the invention, porcine TTV ORF proteins were expressed and purified as recombinant ORF1 capsid protein in *E. coli* expression system (FIG. 10, FIG. 15). Three truncated and His-tagged ORF1 capsid proteins of PTTV1a, PTTV1b and PTTV2, were expressed and purified in *Escherichia coli* (*E. coli*), respectively, and served as recombinant capsid subunit vaccines against PTTV infection.

Four porcine TTV2 strains, TTV-2p, TTV2#472142, PTTV2b-VA and PTTV2c-VA, had available complete genomic sequences to date. Although they are phylogenetically classified into three putative subtypes, a comparative analysis of hydrophilicity profiles of the ORF1 encoding amino acids from four PTTV2 showed that they shared three hydrophilic regions, an arginine-rich region from aa 1-49 at the N-terminal and two particular domains (I and II) located at the middle and C-terminal part, respectively (FIG. 9A). The C-terminal region used for truncated PTTV2c-VA ORF1 expression and the corresponding regions shared in other three PTTV2 strains were indicated by a dashed box. Alignments of amino acid sequences demonstrated high levels of

sequence conservation of domains I (aa 322-349) and II (aa 536-625) across the four PTTV2 strains (FIG. 9B).

Since hydrophilic domains are believed to be important for the antigenicity of many proteins, the C-terminal region (aa 310-625) of the PTTV2c-VA ORF1 SEQ ID NO:16 containing the two domains was chosen for protein expression, which would be used as antigen for PTTV2-specific antibody detection in porcine serum. According to one aspect of the invention, expression of the truncated PTTV2c ORF1 was sufficient for detection of all PTTV2 subtypes (2a, 2b and 2c; also see FIG. 3A).

According to one embodiment of the present invention, the C-terminal part of the PTTV2c ORF1 gene fused with 8xHis-tags was constructed and expressed in *E. coli*. The recombinant protein was insoluble and expressed within the bacterial inclusion bodies. FIG. 10A shows SDS-PAGE of unpurified 2c-ORF1 products. FIG. 10B shows SDS-PAGE of purified 2c-ORF1 products. FIG. 10C shows Western blot analysis of purified 2c-ORF1 products using an anti-His-tagged mAb. White arrowheads indicated the ORF1 protein with the expected size and its truncated product whereas black arrowheads show the putative dimers of the expected and truncated proteins. M: protein markers. In FIG. 10A, two significant polypeptides (white arrowheads) were produced in the 2c-ORF1 unpurified sample in comparison with the control sample. The band of ~40 KDa was consistent with the expected size of 2c-ORF1 whereas the ~30 KDa polypeptide was probably an N-terminally truncated product from the former. After purification with a nickel-affinity column, four polypeptides including the two described significant bands were showed in SDS-PAGE (FIG. 10B). They were also detected by western blot using an anti-His-tagged mAb (FIG. 10C). Two high-molecular-mass bands (black arrowheads) were the homodimers formed by the two polypeptides of ~40 KDa and ~30 KDa, respectively, based on the predicted sizes (~80 KDa and ~60 KDa). The results demonstrated that the purified C-terminal PTTV2c-ORF1 was successfully produced and could be used for porcine TTV2 antibody detection in porcine sera.

According to another aspect of the present invention, porcine TTV2 antibodies in various porcine serum samples can be detected by Western blot using purified C-terminal PTTV2c-ORF1. White arrowheads indicated the ORF1 protein with the expected size and its truncated product. It should be noted that only the bands in green color were recognized as positive. A total of more than 200 serum samples of conventional pigs (healthy or diseased), CD/CD pig's and gnotobiotic pigs from different sources were collected. Samples were randomly selected for detection of anti-PTTV2c-ORF1 IgG antibodies using the purified C-terminal PTTV2c-ORF1 as antigen. FIG. 11A shows results of Western blot analyses of selected porcine serum samples of conventional pigs, FIG. 11B shows CD/CD pigs, and FIG. 11C shows gnotobiotic pigs. Purified PTTV2c-ORF1 products were used as the antigens. The two marked ~40 KDa and ~30 KDa bands were detected in most samples of the conventional pigs (FIG. 11A) and CD/CD pigs (FIG. 11B), indicating widely PTTV2 infection in these pigs. However, all the gnotobiotic pigs from two different sources (Blacksburg, Va. and Ames, Iowa) had no detectable PTTV2 antibody (FIG. 11C). Additional low-molecular-mass bands were also observed (FIGS. 11A and 11B). They were likely from non-specific reactivity in the Western blot.

According to yet another aspect of the present invention, PTTV2-specific ELISA can be used as a porcine TTV2 serological test. Seronegative results were also shown in a few samples from conventional pigs of a Wisconsin farm (FIG.

12). These negative samples were pooled and used as a negative reference in development of a PTTV2-specific ELISA. The remaining samples from this source were positive (FIG. 12, the four lanes in the left). In addition, porcine sera from a commercial company used in cell culture (supposed to be OIE diseases-free) also displayed strong anti-PTTV2-ORF2 positivity (FIG. 12), which was used as a positive control for ELISA. The concentrations of purified 2c-ORF1 antigen, porcine sera and IgG conjugate were determined by check-board titration to present low background signal and give the highest difference of OD<sub>405</sub> value between the positive and negative controls. The optimal antigen amount was 69 ng per well, and the optimal ELISA results were obtained by use of a 1:100 dilution of serum samples and a 1:4000 dilution of IgG conjugates. The ELISA cutoff values ranged from 0.25 to 0.5 in each trial. FIG. 4 shows a representative result reflecting the consistency of serum western blot and the developed ELISA.

138 conventional pig sera samples from 3 herds were chosen to analyze the correlation between PTTV2 viral load by real-time PCR and anti-PTTV2 IgG antibody level by ELISA. The results showed that pigs with undetectable or higher PTTV2 viral load (10<sup>8</sup> copies/ml) were more likely to have a lower serum PTTV2 antibody titer than pigs with middle values of PTTV2 viral load (FIG. 13).

In particular, sera from 10 pigs in the same herd were also analyzed by comparing the PTTV2 viral loads and anti-PTTV2 antibody levels of their sera from their arrival in the new facility to two months after arrival. Nine of the 10 pigs had decreased viral loads (three had no detectable virus) after 2 months whilst the anti-PTTV2 antibody titers increased in nine of 10 pigs (FIGS. 14A and 14B). The results suggested that the 10 pigs acquired PTTV2 infection at early stage, which induced humoral response and produced anti-ORF1 capsid IgG antibody progressively. The PTTV2-ORF1 IgG antibody was able to neutralize or even clear the virus, indicating the ORF1 indeed encode a viral capsid protein and may contain neutralizing epitopes against PTTV2.

According to one embodiment of the present invention, the C-terminal PTTV1a- and PTTV1b-ORF1 proteins were expressed and purified in *E. coli* system, respectively. SDS-PAGE and western blot analysis using an anti His-tagged mAb showed that both 1a- and 1b-ORF products had two polypeptides, one with expected size (~40 KDa) and another as the putative homodimer (~80 KDa) (FIG. 15A-C). FIG. 15A shows SDS-PAGE of unpurified and purified 1a-ORF1 products. FIG. 15B shows SDS-PAGE of purified 1b-ORF1 and 1b-ORF1ctruc products. FIG. 15C shows Western blot analysis of purified 1a- and 1b-ORF1 products using an anti-His-tagged mAb. White arrowheads indicate the ORF1 protein with the expected size whereas black arrowheads show the putative dimer of the ORF1 proteins. Compared to 2c-ORF1 expression, no truncated polypeptide was observed. As a comparative control, expression of a C-terminal-truncated 1b-ORF1 region (1b-ORF1ctruc) resulted in a lower-molecular-mass polypeptide compared to its C-terminal-non-truncated counterpart 1b-ORF1 (FIG. 15B).

According one embodiment of the present invention, the purified C-terminal PTTV1a- and PTTV1b-ORF1 proteins were used to develop genotype-specific serum Western blots and ELISA as described for PTTV2 above. FIG. 16 shows negative (lanes 1-2) and positive (lanes 3-5) examples of serum Western blot using 1a-ORF1 as antigen. The same antigen amount (69 ng), dilution of sera (1:100) and dilution of IgG conjugate (1:4000) as PTTV2-ORF1 were used in PTTV1a- and PTTV1b-specific ELISA (data not shown).

Additionally, the present invention provides a useful diagnostic reagent for detecting the porcine TTV infection which comprise a monoclonal or polyclonal antibody purified from a natural host such as, for example, by inoculating a pig with the porcine TTV or the immunogenic composition of the invention in an effective immunogenic quantity to produce a viral infection and recovering the antibody from the serum of the infected pig. Alternatively, the antibodies can be raised in experimental animals against the natural or synthetic polypeptides derived or expressed from the amino acid sequences or immunogenic fragments encoded by the nucleotide sequence of the isolated porcine TTV. For example, monoclonal antibodies can be produced from hybridoma cells which are obtained from mice such as, for example, Balb/c, immunized with a polypeptide antigen derived from the nucleotide sequence of the isolated porcine TTV. Selection of the hybridoma cells is made by growth in hypoxanthine, thymidine and aminopterin in a standard cell culture medium like Dulbecco's modified Eagle's medium (DMEM) or minimal essential medium. The hybridoma cells which produce antibodies can be cloned according to procedures known in the art. Then, the discrete colonies which are formed can be transferred into separate wells of culture plates for cultivation in a suitable culture medium. Identification of antibody secreting cells is done by conventional screening methods with the appropriate antigen or immunogen. Cultivating the hybridoma cells in vitro or in vivo by obtaining ascites fluid in mice after injecting the hybridoma produces the desired monoclonal antibody via well-known techniques.

For another alternative method, porcine TTV capsid protein can be expressed in a baculovirus expression system or *E. coli* expression system according to procedures known in the art. The expressed recombinant porcine TTV capsid protein can be used as the antigen for diagnosis in an enzyme-linked immunoabsorbent Assay (ELISA). The ELISA assay based on the porcine recombinant capsid antigen, for example, can be used to detect antibodies to porcine TTV in porcine and mammalian species. Although the ELISA assay is preferred, other known diagnostic tests can be employed such as immunofluorescence assay (IFA), immunoperoxidase assay (IPA), etc.

Desirably, a commercial ELISA diagnostic assay in accordance with the present invention can be used to diagnose porcine TTV infection in pigs. The examples illustrate using purified ORF1 and ORF2 proteins of porcine TTV to develop an ELISA assay to detect anti-TTV antibodies in pigs. Sera collected from pigs infected with porcine TTV, and negative sera from control pigs are used to validate the assay. PTTV2 specific, PTTV1a specific, and PTTV1b specific antibodies were demonstrated to specifically recognize PTTV ORF proteins. Further standardization of the test by techniques known to those skilled in the art may optimize the commercialization of a diagnostic assay for porcine TTV.

Another aspect of the present invention is the unique immunogenic composition comprising the isolated porcine TTV or an antigenic protein encoded by an isolated polynucleotide described hereinabove and its use for raising or producing antibodies. The composition contains a nontoxic, physiologically acceptable carrier and, optionally, one or more adjuvants. Suitable carriers, such as, for example, water, saline, ethanol, ethylene glycol, glycerol, etc., are easily selected from conventional excipients and co-formulants may be added. Routine tests can be performed to ensure physical compatibility and stability of the final composition.

In accordance with the present invention, there are further provided infectious molecular and nucleic acid molecules of porcine Torque teno (TTV), live viruses produced from the

nucleic acid molecule and veterinary vaccines to protect pigs from porcine TTV viral infection or disease caused by porcine TTV co-infection with other viruses. The invention further provides immunogenic polypeptide expression products that may be used as vaccines.

The novel infectious DNA molecule of porcine TTV comprises a nucleic acid molecule encoding at least a portion of an infectious PTTV1a-VA (SEQ ID NO:9), PTTV1b-VA (SEQ ID NO:10), PTTV2c-VA (SEQ ID NO:11), or PTTV2c-VA (SEQ ID NO:12) genome. The infectious PTTV DNA clone preferably contains at least one of ORF1, ORF2, ORF1/1, and ORF2/2 gene of the PTTV1 or PTTV2. Multiple copies of the PTTV1a-VA (SEQ ID NO:9), PTTV1b-VA (SEQ ID NO:10), PTTV2c-VA (SEQ ID NO:11), or PTTV2c-VA (SEQ ID NO:12) genome may be inserted into a single DNA molecule to construct tandem infectious PTTV clones.

The cloned genomic DNA of PTTV, particularly PTTV1a-VA, PTTV1b-VA, PTTV2c-VA, and tandem PTTV2b-RR, PTTV2c-RR, described herein is shown to be *in vitro* or *in vivo* infectious when transfected into PK-15 cells and given to pigs. This new, readily reproducible pathogenic agent lends itself to the development of a suitable vaccination program to prevent PTTV infection in pigs.

According to a further embodiment of the present invention, three one-genome-copy PTTV DNA clones were derived from the prototype US isolates PTTV1a-VA, PTTV1b-VA and PTTV2c-VA by fusion PCR, respectively. Each of the full-length genomic DNA was inserted into a cloning vector pSC-B-amp/kan by blunt-end ligation. The restriction site BamH I is the unique site on the three PTTV genomes, which was engineered at both ends of the three genomes to facilitate the generation of concatemers and thus mimic the TTV circular genome. BamH I single digestions of the selected plasmid DNA of each clone clearly resulted in two different fragments of 4.3-Kb and 2.8-Kb in size (FIG. 18A). The 4.3-Kb fragments represented the backbone vector whereas the 2.8-Kb fragments represented the inserted PTTV genomic DNA. The empty vector pSC-B-amp/kan digested with the same enzyme only showed a 4.3-Kb band (FIG. 18A). The resulting PTTV clones were designated pSC-PTTV1a, pSC-PTTV1b and pSC-PTTV2c, respectively (FIG. 17A-C).

Furthermore, two copies of the full-length PTTV2c-VA genome derived from the clone pSC-PTTV2c were ligated in tandem into the pSC-B-amp/kan vector to generate the clone pSC-2PTTV2c-RR (FIG. 17D). Comparison of the Afl II single digestion patterns between pSC-PTTV2c and pSC-2PTTV2c-RR showed that the latter plasmid had an additional 2.8-Kb fragment representing the second copy of PTTV2c genome (FIG. 18B, right panel). Subsequently, we utilized the same cloning strategy to produce a tandem-dimerized PTTV2b DNA clone derived from the Germany TTV clone TTV2-#471942-full. An additional 2.8-Kb fragment representing the second copy of PTTV2b genome was presented in this construct, designated pSC-2PTTV2b-RR (FIG. 17F), which was digested with the Hind III alone when compared to its one-genome-copy counterpart (FIG. 18B, left panel), confirming the successful construction.

The replication competencies of the constructed PTTV infectious clones were tested by *in vitro* transfection of PK-15 cells. IFA using the commercially generated rabbit polyclonal antibodies against PTTV2c ORF1 confirmed that both the concatemers of clones TTV2-#471942-full and pSC-PTTV2c were replication competent, respectively (FIG. 19A and FIG. 20A). Passaging of the transfected cells did not eliminate or reduce the fluorescent signals (FIG. 19B and FIG. 20B), suggesting that the expression of ORF1 proteins

was resulted from the PTTV2 concatemers that mimicked the natural PTTV2b or PTTV2c circular molecules. No fluorescent signals was observed in mock-transfected cells or DNA-transfected cells using pre-immune rabbit serum as the antibody for IFA detection (data not shown). The concatemers of the clone pSC-PTTV1a also showed to be replication-competent using an anti-PTTV1a ORF1 antibody (FIG. 21). The positive fluorescent signals were located in the nucleus of transfected or passaged cells, indicating that porcine TTVs likely replicate in the cell nucleus. It is not unexpected because porcine circovirus (PCV) has a similar expression pattern *in vitro*.

Direct transfection of the tandem-dimerized clone pSC-2PTTV2b-RR or pSC-2PTTV2c-RR in PK-15 cells results in viral replication and produces the ORF1 capsid antigen. IFA using antibodies against PTTV2 ORF1 confirmed that both clones were also replication-competent and the positive ORF1 antigens were localized in the nuclei (FIGS. 22A and B).

According to one embodiment of the present invention, infectious clones of porcine TTV can be used to inoculate pigs, which will then elicit an immune response of the host animal and stimulate production of neutralizing antibodies. In one particular embodiment of the present invention, the two tandem-dimerized PTTV2 clones were infectious when injected into the lymph nodes and muscles of conventional pigs.

To test the *in vivo* infectivity of PTTV2 molecular clones, conventional pigs were inoculated with the clone pSC-2TTV2b-RR or pSC-2TTV2c-RR. Serum samples were collected from animals at 0, 7, 14, 21 and 28 days post-inoculation (DPI). PTTV2 DNA was detected in pSC-2TTV2c-RR-inoculated pigs beginning at 7 DPI (#92), 14 DPI (#188 and #191) and 21 DPI (#180), respectively (FIG. 23A, right panel). PTTV viremia appeared late for pigs inoculated with the clone pSC-2TTV2b-RR: two began at 14 DPI (#189 and #192), one at 21 DPI (#181) and one at 28 DPI (#193) (FIG. 23A, left panel). The viral loads increased during the course in all inoculated pigs that had the highest viral loads at 28 DPI before necropsy, as determined by PTTV2-specific real-time PCR (FIG. 23A). The real-time PCR products amplified from selected pigs were sequenced and found to have identical sequences to the corresponding regions of pSC-2TTV2b-RR or pSC-2TTV2c-RR (data not shown).

All inoculated pigs were negative for PTTV2 ORF1 antibodies at 0 and 7 DPI. At 14 DPI, all the four pSC-2TTV2b-RR-inoculated pigs seroconverted to anti-PTTV2 ORF1 IgG, whereas pigs in pSC-2TTV2c-RR-inoculated group seroconverted at 14 (#92 and #180), 21 (#191) and 28 (#188) DPI, respectively (FIG. 23B). The results indicated that active porcine TTV2b or TTV2c infection had occurred.

Vaccines of the infectious viral and infectious molecular DNA clones, and methods of using them, are also included within the scope of the present invention. Inoculated pigs are protected from viral infection and associated diseases caused by TTV2 infection or co-infection. The novel method protects pigs in need of protection against viral infection by administering to the pig an immunologically effective amount of a vaccine according to the invention, such as, for example, a vaccine comprising an immunogenic amount of the infectious PTTV DNA, a plasmid or viral vector containing the infectious DNA clone of PTTV, the recombinant PTTV DNA, the polypeptide expression products, the bacteria-expressed or baculovirus-expressed purified recombinant ORF1 capsid protein, etc. Other antigens such as PRRSV, PPV, other infectious swine agents and immune stimulants may be given

concurrently to the pig to provide a broad spectrum of protection against viral infections.

The vaccines comprise, for example, the infectious viral and molecular DNA clones, the cloned PTTV infectious DNA genome in suitable plasmids or vectors such as, for example, the pSC-B vector, an avirulent, live virus, an inactivated virus, expressed recombinant capsid subunit vaccine, etc. in combination with a nontoxic, physiologically acceptable carrier and, optionally, one or more adjuvants. The vaccine may also comprise the infectious TTV2 molecular DNA clone described herein. The infectious PTTV DNA, the plasmid DNA containing the infectious viral genome and the live virus are preferred with the live virus being most preferred. The avirulent, live viral vaccine of the present invention provides an advantage over traditional viral vaccines that use either attenuated, live viruses which run the risk of reverting back to the virulent state or killed cell culture propagated whole virus which may not induce sufficient antibody immune response for protection against the viral disease.

Vaccines and methods of using them are also included within the scope of the present invention. Inoculated mammalian species are protected from serious viral infection, may also provide protection for disease related to co-infection of PTTV, such as porcine dermatitis and nephropathy syndrome (PDNS), postweaning multisystemic wasting syndrome (PMWS), and other related illness. The vaccines comprise, for example, an inactivated or attenuated porcine TTV virus, a nontoxic, physiologically acceptable carrier and, optionally, one or more adjuvants.

The adjuvant, which may be administered in conjunction with the vaccine of the present invention, is a substance that increases the immunological response of the pig to the vaccine. The adjuvant may be administered at the same time and at the same site as the vaccine, or at a different time, for example, as a booster. Adjuvants also may advantageously be administered to the pig in a manner or at a site different from the manner or site in which the vaccine is administered. Suitable adjuvants include, but are not limited to, aluminum hydroxide (alum), immunostimulating complexes (ISCOMS), non-ionic block polymers or copolymers, cytokines (like IL-1, IL-2, IL-7, IFN- $\alpha$ , IFN- $\beta$ , IFN- $\gamma$ , etc.), saponins, monophosphoryl lipid A (MLA), muramyl dipeptides (MDP) and the like. Other suitable adjuvants include, for example, aluminum potassium sulfate, heat-labile or heat-stable enterotoxin isolated from *Escherichia coli*, cholera toxin or the B subunit thereof, diphtheria toxin, tetanus toxin, pertussis toxin, Freund's incomplete or complete adjuvant, etc. Toxin-based adjuvants, such as diphtheria toxin, tetanus toxin and pertussis toxin may be inactivated prior to use, for example, by treatment with formaldehyde.

The vaccines may further contain additional antigens to promote the immunological activity of the infectious PTTV DNA clones such as, for example, porcine reproductive and respiratory syndrome virus (PRRSV), porcine parvovirus (PPV), other infectious swine agents and immune stimulants.

The new vaccines of this invention are not restricted to any particular type or method of preparation. The cloned viral vaccines include, but are not limited to, infectious DNA vaccines (i.e., using plasmids, vectors or other conventional carriers to directly inject DNA into pigs), live vaccines, modified live vaccines, inactivated vaccines, subunit vaccines, attenuated vaccines, genetically engineered vaccines, etc. These vaccines are prepared by standard methods known in the art.

As a further benefit, the preferred live virus of the present invention provides a genetically stable vaccine that is easier to make, store and deliver than other types of attenuated vaccines.

Another preferred vaccine of the present invention utilizes suitable plasmids for delivering the nonpathogenic DNA clone to pigs. In contrast to the traditional vaccine that uses live or killed cell culture propagated whole virus, this invention provides for the direct inoculation of pigs with the plasmid DNA containing the infectious viral genome.

Additional genetically engineered vaccines, which are desirable in the present invention, are produced by techniques known in the art. Such techniques involve, but are not limited to, further manipulation of recombinant DNA, modification of or substitutions to the amino acid sequences of the recombinant proteins and the like.

Genetically engineered vaccines based on recombinant DNA technology are made, for instance, by identifying alternative portions of the viral gene encoding proteins responsible for inducing a stronger immune or protective response in pigs (e.g., proteins derived from ORF1, ORF1/1, ORF2, ORF2/2, etc.). Such identified genes or immuno-dominant fragments can be cloned into standard protein expression vectors, such as the baculovirus vector, and used to infect appropriate host cells (see, for example, O'Reilly et al., "Baculovirus Expression Vectors: A Lab Manual," Freeman & Co., 1992). The host cells are cultured, thus expressing the desired vaccine proteins, which can be purified to the desired extent and formulated into a suitable vaccine product. The recombinant subunit vaccines are based on bacteria-expressed (FIG. 10, FIG. 15) or baculovirus-expressed ORF1 capsid proteins of PTTV1a, PTTV1b and PTTV2.

If the clones retain any undesirable natural abilities of causing disease, it is also possible to pinpoint the nucleotide sequences in the viral genome responsible for any residual virulence, and genetically engineer the virus avirulent through, for example, site-directed mutagenesis. Site-directed mutagenesis is able to add, delete or change one or more nucleotides (see, for instance, Zoller et al., DNA 3:479-488, 1984). An oligonucleotide is synthesized containing the desired mutation and annealed to a portion of single stranded viral DNA. The hybrid molecule, which results from that procedure, is employed to transform bacteria. Then double-stranded DNA, which is isolated containing the appropriate mutation, is used to produce full-length DNA by ligation to a restriction fragment of the latter that is subsequently transfected into a suitable cell culture. Ligation of the genome into the suitable vector for transfer may be accomplished through any standard technique known to those of ordinary skill in the art. Transfection of the vector into host cells for the production of viral progeny may be done using any of the conventional methods such as calcium-phosphate or DEAE-dextran mediated transfection, electroporation, protoplast fusion and other well-known techniques (e.g., Sambrook et al., "Molecular Cloning: A Laboratory Manual," Cold Spring Harbor Laboratory Press, 1989). The cloned virus then exhibits the desired mutation. Alternatively, two oligonucleotides can be synthesized which contain the appropriate mutation. These may be annealed to form double-stranded DNA that can be inserted in the viral DNA to produce full-length DNA.

An immunologically effective amount of the vaccines of the present invention is administered to a pig in need of protection against viral infection. The immunologically effective amount or the immunogenic amount that inoculates the pig can be easily determined or readily titrated by routine testing. An effective amount is one in which a sufficient immunological response to the vaccine is attained to protect the pig exposed to the PTTV virus. Preferably, the pig is protected to an extent in which one to all of the adverse physiological symptoms or effects of the viral disease are significantly reduced, ameliorated or totally prevented.

The vaccine can be administered in a single dose or in repeated doses. Dosages may range, for example, from about 1 microgram to about 1,000 micrograms of the plasmid DNA containing the infectious chimeric DNA genome (dependent upon the concentration of the immuno-active component of the vaccine), preferably 100 to 200 micrograms of the porcine TTV DNA clone, but should not contain an amount of virus-based antigen sufficient to result in an adverse reaction or physiological symptoms of viral infection. Methods are known in the art for determining or titrating suitable dosages of active antigenic agent to find minimal effective dosages based on the weight of the pig, concentration of the antigen and other typical factors. Preferably, the infectious viral DNA clone is used as a vaccine, or a live infectious virus can be generated *in vitro* and then the live virus is used as a vaccine. In that case, from about 50 to about 10,000 of the 50% tissue culture infective dose (TCID<sub>50</sub>) of live virus, for example, can be given to a pig.

The new vaccines of this invention are not restricted to any particular type or method of preparation. The vaccines include, but are not limited to, modified live vaccines, inactivated vaccines, subunit vaccines, attenuated vaccines, genetically engineered vaccines, etc.

The advantages of live vaccines are that all possible immune responses are activated in the recipient of the vaccine, including systemic, local, humoral and cell-mediated immune responses. The disadvantages of live virus vaccines, which may outweigh the advantages, lie in the potential for contamination with live adventitious viral agents or the risk that the virus may revert to virulence in the field.

To prepare inactivated virus vaccines, for instance, the virus propagation and virus production can occur in cultured porcine cell lines such as, without limitation PK-15 cells. Serial virus inactivation is then optimized by protocols generally known to those of ordinary skill in the art or, preferably, by the methods described herein.

Inactivated virus vaccines may be prepared by treating the porcine TTV with inactivating agents such as formalin or hydrophobic solvents, acids, etc., by irradiation with ultraviolet light or X-rays, by heating, etc. Inactivation is conducted in a manner understood in the art. For example, in chemical inactivation, a suitable virus sample or serum sample containing the virus is treated for a sufficient length of time with a sufficient amount or concentration of inactivating agent at a sufficiently high (or low, depending on the inactivating agent) temperature or pH to inactivate the virus. Inactivation by heating is conducted at a temperature and for a length of time sufficient to inactivate the virus. Inactivation by irradiation is conducted using a wavelength of light or other energy source for a length of time sufficient to inactivate the virus. The virus is considered inactivated if it is unable to infect a cell susceptible to infection.

The preparation of subunit vaccines typically differs from the preparation of a modified live vaccine or an inactivated vaccine. Prior to preparation of a subunit vaccine, the protective or antigenic components of the vaccine must be identified. In the present invention, antigenic components of PTTV were identified as the ORF1 capsid proteins of PTTV1a, PTTV1b and PTTV2, which were expressed and purified in *Escherichia coli* (*E. coli*) in this invention, and other expression system, such as baculovirus expression system, for use as subunit recombinant capsid vaccines. Such protective or antigenic components include certain amino acid segments or fragments of the viral capsid proteins which raise a particularly strong protective or immunological response in pigs; single or multiple viral capsid proteins themselves, oligomers thereof, and higher-order associations of the viral capsid pro-

teins which form virus substructures or identifiable parts or units of such substructures; oligoglycosides, glycolipids or glycoproteins present on or near the surface of the virus or in viral substructures such as the lipoproteins or lipid groups associated with the virus, etc. Preferably, the ORF1 protein is employed as the antigenic component of the subunit vaccine. Other proteins may also be used such as those encoded by the nucleotide sequence in the ORF2, ORF1/1, and ORF2/2 gene. These immunogenic components are readily identified by methods known in the art. Once identified, the protective or antigenic portions of the virus (i.e., the "subunit") are subsequently purified and/or cloned by procedures known in the art. The subunit vaccine provides an advantage over other vaccines based on the live virus since the subunit, such as highly purified subunits of the virus, is less toxic than the whole virus.

If the subunit vaccine is produced through recombinant genetic techniques, expression of the cloned subunit such as the ORF1, ORF2, ORF1/1, and ORF2/2 genes, for example, may be expressed by the method provided above, and may also be optimized by methods known to those in the art (see, for example, Maniatis et al., "Molecular Cloning: A Laboratory Manual," Cold Spring Harbor Laboratory, Cold Spring Harbor, Mass. (1989)). On the other hand, if the subunit being employed represents an intact structural feature of the virus, such as an entire capsid protein, the procedure for its isolation from the virus must then be optimized. In either case, after optimization of the inactivation protocol, the subunit purification protocol may be optimized prior to manufacture.

To prepare attenuated vaccines, the live, pathogenic virus is first attenuated (rendered nonpathogenic or harmless) by methods known in the art or, preferably, as described herein. For instance, attenuated viruses may be prepared by the technique of the present invention which involves the novel serial passage through embryonated pig eggs. Attenuated viruses can be found in nature and may have naturally-occurring gene deletions or, alternatively, the pathogenic viruses can be attenuated by making gene deletions or producing gene mutations. The attenuated and inactivated virus vaccines comprise the preferred vaccines of the present invention.

Genetically engineered vaccines, which are also desirable in the present invention, are produced by techniques known in the art. Such techniques involve, but are not limited to, the use of RNA, recombinant DNA, recombinant proteins, live viruses and the like.

For instance, after purification, the wild-type virus may be isolated from suitable clinical, biological samples such as serum, fecal, saliva, semen and tissue samples by methods known in the art, preferably by the method taught herein using infected pigs or infected suitable cell lines. The DNA is extracted from the biologically pure virus or infectious agent by methods known in the art, and purified by methods known in the art, preferably by ultracentrifugation in a CsCl gradient. The cDNA of viral genome is cloned into a suitable host by methods known in the art (see Maniatis et al., *id.*), and the virus genome is then analyzed to determine essential regions of the genome for producing antigenic portions of the virus. Thereafter, the procedure is generally the same as that for the modified live vaccine, an inactivated vaccine or a subunit vaccine.

Genetically engineered vaccines based on recombinant DNA technology are made, for instance, by identifying the portion of the viral gene which encodes for proteins responsible for inducing a stronger immune or protective response in pigs (e.g., proteins derived from ORF1, ORF2, ORF1/1, and ORF2/2, etc.). Such identified genes or immuno-dominant fragments can be cloned into standard protein expression

vectors, such as the baculovirus vector, and used to infect appropriate host cells (see, for example, O'Reilly et al., "Baculovirus Expression Vectors: A Lab Manual" Freeman & Co. (1992)). The host cells are cultured, thus expressing the desired vaccine proteins, which can be purified to the desired extent and formulated into a suitable vaccine product.

Genetically engineered proteins, useful in vaccines, for instance, may be expressed in insect cells, yeast cells or mammalian cells. The genetically engineered proteins, which may be purified or isolated by conventional methods, can be directly inoculated into a porcine or mammalian species to confer protection against porcine TTV.

An insect cell line (like sf9, sf21, or HIGH-FIVE) can be transformed with a transfer vector containing polynucleic acids obtained from the virus or copied from the viral genome which encodes one or more of the immuno-dominant proteins of the virus. The transfer vector includes, for example, linearized baculovirus DNA and a plasmid containing the desired polynucleotides. The host cell line may be co-transfected with the linearized baculovirus DNA and a plasmid in order to make a recombinant baculovirus.

Alternatively, DNA from the isolated porcine TTV which encode one or more capsid proteins can be inserted into live vectors, such as a poxvirus or an adenovirus and used as a vaccine.

An immunologically effective amount of the vaccine of the present invention is administered to an porcine or mammalian species in need of protection against said infection or syndrome. The "immunologically effective amount" can be easily determined or readily titrated by routine testing. An effective amount is one in which a sufficient immunological response to the vaccine is attained to protect the pig or other mammal exposed to the porcine TTV virus, or porcine TTV co-infection, which may cause porcine dermatitis and nephropathy syndrome (PDNS), postweaning multisystemic wasting syndrome (PMWS) or related illness. Preferably, the pig or other mammalian species is protected to an extent in which one to all of the adverse physiological symptoms or effects of the viral disease are found to be significantly reduced, ameliorated or totally prevented.

The vaccine can be administered in a single dose or in repeated doses. Dosages may contain, for example, from 1 to 1,000 micrograms of virus-based antigen (dependent upon the concentration of the immuno-active component of the vaccine), but should not contain an amount of virus-based antigen sufficient to result in an adverse reaction or physiological symptoms of viral infection. Methods are known in the art for determining or titrating suitable dosages of active antigenic agent based on the weight of the bird or mammal, concentration of the antigen and other typical factors.

The vaccine can be administered to pigs. Also, the vaccine can be given to humans such as pig farmers who are at high risk of being infected by the viral agent. It is contemplated that a vaccine based on the porcine TTV can be designed to provide broad protection against both porcine and human TTV. In other words, the vaccine based on the porcine TTV can be preferentially designed to protect against human TTV infection through the so-called "Jennerian approach" (i.e., cowpox virus vaccine can be used against human smallpox by Edward Jenner). Desirably, the vaccine is administered directly to a porcine or other mammalian species not yet exposed to the TTV virus. The vaccine can conveniently be administered orally, intrabuccally, intranasally, transdermally, parenterally, etc. The parenteral route of administration includes, but is not limited to, intramuscular, intravenous, intraperitoneal and subcutaneous routes.

When administered as a liquid, the present vaccine may be prepared in the form of an aqueous solution, a syrup, an elixir, a tincture and the like. Such formulations are known in the art and are typically prepared by dissolution of the antigen and other typical additives in the appropriate carrier or solvent systems. Suitable carriers or solvents include, but are not limited to, water, saline, ethanol, ethylene glycol, glycerol, etc. Typical additives are, for example, certified dyes, flavors, sweeteners and antimicrobial preservatives such as thimerosal (sodium ethylmercurithiosalicylate). Such solutions may be stabilized, for example, by addition of partially hydrolyzed gelatin, sorbitol or cell culture medium, and may be buffered by conventional methods using reagents known in the art, such as sodium hydrogen phosphate, sodium dihydrogen phosphate, potassium hydrogen phosphate, potassium dihydrogen phosphate, a mixture thereof, and the like.

Liquid formulations also may include suspensions and emulsions which contain suspending or emulsifying agents in combination with other standard co-formulants. These types of liquid formulations may be prepared by conventional methods. Suspensions, for example, may be prepared using a colloid mill. Emulsions, for example, may be prepared using a homogenizer.

Parenteral formulations, designed for injection into body fluid systems, require proper isotonicity and pH buffering to the corresponding levels of mammalian body fluids. Isotonicity can be appropriately adjusted with sodium chloride and other salts as needed. Suitable solvents, such as ethanol or propylene glycol, can be used to increase the solubility of the ingredients in the formulation and the stability of the liquid preparation. Further additives which can be employed in the present vaccine include, but are not limited to, dextrose, conventional antioxidants and conventional chelating agents such as ethylenediamine tetraacetic acid (EDTA). Parenteral dosage forms must also be sterilized prior to use.

The following examples demonstrate certain aspects of the present invention. However, it is to be understood that these examples are for illustration only and do not purport to be wholly definitive as to conditions and scope of this invention. It should be appreciated that when typical reaction conditions (e.g., temperature, reaction times, etc.) have been given, the conditions both above and below the specified ranges can also be used, though generally less conveniently. The examples are conducted at room temperature (about 23° C. to about 28° C.) and at atmospheric pressure. All parts and percents referred to herein are on a weight basis and all temperatures are expressed in degrees centigrade unless otherwise specified.

#### EXAMPLE 1

##### Viral DNA Extraction, Nested PCR and Genomic PCR

Convenient serum and semen samples from 20 conventional adult boars from a Virginia pig farm were used in the study. Total DNA was isolated from 20 serum and 19 semen samples using QIAamp DNA mini kit (Qiagen). To screen for the positive PTTV-containing samples, nested PCR amplifications of the conserved regions in the UTR of PTTV1 and PTTV2 were initially performed by using AmpliTag Gold polymerase (Applied Biosystems). The two primer pairs used to amplify the fragment A of PTTV1 were TTV1-mF (SEQ ID NO:45)/TTV1-mR (SEQ ID NO:46) (for the first-round PCR) and TTV1-nF (SEQ ID NO:47)/TTV1-nR. (SEQ ID NO:48) (for the second-round PCR), whereas the two primer pairs used to amplify the fragment D of PTTV2 were TTV2-mF (SEQ ID NO:49)/TTV2-mR (SEQ ID NO:50) (for the

first-round PCR) and TTV2-nF (SEQ ID NO:51)/TTV2-nR (SEQ ID NO:52) (for the second-round PCR; FIG. 1A and Table 1).

In order to amplify the full-length genomic sequences of both PTTV1 and PTTV2, we first performed an inverse genomic PCR using a pair of conserved gene-specific primers TTV1-IF (SEQ ID NO:1)/TTV (SEQ ID NO:4) located in region A for PTTV1 and another pair of gene-specific primers TTV2-IF (SEQ ID NO:5)/TTV2-IR (SEQ ID NO:8) located in region D for PTTV2, respectively, with Hercules II Fusion DNA Polymerase (Stratagene) according to the manufacturer's instructions. No PCR products with expected sizes were detected. Subsequently we designed new sets of primers to amplify two regions covering the complete PTTV1 and PTTV2 genomes in the second-round PCR, respectively (FIG. 1A). The primer pairs used to amplify fragments B and C of PTTV1 were TTV1-IF (SEQ ID NO:1)/TTV1-2340R (SEQ ID NO:2) and TTV1-2311F (SEQ ID NO:3)/TTV1-IR (SEQ ID NO:4), respectively, whereas the primer pairs used to amplify fragments E and F of PTTV2 were TTV2-IF (SEQ ID NO:5)/TTV2-2316R (SEQ ID NO:6) and TTV2-GCF (SEQ ID NO:7)/TTV2-IR (SEQ ID NO:8), respectively (FIG. 1A and Table 1). Fragments C and F contain the GC-rich regions of PTTV1 and PTTV2, respectively. The amplified PCR products were individually excised, purified, and subsequently cloned into a pSC-B-amp/kan vector (Stratagene) by StrataClone Blunt PCR cloning strategy according to the manufacturer's instructions (Stratagene) followed by DNA sequencing.

#### EXAMPLE 2

Screening for Porcine TTV Positive Samples Collected from Boars in a Farm from Virginia

Porcine TTV DNA was previously detected from pigs in different geographic regions by nested-PCR based on the UTR sequence of a Japanese PTTV1 strain Sd-TTV31 (McKeown et al., 2004, supra). With the recent identification of PTTV2, two different sets of nested-PCR primers have been used to amplify region A of PTTV1 and region D of PTTV2, respectively (FIG. 1A) (Ellis et al., 2008, supra; Kekarainen, T., Sibila, M., and Segales, J. (2006). Prevalence of swine Torque teno virus in post-weaning multisystemic wasting syndrome (PMWS)-affected and non-PMWS-affected pigs in Spain. *J Gen Virol* 87(Pt 4), 833-7; Krakowka et al., 2008, supra). A similar detection approach was also utilized in the present study to identify PTTV strains from pigs in the United States. In order to screen for indigenous PTTV1- or PTTV2-positive samples for subsequent use to determine the full-length genomic sequences, 20 sera (SR#1-20) and 19 semen samples (SM#1-18, and SM#20) collected from 20 boars in a farm of Virginia were subjected to nested-PCR analyses. Surprisingly, all the 20 serum samples were positive for PTTV1 and 19 were also positive for PTTV2 (except for SR#18). In contrast, only 1 semen sample (SM#6) was PTTV1-positive and 3 semen samples (SM#8, 9 and 20) were PTTV2-positive. The result was consistent with a recent study in that boar semen samples were shown to be positive for PTTV DNA in Spain (Kekarainen, T., Lopez-Soria, S., and Segales, J. (2007). Detection of swine Torque teno virus genogroups 1 and 2 in boar sera and semen. *Theriogenology* 68(7), 966-71), and thus suggesting a potential vertical transmission of PTTV. However, the prevalence rates of both PTTV1 and PTTV2 in semen were much lower than that in

sera, suggesting that there is no direct association for the presence of PTTV DNAs in sera and semen of the same pig.

#### EXAMPLE 3

Sequence and Phylogenetic Analyses

Generic analyses and alignment of DNA and amino acid sequences were performed using Lasergene package (DNASTAR Inc., Madison, Wis.). The genomic sequences of three known PTTV strains and their corresponding GenBank accession numbers used for the alignment and comparison are Sd-TTV31 (AB076001), TTV-1p (AY823990) and TTV-2p (AY823991). Pairwise sequence comparisons (PASC) were performed using 121 full-length genomic sequences of human and animal TTV-related strains available in GenBank with an online program PASC (<http://www.ncbi.nlm.nih.gov/sutils/pasc/viridty.cgi?textpage=overview>) (Bao et al., 2008).

Phylogenetic trees were constructed by the neighbor-joining method in the PAUP 4.0 program (David Swofford, Smithsonian Institute, Washington, D.C., distributed by Sinauer Associate Inc.) based upon the full-length genomic sequences and the deduced amino acid sequences of 4 ORFs of seven PTTV strains. The data were obtained from 1000 re-sampling.

#### EXAMPLE 4

Design of PCR Primers for Diagnosing Porcine PTTV Infection

Analyses and alignment of DNA sequences were performed using Lasergene package (DNASTAR Inc., Madison, Wis.). Full-length genomic sequences of ten porcine TTV strains and their corresponding GenBank accession numbers used for the alignment were as follows. Species PTTV1: Sd-TTV31 (AB076001), PTTV1a-VA (GU456383), TTV-1p (AY823990), PTTV1b-VA (GU456384), swSTHY-TT27 (GQ120664) and TTV1 #471819 (GU188045). Species PTTV2: PTTV2b-VA (GU456385), PTTV2c-VA (GU456386), TTV-2p (AY823991) and TTV2 #472142 (GU188046). The conserved sequences among the 6 PTTV1 and 4 PTTV2 genomes were identified, respectively, and subsequently used to guide real-time PCR primer selections using the Beacon Designer program (PREMIER Biosoft International, Palo Alto, Calif.). Primers used for the duplex nested PCR of PTTV1 were designed by the Lasergene package.

#### EXAMPLE 5

Standard Curves of PTTV1 and PTTV2 Real-Time PCR

A region of 2091 bp corresponding to the PCR fragment B of PTTV1b-VA genome was re-amplified from the same PCR fragment using primers TTV1-IF (5'-CAIAGGGTGTAAC-CAATCAGATTAAGGCGTT-3') and TTV1-2340R (5'-GGTCATCAGACGATCCATCTCCCTCAG-3') as described previously (Huang et al., 2010). The resulting amplicon was gel-purified by QIAquick Gel Extraction Kit (Qiagen) and quantified by a NanoDrop spectrophotometer that was used for the real-time PCR standard template of porcine TTV species 1. A full-length DNA clone of PTTV2c-VA strain, pSC-PTTV2c, was constructed by assembling PCR fragments E and F from PTTV2c-VA in the vector pSC-B-amp/kan (Huang et al., unpublished data). Plasmid pSC-PTTV2c (7082 bp) was used for the real-time PCR standard template of porcine TTV species 2 and the plasmid DNA concentration was measured by a NanoDrop spectrophotometer. A 10-fold dilution series of the two templates was used to generate the real-time PCR standard curves, respectively.

## 41

## EXAMPLE 6

## Extraction of Viral DNA for PCR Assays

Total DNA was isolated from 20 serum and 19 semen samples collected from 20 conventional adult boars (with no clinical syndromes) from a Virginia pig farm using QIAamp DNA mini kit (Qiagen) as described previously (Huang et al., 2010). A sample volume of 400  $\mu$ l for sera and semen was used to extract DNA with a final eluate of 50  $\mu$ l sterile water. All extracted DNA samples were stored at  $-20^{\circ}$  C. until real-time PCR testing. Detection of porcine TTVs in these samples by conventional nested PCR had been described previously (Huang et al., 2010). Total DNA extracted from a goat serum sample with the same procedure was used as the negative control.

## EXAMPLE 7

## SYBR Green Real-Time Quantitative PCR Assays

PTTV1- and PTTV2-specific real-time PCR were performed, respectively, using SensiMix SYBR & Fluorescein kit (Quantace Ltd) and the MyiQ iCYCLER Real Time PCR instrument (BIO-RAD Laboratories). Each 25- $\mu$ l reaction contained 12.5  $\mu$ l of SYBR green Master Mix, 4  $\mu$ l of extracted DNA, 0.5  $\mu$ l of each primer (10 nM) and 7.5  $\mu$ l of sterile water. The PCR condition for PTTV1 was 10 min at  $95^{\circ}$  C. followed by 40 cycles of amplification (15 sec at  $95^{\circ}$  C., 30 sec at  $59.4^{\circ}$  C., 10 sec at  $72^{\circ}$  C.). This was immediately followed by a melting point analysis obtained by gradually increasing the temperature from  $55^{\circ}$  C. to  $95^{\circ}$  C. with the fluorescence signal being measured every  $0.5^{\circ}$  C. The PCR condition for PTTV2 was the same as PTTV1 except that the annealing temperature was  $56^{\circ}$  C. PTTV1 and PTTV2 standard templates were included as positive controls in every

## 42

A PTTV1 standard curve was established over a range of target DNA concentrations per 25  $\mu$ l. The linear range was shown to span  $4.4 \times 10^1$  to  $4.4 \times 10^8$  copies. The minimum detection limit (44 copies) corresponded to a threshold cycle ( $C_t$ ) of 37.57. Tested samples with  $C_t > 37.57$  were considered as below the detection limit and were not quantifiable. Similarly, a PTTV2 standard curve was generated and used to detect DNA concentration ranging from  $8.6 \times 10^0$  to  $8.6 \times 10^8$  copies per 25  $\mu$ l reaction. The corresponding  $C_t$  of minimum detection limit (8.6 copies) was 36.53. All samples that were considered as PTTV1- or PTTV2-positive had copy numbers lower than the respective maximum detection limit. Melting curves using a 10-fold dilution of PTTV1 or PTTV2 standard template (FIGS. 6a & 6b; blue curves), as well as 20 boar serum samples, displayed melting temperatures ( $T_m$ ) of  $87.0^{\circ}$  C. for PTTV1 and  $80.0^{\circ}$  C. for PTTV2, respectively (FIGS. 6a & 6b; red curves). No peaks were observed for the negative controls using sterile water or goat serum DNA as templates (FIGS. 6a & 7b; black lines).

## EXAMPLE 9

## Quantification of Porcine TTV1 and TTV2 in Boar Serum and Semen Samples

Viral load was expressed as copy numbers of PTTV1 or PTTV2 genomes per ml of original boar serum samples. PTTV1 DNA were detected in all 20 serum samples ranging from  $1.91 \times 10^3$  to  $3.25 \times 10^5$  copies/ml whereas PTTV2 DNA were detected in 19 serum samples (except #10) ranging from  $3.59 \times 10^2$  to  $1.39 \times 10^6$  copies/ml. The result was consistent to our previous study by using conventional nested PCR (Table 5). None of the semen samples were PTTV1-positive whereas three semen samples were PTTV2-positive with very low viral loads (230, 244 and 357 copies/ml, respectively).

TABLE 5

Comparison of porcine TTVs detection by different assays in 20 serum and 19 semen samples from adult boars in a Virginia Farm.					
Samples	No. of positive/total no. tested by different assay				
	PTTV1 real-time PCR	PTTV1 nested PCR	PTTV2 real-time PCR	PTTV2 nested PCR	PTTV1/PTTV2 duplex real-time PCR
Serum PTTV1	20/20	20/20	—	—	20/20
Serum PTTV2	—	—	19/20	19/20	19/20
Semen PTTV1	0/19	1/19	—	—	—
Semen PTTV2	—	—	3/19	3/19	—

run. Amplification and data analysis were carried out using MyiQ System software (BIO-RAD Laboratories). All samples were run in duplicate on the same plate.

## EXAMPLE 8

## Specificity and Sensitivity of Two Singleplex Assays

The optimal annealing temperatures for amplification of PTTV1- and PTTV2-specific assays were  $59.4^{\circ}$  C. and  $56^{\circ}$  C., respectively, as determined by a 10-fold dilution of amplifications using a gradient of annealing temperatures. Amplification of the 118-bp product using primers TTV1F/TTV1R was obtained only with PTTV1 template whereas amplification of the 200-bp product with PTTV2 template was only observed when primers TTVF4/TTVR4 were used. Neither assay yielded any cross-amplification from the other, confirming the specificity of the primers and targets (data not shown).

## EXAMPLE 10

## PTTV1/PTTV2 Duplex Real-Time PCR Assay

PTTV1/PTTV2 duplex real-time PCR assay was performed in a 25- $\mu$ l PCR system containing 12.5  $\mu$ l of SYBR green Master Mix, 0.5  $\mu$ l of each PTTV1 primers, 0.5  $\mu$ l of each PTTV2 primers, 4  $\mu$ l of DNA and 6.5  $\mu$ l of sterile water. The duplex PCR condition and melting point analysis were the same as PTTV1 except that the annealing temperature was  $58^{\circ}$  C. The melting peaks were analyzed to distinguish the PTTV1- and PTTV2-specific amplicons.

## EXAMPLE 11

## Duplex Nested PCR

The first-round PCR was performed with a Platinum PCR HiFi Supermix (Invitrogen) using 4  $\mu$ l of extracted DNA in a total volume of 50  $\mu$ l. The PCR condition was 30 cycles of  $94^{\circ}$  C. for 30 sec,  $55^{\circ}$  C. for 30 sec,  $72^{\circ}$  C. for 30 sec with an initial

denaturation of the template DNA at 94° C. for 2 min. A 4- $\mu$ l aliquot of the first-round PCR product was used for the second-round PCR with the same PCR reagents and condition. One pair of primers P1ab-mF/P1ab-mR was used in the first-round PCR whereas a mixture of two pairs of primers, P1a-nF/P1a-nR for detection of PTTV1a, and P1b-nF/P1b-nR for detection of PTTV1b, were used in the second-round PCR (Table 1). The amplification products were visualized by gel electrophoresis on a 1% agarose gel stained with ethidium bromide and two bands specific for each type were differentiated by UV light.

## EXAMPLE 12

## Construction of PTTV1 and PTTV2 ORF Expression Plasmids

The C-terminal parts of ORF1 of PTTV1a, PTTV1b and PTTV2c were amplified from the respective full-length DNA clones (pSC-PTTV1a, pSC-PTTV1b and pSC-PTTV2c; described elsewhere). The amplified fragments were expected to encode protein products with 319 aa for PTTV1a (ORF1 aa positions 317-635 (SEQ ID NO:13); GenBank accession no. GU456383), 318 aa for PTTV1b (ORF1 aa positions 322-639 (SEQ ID NO:14); GenBank accession no. GU456384), and 316 aa for PTTV2c (ORF1 aa positions 310-625 (SEQ ID NO:16); GenBank accession no. GU456386), respectively. A C-terminal truncated fragment of PTTV1b encoding 248 aa (ORF1 aa positions 322-569 (SEQ ID NO:14)) was also amplified and used as a comparison control for SDS-PAGE analysis. All the plasmids were constructed by cloning of the PCR products into an *E. coli* baculovirus/mammalian cells triple expression vector pTriEx1.1-Neo (Novagen) between the NcoI and XhoI restriction sites to generate C-terminally 8 $\times$ His-tagged fusion proteins. The four recombinant plasmids were designated pTri-PTTV1a-ORF1, pTri-PTTV1b-ORF1, pTri-PTTV1b-ORF1ctruc and pTri-PTTV2c-ORF1. All cloned sequences were confirmed by DNA sequencing.

## EXAMPLE 13

## Expression of Recombinant PTTV1 and PTTV2 Proteins

The four expression plasmids were transformed into Rosetta 2 (DE3) pLacI competent cells (Novagen), respectively, and the bacteria were plated on LB agar plates containing 100  $\mu$ g/ml ampicillin overnight at 37° C. A single transformation colony for each construct was used to inoculate 3 ml of LB medium containing 100  $\mu$ g/ml of ampicillin (LB/amp), and grown 6-8 hours at 37° C. The turbid 3 ml culture for each construct was then used to make bacterial stocks by adding 25% filter sterilized glycerol, and freezing the culture down at -80° C. Prior to purification, 10  $\mu$ l of the frozen bacterial stock for each construct was used to inoculate a 3 ml starter culture of LB/amp, and grown for 6-8 hours at 37° C. A 100-ml of Overnight Express TB Media (Novagen) was inoculated with the starter culture to induce protein expression, and was grown 16-18 hours at 37° C. After incubation, the autoinduction culture underwent centrifugation at 3400 rpm for 15 minutes at 4° C. The resulting supernatant for each construct was discarded, and each of the bacterial pellets was reserved at -20° C. until use.

## EXAMPLE 14

## Purification and Dialysis of Recombinant Proteins

The recombinant proteins were insoluble and expressed within the bacterial inclusion bodies. Each of the bacterial

pellets was treated with BugBuster and rLysozyme according to the manufacture's protocol (Novagen), and Benzonase Nuclease (Novagen) was added for degradation of DNA and RNA. Each of the inclusion body pellets was subsequently resuspended with 840  $\mu$ l of lysis buffer (6M Guanidine Hydrochloride, 0.1M sodium phosphate, 0.01M Tris-Chloride, 0.01M imidazole, pH 8.0), and frozen at -80° C. for at least 30 minutes. It was then thawed, diluted with an additional 2.5 ml of lysis buffer and gently rotated for 30 minutes at room temperature. The lysate supernatants were collected by centrifugation at 15,000 $\times$ g for 30 minutes at room temperature. A 50%-Ni-NTA His-bind slurry (Novagen) was added to each of the decanted supernatants, and the mixtures were shaken for 60 minutes at room temperature to promote his-tag binding. The lysate/resin mixtures were loaded into an empty chromatography column. After the initial flow-through, a 7-ml of lysis buffer was added to the column and allowed to flow through. Each column was then washed 2 times with 7 mL of wash buffer (8M Urea, 0.1M Sodium Phosphate, 0.15M Sodium Chloride, 0.02M imidazole, pH 8.0). Elution of the target protein was achieved by adding 4 separate 1 ml aliquots of elution buffer (8M Urea, 0.05M Sodium Phosphate, 1M Sodium Chloride, 0.5M Imidazole, pH 8.0) to the column. The four elution fractions were analyzed by SDS Page and Coomassie Blue Staining.

The elutions containing significant concentrations of the target protein were injected into a 0.5 ml-3 ml dialysis cassette with a 20,000 molecular weight cut-off (Pierce). A series of 4 dialysis buffers were used for dialysis; dialysis buffer 1 (6M Urea, 0.05M Sodium Phosphate, 0.8M Sodium Chloride, 0.3M Imidazole, pH 8.0), dialysis buffer 2 (4M Urea, 0.033M Sodium Phosphate, 0.533M Sodium Chloride, 0.2M Imidazole, pH 8.0), dialysis buffer 3 (2.67M Urea, 0.022M Sodium Phosphate, 0.356M Sodium Chloride, 0.133M Imidazole, pH 8.0) and dialysis buffer 4 (1.5M Urea, 0.0148M Sodium Phosphate, 0.237M Sodium Chloride, 0.089M Imidazole, pH 8.0). The dialysis cassette was sequentially submerged and rotated in each dialysis buffer for over 6 hours at 4° C. When dialysis was complete, the recombinant His-tagged fusion proteins were each removed from the cassettes, quantified using a NanoDrop and frozen at -80° C.

## EXAMPLE 15

## SDS-PAGE and Anti-His-Tagged Western Blot

A western blot was developed to detect purified recombinant proteins by using an anti-6 $\times$ His-tagged monoclonal antibody (Rockland). Equal volumes of each of the purified truncated ORF1 proteins and LDS/10%  $\beta$ -ME were mixed, and boiled at 95° C. for 10 minutes. A 10- $\mu$ l of the boiled sample was added to each appropriate well of a 4-12% Bis-Tris Polyacrylamide Gel (Invitrogen), and was run at 200 volts for 43 minutes in 1 $\times$ MES running buffer (Invitrogen). The proteins were transferred to a PVDF membrane (Bio-Rad) using a Trans blot semi dry transfer apparatus and 1 $\times$  transfer buffer (Invitrogen). Once transfer was complete, the PVDF membrane was incubated in Odyssey blocking buffer (Li-Cor) at room temperature for 1 hour. The anti-6 $\times$ His-tagged MAb was diluted at 1:1000 in Odyssey blocking buffer/0.2% tween 20, and transferred to the membrane after the previous Odyssey blocking buffer was removed. The MAb was left on a rocker to incubate with the membrane for either 2 hours at room temperature or 4° C. overnight, and then the membrane was washed 3 times with tris buffered saline/0.05% tween 20 (TBS-T, Sigma). A Goat anti-rabbit IgG IRDye 800 (Li-Cor) antibody was diluted at 1:5000 in Odyssey blocking buffer/0.2% tween 20/0.1 SDS. It was transferred to the freshly

## 45

washed PVDF membrane, and allowed to incubate for 1 hour at room temperature while gently rocking. The membrane was washed 3 times with TBS-T, 1 time with TBS and imaged with the Li-Cor Odyssey.

## EXAMPLE 16

## Serum Western Blot

A serum western blot was developed, and used to identify positive and negative serum controls for ELISA development. After SDS-PAGE as described above, the proteins were transferred to a PVDF membrane that was subsequently incubated in Odyssey blocking buffer (Li-Cor) at room temperature for 1 hour. A selected serum sample was diluted at 1:100 in Odyssey blocking buffer/0.2% tween 20, and transferred to the membrane after the previous Odyssey blocking buffer was removed. The serum sample was left on a rocker to incubate with the membrane for 2 hours at room temperature, and then the membrane was washed 3 times with tris buffered saline/0.05% tween 20 (TBS-T, Sigma). A goat anti-swine IgG IRDye 800 antibody (Rockland) was diluted at 1:2500 in Odyssey blocking buffer/0.2% tween 20/0.1% SDS. It was transferred to the freshly washed PVDF membrane, and allowed to incubate for 1 hour at room temperature while gently rocking. The membrane was washed 3 times with TBS-T, 1 time with TBS and imaged with the Li-Cor Odyssey.

## EXAMPLE 17

## Indirect PTTV1a-, PTTV1b- and PTTV2-Specific ELISA

The optimal concentrations of the antigens used to coat the plates and dilutions of antisera and conjugates were determined by checkboard titration. The ELISA was initiated by diluting each of the purified recombinant His-tagged fusion proteins (PTTV1a, PTTV1b and PTTV2c, respectively) to 680 ng/ml in 1×Carbonate Coating Buffer (CCB) at a pH of 9.6, and coating medium binding ELISA plates (Greiner) with 100 µl/well. The plates were covered, and allowed to incubate at 37° C. for 2 hours. After coating, the diluted proteins were removed, and each well was washed 3 times with 300 µl of 1×TBS-T. Protein Free Blocking Buffer (Pierce) was then added at a volume of 300 µl/well, and the plates were allowed to incubate at 37° C. for 1 hour. Meanwhile, in a 96-well dilution block, the serum samples were diluted at 1:100 in 150 µl of protein free blocking buffer. The block was then removed, and 100 µl of each diluted serum sample was transferred to each corresponding well on the ELISA plates. The plates were allowed to incubate at 37° C. for 2 hours, after which each well was washed 3 times with 300 µl of TBS-T. Next, the HRP-conjugated anti-swine IgG antibody (Rockland) was diluted at 1:4000 in 12 ml of protein free block, and 100 µl was added to each well of the plates. This was incubated at 37° C. for 1 hour, and then each well was washed 3 times with 300 µl of TBS-T. In order to develop the ELISA, 100 µl of Sure Blue Reserve 1-Component (KPL) was added to each well of the plates. After 20 minutes, 100 µl of 1N HCL was added to each well to stop development. The plates were then read at 450 nm.

## EXAMPLE 18

## Data Analyses

Porcine sera used in cell culture research from a commercial company (manufactured in New Zealand and considered free from all OIE diseases) were used as a positive control for the three ELISA protocols because the sera were all

## 46

PTTV1a-, PTTV1b- and PTTV2-positive as detected by serum western blot and displayed high OD values (>2.0). We initially used pooled gnotobiotic pig sera as a negative control as they were negative in western blot detection. Subsequently, in comparison of the negative gnotobiotic pig sera, we screened some porcine sera collected from a conventional pig farm in Wisconsin. They were also negative in western blot detection and their OD values corresponded to that of negative gnotobiotic pig sera. These conventional porcine sera were pooled and used as a negative control. The cutoff value for each ELISA was calculated as the mean OD value of the negative control group (n=4) plus 3 times of the standard deviation.

## EXAMPLE 19

## Construction of Full-Length Genomic DNA Clones of Porcine TTV1a, 1c and 2c

PCR fragments B and C from the US isolate PTTV1a-VA (GenBank accession no. GU456383) were re-amplified from the constructs described previously, and were subsequently assembled into a full-length genomic DNA with a BamH I site at the both ends of the genome by overlapping PCR using the Herculase II Fusion DNA Polymerase (Stratagene) on the vector pSC-B-amp/kan (Stratagene). The resulting construct was designated pSC-PTTV1a (FIG. 17A). Using the same strategy, the clone pSC-PTTV1b (FIG. 17B) originated from the US isolate PTTV1b-VA (GenBank accession no. GU456384) and the clone pSC-PTTV2c (FIG. 17C) originated from the US isolate PTTV2c-VA (GenBank accession no. GU456386) were constructed with the same restriction sites (BamH I) on the same backbone vector. Plasmid TTV2-#471942-full (FIG. 17E) containing a full-length genomic DNA originated from a Germany pathogenic porcine TTV2 isolate. TTV2-#471942 was a gift from Dr. Andreas Gallei (BIVI, Germany). TTV2-#471942 was classified into the porcine TTV subtype 2b together with the US isolate PTTV1b-VA based upon the phylogenetic analysis (data not shown).

## EXAMPLE 20

## Construction of Tandem-Dimerized DNA Clones of Porcine TTV2b and 2c

The full-length PTTV2c genome was excised from the clone pSC-PTTV2c by BamH I digestion, purified and ligated to form concatemers. Ligated concatemers were cloned into the BamH I-pre-digested pSC-B-amp/kan vector to produce a tandem-dimerized DNA clone, pSC-2PTTV2c-RR (FIG. 1D). Similarly, a tandem-dimerized DNA clone, pSC-2PTTV2b-RR, was generated from the clone TTV2-#471942-full using EcoR V restriction sites (FIG. 1F).

## EXAMPLE 21

## Generation of PTTV1a-, PTTV1b- and PTTV2-Specific Anti-ORF1 Polyclonal Antibodies

The ORF1-encoding product is the putative capsid protein of TTV. To generate PTTV1a-, PTTV1b- and PTTV2-specific anti-ORF1 polyclonal antibodies to detect the expression of PTTV ORF1 proteins and to determine the infectivity of PTTV DNA clones, the three ORF1 proteins from PTTV1a, PTTV1b and PTTV2c were expressed in *E. coli*, purified and were subsequently used to immunize New Zealand white rabbits, respectively, as a custom antibody production service at Rockland Immunochemicals (Gilberts-

ville, Pa.). Each anti-ORF1 polyclonal antibody was produced from serum of immunized rabbits.

## EXAMPLE 22

## In Vitro Transfection of PTTV Infectious Clones

PK-15 cells were seeded at  $2 \times 10^5$  cells per well onto a 6-well plate and grown until 60%-70% confluency before transfection. The DNA clones pSC-2PTTV2b-RR and pSC-2PTTV2c-RR were directly transfected into PK-15 cells, respectively, using Lipofectamine LTX (Invitrogen) according to the manufacturer's protocol. For clones pSC-PTTV1a, pSC-PTTV2c and TTV2-#471942-full, their ligated concatamers, produced as described above, were used for transfection, respectively. Cells were cultured for 3 to 5 days, and were then applied to an immunofluorescence assay (IFA) to detect the expression of ORF1 of porcine TTVs. Alternatively, transfected cells were passaged into new 6-well plates and continued to culture for 3 days before the IFA detection.

## EXAMPLE 23

## Immunofluorescence Assay (IFA)

Transfected or passaged cells were washed 2 times with PBS and fixed with acetone. Five hundred microliters of the antibodies, specific to PTTV1a or PTTV2 at 1:500 dilution in PBS, was added over the cells and incubated for 1 hour at room temperature. Cells were washed 3 times with PBS and 500  $\mu$ l Texas red- or Alexa Fluor 488-labeled goat anti-rabbit IgG (Invitrogen) at 1:200 dilution was then added. After 1-hour incubation at room temperature and washed with PBS, the cells were stained with 500  $\mu$ l DAPI (KPL, Inc.) at 1:1000 dilution and visualized under a fluorescence microscope.

## EXAMPLE 24

## In Vivo Inoculation of Conventional Pigs with the Tandem-Dimerized Porcine TTV2 Clones

5 A pig inoculation study was performed to determine the infectivities of the two tandem-dimerized porcine TTV2 clones: pSC-2TTV2b-RR and pSC-2TTV2c-RR. Briefly, eight 4-week-old conventional pigs that were seronegative and viral DNA negative for porcine TTV2 were randomly assigned into two groups of four each. Each group of pigs was housed separately and maintained under conditions that met all requirements of the Institutional Committee on Animal Care and Use.

15 All pigs in each group were injected by a combination of both the intra-lymph node route and intramuscular route. The four pigs (nos. 181, 189, 192 and 193) were each injected with 200  $\mu$ g of the pSC-2TTV2b-RR plasmid DNA whereas another four pigs (nos. 92, 180, 188 and 191) were each inoculated with 200  $\mu$ g of the pSC-2TTV2c-RR clone. Pigs were monitored daily for clinical signs of disease for a total of 28 days. All pigs were necropsied at 28 days postinoculation.

20 While the present invention has been illustrated by description of several embodiments and while the illustrative embodiments have been described in detail, it is not the intention of the applicants to restrict or in any way limit the scope of the appended claims to such detail. Additional modifications will readily appear to those skilled in the art. The invention in its broader aspects is therefore not limited to the specific details, representative apparatus and methods, and illustrative examples shown and described. Accordingly, departures may be made from such details without departing from the spirit or scope of applicants' general inventive concept.

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 62

<210> SEQ ID NO 1  
<211> LENGTH: 32  
<212> TYPE: DNA  
<213> ORGANISM: Torque teno virus

<400> SEQUENCE: 1

catagggtgt aaccaatcag atttaaggcg tt

??

<210> SEQ ID NO 2  
<211> LENGTH: 27  
<212> TYPE: DNA  
<213> ORGANISM: Torque teno virus

<400> SEQUENCE: 2

ggtcacaga cgatccatct cctcag

27

<210> SEQ ID NO 3  
<211> LENGTH: 28  
<212> TYPE: DNA  
<213> ORGANISM: Torque teno virus

<400> SEQUENCE: 3

cttctgaggg agatggatcg tctgatga

<210> SEQ ID NO 4  
<211> LENGTH: 28  
<212> TYPE: DNA  
<213> ORGANISM: Torque teno virus

-continued

---

<400> SEQUENCE: 4  
 ttgagctccc gaccaatcag aattgact 28

<210> SEQ ID NO 5  
 <211> LENGTH: 22  
 <212> TYPE: DNA  
 <213> ORGANISM: Torque teno virus

<400> SEQUENCE: 5  
 ttgtgccgga gctectgaga gc 22

<210> SEQ ID NO 6  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Torque teno virus

<400> SEQUENCE: 6  
 aggtgcttga ggagtcgctg cttg 24

<210> SEQ ID NO 7  
 <211> LENGTH: 27  
 <212> TYPE: DNA  
 <213> ORGANISM: Torque teno virus

<400> SEQUENCE: 7  
 taccagcg gttagacact cagctct 27

<210> SEQ ID NO 8  
 <211> LENGTH: 27  
 <212> TYPE: DNA  
 <213> ORGANISM: Torque teno virus

<400> SEQUENCE: 8  
 ctcaagcacg agcagtgat cctctca 27

<210> SEQ ID NO 9  
 <211> LENGTH: 2878  
 <212> TYPE: DNA  
 <213> ORGANISM: Torque teno virus

<400> SEQUENCE: 9  
 tacacttcg ggttcaggag gctcaatttg gctagcttcg ctgcaccac gtttgetgcc 60  
 aggcggacct gattgaagac tgaaaaccgt taaattcaaa tttgaaaatg gcgggcaaaa 120  
 tggcggaagg gggcgaggat ttatgcaaat taatttatgc aaagtaggag gagctcgatt 180  
 ttaatttatg caaagtagga ggagtcaatt ctgattggtc gggagctcaa gtcctcattt 240  
 gcataggggtg taaccaatca gatttaaggc gttcccccaa aagtgaatat aagtaagcgc 300  
 agttccgaat ggctgagttt atgccgccag cggtagacag aactgtctag cgactgggcg 360  
 ggtgccggag gatccctgat ccggagtcaa gggcctatc gggcaggagc agctgagcgg 420  
 agggcctaca tgaaggagaa agactactgg gaggaagcct ggctgaccag ctgtacatct 480  
 atacacgacc accactgcaa ctgctgtagc tggagagacc acctgtggac gctatgcgct 540  
 ttagacgacg cagatttggc cgccgccga gatattatag aaagagaaga ggcggatgga 600  
 ggagaagatt tcgattcgt agacggagac cctggagacg ctggcgggta aggagatggc 660  
 ggcgttccgt ctccgtaga gggggacgta gagcgcgccc ctaccgcatt agcgcttga 720  
 accctaaggt tctcagaaac tgccgcatca cgggatgggt gccagttata cagtgtatgg 780

-continued

---

```

acgggatgga gtggataaaa tacaagccta tggacttaag agtcgaggca aactggatat 840
tcaataaaca ggacagtaaa atagagacag aacagatggg atacctgatg cagtatggag 900
gagggtggtc aagcggagta atcagcttag agggactatt caatgaaaac agactgtgga 960
gaaatataat gtcaaaaagc aatgacggga tggacttggg cagatacttt ggctgtagaa 1020
ttagactata tccaacagag aatcaggact acttgttctg gtatgacaca gaatttgacg 1080
aacagcaaag gagaatgcta gatgaataca cacaacctag tgtgatgctg caggctaaaa 1140
actcgcgtct aatagtgtgt aaacagaaga tgccaattag acgcagagta aaaagtattt 1200
ttataccgcc gcctgcacag ttaacaactc agtggaaatt tcaacaggaa ctatgtcagt 1260
ttccactggt taactgggcc tgtatctgca tagacatgga cacgccgttc gactacaacg 1320
gcgcatggcg aaatgcctgg tggctaata gaagcctgca aaacggaaac atggagtaca 1380
tagaaagatg gggcagaata ccaatgacag gagacacaga actaccacca gcagacgact 1440
tcaaggcagg aggggtgaac aaaaacttca aaccgacagg tattcaaga atatacccgga 1500
tagtagcggg atgccttcta gaagggaaca aaagagtagt caaatgggcc acagtacaca 1560
atggtcccat agacagatgg agaaaaaac agacaggaac ttaaaagctc tctaacttga 1620
gaggcctagt actgagagta tgcacagaga gtgaaacata ctataagtgg acaggatcag 1680
aatttacagg ggcatttcaa caagactggt ggcagtagg cggaaacagaa taccgcttt 1740
gtaccattaa aatggacca gaatatgaaa accctacagt agaggtagg tcctggaag 1800
caaatatacc gacatcaggg actcttaaag actacttcgg actgagtaca gggcaacagt 1860
ggaaagacac tgactttgcg aggctgcaac tacctagaag cagccacaat gtggactttg 1920
gacataaagc tagatttggg ccatthtgcg ttaaaaagcc tccagtagag ttcagagata 1980
cagcccaaaa cccactaaat atatgggtaa aatacacggt ctatthttag ttcggcggca 2040
tgtaccagcc tcccaccgga atccaagatc cctgcacttc taaccgacc tatcctgtca 2100
gaatggctcg agcagttaca caccacaaat acgcccggca aggcggaatc acgacccaaa 2160
ttggagatca aggtatcacc gctgcctcta tccgtgccat cagtgcagct ccaccagata 2220
cctacacgca gtcggcgttc ctcaaagccc cggaaaccga gaaagaagag gaaagagaga 2280
gtgagaccag tttcacagat gccgaaagct cttctgaggg agatggatcg tctgatgacc 2340
aagcagagag acgcgctgcc agaaagcagag tcatcaagtt acttctcaag cgactcgtg 2400
acagaccctg ggacaacaag cgacgacgat tttcagagt accctgacc cctcaccat 2460
aaacgcaaaa aacgcttga attctaactc tgtctctgtg acttcattgg gggggtccgg 2520
gggggcttgc cccccgta gttgggttct cgcactcccg cctgccaagt gaaactcggg 2580
gaggagttag tgcgggacat cccgtgtaat ggctacataa ctaccggct ttgcttcgac 2640
agtggcctg gctcgaccct cacacaacac tgcagatagg gggcgcaatt gggatcgta 2700
gaaaactatg gccgagcatg gggggggctc cggccccccc aaccccccg gtgggggggc 2760
caaggcccc cctacacccc cccatggggg gctgcccggc cccaaacccc ccgctcgga 2820
tggggggggc tgcgcccccc ccaaaccccc cttgcccggg gctgtgcccc ggaccccc 2878

```

&lt;210&gt; SEQ ID NO 10

&lt;211&gt; LENGTH: 2875

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Torque teno virus

&lt;400&gt; SEQUENCE: 10

```

tacacttcgg ggttcaggag gctcaatttg gctcgttcg ctcgaccac gtttgetgcc 60

```

-continued

---

aagcggacct	gattgaagac	tgacaaccgt	tacattcaaa	tttgaaaatg	gcgcccbaac	120
atggcggcgg	ggggcggagt	ttatgcaaat	taatttatgc	aaagtaggag	gagctccatt	180
ttaatttatg	caaagtagga	ggagtcatt	ctgattggtc	gggagctcaa	gtcctcattt	240
gcataggggt	taaccaatca	gatttaaggc	gttccatta	aagcgaatat	aagtaagtga	300
ggttccgaat	ggctgagttt	atgccgccag	cggtagacag	aactgtctag	cgactgggcg	360
ggtgcccggag	gatccctgat	ccggagtcaa	ggggcctatc	gggcaggagc	agctgagcgg	420
agggcctatg	ccggaacact	gggaggaagc	ctggttggaa	gctaccaagg	gctggcacga	480
tctcgactgc	cgctgcggta	actggcagga	ccacctatgg	ctcctactcg	ccgatggaga	540
cgccgctttg	gccccgccc	tagacgctat	agaaagagac	gctatgggtg	gagaagacgt	600
tactaccgct	acagaccgcg	ttactatagg	agacgatggc	tggttaaggag	aaggcggcgt	660
tccgtctacc	gtagaggtgg	acgtagagcg	cgcccctacc	gaataagtgc	ttttaacca	720
aaagtaatgc	ggaggggtgt	gatttagaggt	tggtggccaa	tattacagtg	tctaaaagga	780
caggaatcac	taagatatag	accactgcag	tgggacactg	aaaaacagtg	gagagtaaa	840
aaagactatg	aggacaacta	cggtactctg	gtgcagtacg	gaggaggttg	ggggagtgg	900
gaagtgacat	tggagggtt	atatcaggaa	cacttactct	ggagaaactc	ttggtcaaag	960
ggaaatgatg	gcatggacct	agtgagatac	tttggtgca	tagtataact	gtaccactg	1020
caggaccaag	actactggtt	ttggtgggat	acagacttta	aagaactata	cgagagagc	1080
atcaaagaat	actcccagcc	aagtgttatg	atgatggcca	aacgcactag	actagtaata	1140
gctagagaca	gagcaccaca	cagaagaaga	gtaagaaaa	tttccatacc	cccgccaaagc	1200
agagacacca	cacaatggca	atttcagaca	gacttctgca	aaagccact	attcacatgg	1260
gcggcaggat	taatagacat	gcagaaacca	ttgatgcaa	acggagcgtt	tagaaacgcc	1320
tggtggctag	aaacaaggaa	tgaccaggga	gaaatgaaat	acattgaaat	atggggaagg	1380
gtgccaccac	agggtagcac	agaactgcc	aaacagagtg	agtttaagaa	gggagataat	1440
aaccctaact	ataacataac	ggaaggacat	gaaaaaata	tttaccat	aatcatatac	1500
gttgaccaga	aagaccagaa	aacaagaaaa	aaatactgtg	tatgctacaa	caaaacttta	1560
aatagatgga	gaaaagccca	ggcagtagca	ttagcaatag	gagatcttca	aggactagta	1620
ctgctcagc	ttatgaatca	ggagatgaca	tactactgga	aatcgggaga	gttttctca	1680
ccattcctgc	aaagatggaa	aggaactagg	ctaataacca	tagacgcaag	aaaggcagac	1740
acagaaaacc	caaaagtaag	ttcgtgggaa	tgggggcaaa	actggaacac	aagcggaaaca	1800
gtgctacagg	aggtattcaa	catttcactg	aacaacactc	aaataagaca	ggatgacttt	1860
gcaaaattga	cactgccaaa	gtcaccacat	gacatagact	ttggacatca	cagcagattt	1920
ggaccattct	gtgttaaaaa	cgaaccacta	gaattccaac	tactgcctcc	aacaccaact	1980
aacctatgg	ttcagtacaa	atttctcttt	cagtttggcg	gtgaatacca	gccaccaaca	2040
ggtatccgcg	atccctgcat	tgatacacca	gcctatcctg	tgccgcagtc	aggaagtgtt	2100
acacacccca	aattcgccgg	aaagggcgga	atgctcacgg	aaacagaccg	ttggggtatc	2160
actgtgcct	cttcagaac	cctcagtgc	gatacaccca	ccgaagcagc	gcaaagtgca	2220
cttctcagag	gggacgcgga	aaagaaagga	gaggaaccg	aggaaccgc	gtcatcgtcc	2280
agtatcacga	gtgccgaaag	ctcactgag	ggagatggat	cgtctgatga	tgaagagaca	2340
atcagacgca	gaaggaggac	ctggaagcga	ctcagacgga	tggtcagaca	gcagcttgac	2400

-continued

---

```

cgacgaatgg accacaagcg acagcgactt cattgatacc cccataagag aaagatgcct 2460
caataaaaaa caaaaaaac gctaaacagt gtccgctat tagtgggggg gtccgggggg 2520
gcttgcccc cggtaagcgg ggttacgca ctaactcct gccaaagtga actcggggac 2580
gagtgagtgc gggacatccc gtgtaatgc tacataacta cccggcttg ctccgacagt 2640
ggccgtggct cgaccctcgc acaacactgc aggtaggggg cgcaattggg atcgttagaa 2700
aactatggcc gagcatgggg ggggctccgc ccccccaac cccccggtg ggggggcca 2760
ggccctcct acaccccc atggggggct gccgcccc aaaccccc cgtcggatgg 2820
ggggggctgc ccccccca aacccccct gccgggggt gtgccccga cccc 2875

```

&lt;210&gt; SEQ ID NO 11

&lt;211&gt; LENGTH: 2750

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Torque teno virus

&lt;400&gt; SEQUENCE: 11

```

taatgacagg gttcaccgga aaggctgcaa aattacagct aaaaccacaa tcataacaca 60
ataaaccaca aactattaca ggaactgca ataaattaag aaataaatta cacataacca 120
cctaaccaca ggaactttg caaaaaggg gaaataaatc tcattggctg ggcagaagt 180
cctcattaga ataagaaaag aaccaatcag aaacacttc tcttttagag tatataagta 240
agtgcgcaga cgaatggctg agtttatgcc gctggtgta gacacgaaca gagctgagt 300
tctaaccgcc tgggggggtg ccggagctcc tgagagcggg gtcaaggggc ctatcgggca 360
ggcggtaatc cagcggaaact gggccccct ccatggaaga aagatggctg acggtagcgt 420
actgcgcgca cggattatc tgcgactgta aaaaccgaa aaaacatctt gaaaaatgcc 480
ttacagacgc tatcgccgac gccgaaggag accgacacgg agatggaggc accggagggtg 540
gagacgctac tttcgatata ggtatcgacg cgctcctcgc gcccgccgc caaaggtaa 600
gagacggagg aggaaagctc cggtcattca atggttcct cctagccgga gaacctgcct 660
catagagggc ttctggcgtg tgagctacgg acaactggtc cgtacctgct tccctatgag 720
aaggctaaac ggactgattt tcacgggtgg aggatgtgac tggactcaat ggagtttaca 780
aaatttatc catgaaaaat taaactggag aaatatatgg acagcttcta atgtaggcat 840
ggagtttct agattttta gaggaaaatt ttacttctc agacaccctt ggagaagcta 900
tatagtaaca tgggaccaag acataccctg taaaccgctc ccatatcaaa acttacaacc 960
tctattaatg ctctcaaaa aacagcataa attagtcctc tctcaaaaag attgcaacc 1020
gaacagaaaa caaaaaccg ttacattaaa attcaggcct ccacaaaaat taacatcaca 1080
gtggagacta agcagagaac tctcaaaaat acccttaata agactaggaa taagtctcat 1140
agacctgca gaacctgggt tagaaggctg gggaaatgct ttttacagtg tactaggata 1200
tgaagctagt aaacacagtg gcagatggtc caactggaca caaatgaaat atttttggat 1260
ctatgacaca ggcgtgggaa acgcagtcta cgttatttta ctgaaaaaag acgtgagtga 1320
caatccagga gacatggcta cacagttgt aacaggctca ggacaacacc cagacgcaat 1380
agatcatata gaaatggtaa acgaaggatg gccttactgg ctattttttt atggacaatc 1440
agaacaagat ataaaaaac tagcacatga ccaagatata gtcagagaat atgccagaga 1500
ccctaaatca aaaaaattaa aaataggagt cataggatgg gccagcagta actacacaac 1560
agcagggagc aacaaaaaca gtgtacttca aacccagaa gcaatacaag gtggatatgt 1620
agcttatgca ggatccagaa taccaggcgc aggatctatc acaaatttat ttcaaatggg 1680

```

-continued

---

```

atggccagga gatcaaaact ggccaccac aaaccaagac caaaccaatt ttaactgggg 1740
actcagagga ctttgtgtat taagagataa catgaaacta ggagcacaag agctagacga 1800
tgaatgcaca atgctctcct tatttggacc atttgttgaa aaagcaaaca cagcttttgc 1860
tacaacgcac ccaaaatatt ttaggcctga actaaaggac tacaacgtag taatgaaata 1920
tgcttttaaa tttcagtggg gaggacatgg caccgaaaga tttaaaacaa ccatcggaga 1980
tcccagcacc ataccatgtc cctttgaacc cggggaacgg taccaccacg gggtaacaaga 2040
ccccgccaa gtaaaaaaca cagtctcaa cccttgggac tatgactgtg acgggattgt 2100
tagaacagat actctcaaaa gacttctcga actccccaca gagacggagg agacggagaa 2160
ggcgtacca ctccttgac aaaaaacaga gaaagagcca ttatcagact ccgacgaaga 2220
gagcgttata tcaagcaga gcagtggatc ctctcaagaa gaagagacgc agagacgaaa 2280
gcaccacaag ccaagcaagc gacgactcct caagcacctc cagcgggtgg taaagaggat 2340
gaaaacactg tgatagataa atacagaaac ctacagacc cctcactcaa tgtcacagga 2400
cacatgaaa aattcatgca actacacata caaacatac aagaaataag agctaaaaat 2460
gctaaaaaat ccctcaataa actttacttt tctgattaat agcggcctcc tgtgtccaat 2520
ctatttttcc tacaccctt caaaatggcg ggaggaacac aaaaatggcgg agggactaag 2580
gggggggcaa gcccccccc ggggggtgag ggggggttcc cccccctcc cccgggtcag 2640
ggggcggagc ccccgacccc cccatgcggg ggctccgccc cctgcacccc cgggaggggg 2700
ggaaaccccc cctcaacccc ccgcgggggg caagcccccc tgcaccccc 2750

```

&lt;210&gt; SEQ ID NO 12

&lt;211&gt; LENGTH: 2803

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Torque teno virus

&lt;400&gt; SEQUENCE: 12

```

tcatgacagg gttcaccgga agggctgcaa aattacagct aaaaccacaa gtctaacaca 60
ataaaccaca aagtattaca ggaactgca ataaatttag aaatagatta cacataacca 120
ccaaaccaca ggaacctac acataaccac caaacacag gaaacataac caccaaacca 180
caggaaactg tgcaaaaaag gggaaataaa ttctattggc tgggcctgaa gtctcatta 240
gaataataaa agaaccaatc agaagaactt cctcttttag agtatataag taagtgcgca 300
gacgaatggc tgagtttatg ccgctgggtg tagacacgaa cagagctgag tgtctaaccg 360
cctggcgggg tgccggagct cctgagagcg gactcaaggg gcttatcggg caggcggtaa 420
tccagcggaa cggggcccc ctcgatggaa gaaagatggc tgacggtagc gtactgcgcc 480
cacggattat tctgcggatg taaagacccc aaaaaacacc ttgaaaaatg ccttacagac 540
gctatcgcag acgccgaagg agaccgacac ggagatggag gcaccggagg tgagacgct 600
tctttcgata tcggtatcga cgcgctcctc gccgcgccc cacaaggta aggagacgga 660
ggagaaaagc tccggtcata caatggttcc ctctagccg gaggacctgc ctcatagagg 720
gcttctggcc gttgagctac ggacactggg tccgtacctg tctccctatg agaaggctga 780
acggactcat tttcacgggt ggcggttggt actggacaca gtggagtta caaaacttat 840
accatgaaaa acttaactgg agaaatatat ggacagcttc taatggtggc atggaatttg 900
ctagattttt aagaggaaaa ttttacttct tcagaccccc ctggagaagc tatattatta 960
ctgggacca agacattcct tgcaaacctt taccatacca aaacttacat ccactactta 1020

```

-continued

---

```

tgctatataa aaaacaacat aaacttgtag tatctcaaaa agactgtaat ccaaacagaa 1080
gacaaaaacc agtaacttta aaaataagac ctocacccaa attaacatca cagtggagat 1140
taagcagaga actagcaaaa atgccacttg tcagactagg agtcagtcta atagacctct 1200
cagaacctag gttagaagge tggggaaatg ctttttacag cgtactggga tatgaagcta 1260
gtaaaccactc agggagatgg tcaactgga cacaaataa atacttctgg atatatgaca 1320
caggagttagg aatgcagtt tatgtcattt tattaaaaa agaggtggat gataatccag 1380
gggcaatggc aacaaaatth gtaactggac caggacaaca cccagatgcc atagacagga 1440
tcgaacaaat aatgaagga tggccttact ggcttttctt ttacggacag tcagaacaag 1500
acataaaaaa attagcacac gatcaagaaa tagcaagggg atatgcaaac aatccaaaat 1560
ctaaaaaatt aaaaatagga gtgataggat gggctagcag taactttaca acagcaggca 1620
gctcacaaaa tcaaacacca caaacaccag aagccataca agggagatac gtagcatatg 1680
caggctcaaa aatacaagga gcaggagcaa ttacaaaactt atacacagat gcatggccgg 1740
gagacccaaa ttggccacct ctaaataagag aacaaacaaa ctttaactgg ggcttaagag 1800
gactctgtat aatgagagat aatatgaaac tgggagctca agaactagat gatgaatgta 1860
caatgctcac actttttgga ctttttggg aaaaagcaaa cacagctttt gctacaaatg 1920
accctaaata cttcagacca gaactcaaag actataacat agtaatgaaa tatgccttta 1980
aatttcagtg gggaggccac ggaaccgaaa gattcaaac aaccatcgga gatcccagca 2040
ccataccatg tcctttgaa cccggggaac ggtaccacca cggggtacaa gaccccgcca 2100
aggtacaaaa cacagtcctc aaccctggg actatgactg tgacgggatt gttagaacag 2160
atactctcaa aagacttctc gaactcccca cagagacgga ggagacggag aaggcgtacc 2220
cactccttgg acaaaaaaca gagaaagagc cattatcaga ctccgacgaa gagagcgta 2280
tctcaagcac gagcagtgga tcctctcaag aagaagagac gcagagaaga agacagcaca 2340
agccaagcaa gcgacgactc ctcaagcacc tccagcgggt ggtaaagaga atgaagacac 2400
tgtgatagat aaatatagaa acctagcaga cccctcactc aatgtcacag gacacatgga 2460
aaaattcatg caactgcaca tacaaaacgt acaagaaata agagctaaaa atgctaaaaa 2520
atccctcaat aaactttact tttctgatta ataccggcct cctgtgtcca atctattht 2580
cctacacccc ttcaaatgg cgggcgggac acaaaatggc ggaggaaact aagggggggg 2640
caagcccccc cccgggggtt gaggggggtt ttccccct cccccggtg cagggggcgg 2700
agcccccgca cccccctgc gggggtccg cccctgcac ccccgaggg gggggaaacc 2760
ccccctcaac cccccgagg gggcaagccc cctgcaccc ccc 2803

```

&lt;210&gt; SEQ ID NO 13

&lt;211&gt; LENGTH: 635

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Torque teno virus

&lt;400&gt; SEQUENCE: 13

```

Met Arg Phe Arg Arg Arg Arg Phe Gly Arg Arg Arg Arg Tyr Tyr Arg
1           5           10           15

```

```

Lys Arg Arg Gly Gly Trp Arg Arg Arg Phe Arg Ile Arg Arg Arg Arg
20           25           30

```

```

Pro Trp Arg Arg Trp Arg Val Arg Arg Trp Arg Arg Ser Val Phe Arg
35           40           45

```

```

Arg Gly Gly Arg Arg Ala Arg Pro Tyr Arg Ile Ser Ala Trp Asn Pro
50           55           60

```

-continued

---

Lys Val Leu Arg Asn Cys Arg Ile Thr Gly Trp Trp Pro Val Ile Gln  
 65 70 75 80  
 Cys Met Asp Gly Met Glu Trp Ile Lys Tyr Lys Pro Met Asp Leu Arg  
 85 90 95  
 Val Glu Ala Asn Trp Ile Phe Asn Lys Gln Asp Ser Lys Ile Glu Thr  
 100 105 110  
 Glu Gln Met Gly Tyr Leu Met Gln Tyr Gly Gly Gly Trp Ser Ser Gly  
 115 120 125  
 Val Ile Ser Leu Glu Gly Leu Phe Asn Glu Asn Arg Leu Trp Arg Asn  
 130 135 140  
 Ile Trp Ser Lys Ser Asn Asp Gly Met Asp Leu Val Arg Tyr Phe Gly  
 145 150 155 160  
 Cys Arg Ile Arg Leu Tyr Pro Thr Glu Asn Gln Asp Tyr Leu Phe Trp  
 165 170 175  
 Tyr Asp Thr Glu Phe Asp Glu Gln Gln Arg Arg Met Leu Asp Glu Tyr  
 180 185 190  
 Thr Gln Pro Ser Val Met Leu Gln Ala Lys Asn Ser Arg Leu Ile Val  
 195 200 205  
 Cys Lys Gln Lys Met Pro Ile Arg Arg Arg Val Lys Ser Ile Phe Ile  
 210 215 220  
 Pro Pro Pro Ala Gln Leu Thr Thr Gln Trp Lys Phe Gln Gln Glu Leu  
 225 230 235 240  
 Cys Gln Phe Pro Leu Phe Asn Trp Ala Cys Ile Cys Ile Asp Met Asp  
 245 250 255  
 Thr Pro Phe Asp Tyr Asn Gly Ala Trp Arg Asn Ala Trp Trp Leu Met  
 260 265 270  
 Arg Arg Leu Gln Asn Gly Asn Met Glu Tyr Ile Glu Arg Trp Gly Arg  
 275 280 285  
 Ile Pro Met Thr Gly Asp Thr Glu Leu Pro Pro Ala Asp Asp Phe Lys  
 290 295 300  
 Ala Gly Gly Val Asn Lys Asn Phe Lys Pro Thr Gly Ile Gln Arg Ile  
 305 310 315 320  
 Tyr Pro Ile Val Ala Val Cys Leu Val Glu Gly Asn Lys Arg Val Val  
 325 330 335  
 Lys Trp Ala Thr Val His Asn Gly Pro Ile Asp Arg Trp Arg Lys Lys  
 340 345 350  
 Gln Thr Gly Thr Leu Lys Leu Ser Asn Leu Arg Gly Leu Val Leu Arg  
 355 360 365  
 Val Cys Ser Glu Ser Glu Thr Tyr Tyr Lys Trp Thr Gly Ser Glu Phe  
 370 375 380  
 Thr Gly Ala Phe Gln Gln Asp Trp Trp Pro Val Gly Gly Thr Glu Tyr  
 385 390 395 400  
 Pro Leu Cys Thr Ile Lys Met Asp Pro Glu Tyr Glu Asn Pro Thr Val  
 405 410 415  
 Glu Val Trp Ser Trp Lys Ala Asn Ile Pro Thr Ser Gly Thr Leu Lys  
 420 425 430  
 Asp Tyr Phe Gly Leu Ser Thr Gly Gln Gln Trp Lys Asp Thr Asp Phe  
 435 440 445  
 Ala Arg Leu Gln Leu Pro Arg Ser Ser His Asn Val Asp Phe Gly His  
 450 455 460  
 Lys Ala Arg Phe Gly Pro Phe Cys Val Lys Lys Pro Pro Val Glu Phe  
 465 470 475 480

-continued

---

```

Arg Asp Thr Ala Pro Asn Pro Leu Asn Ile Trp Val Lys Tyr Thr Phe
      485                               490                               495
Tyr Phe Gln Phe Gly Gly Met Tyr Gln Pro Pro Thr Gly Ile Gln Asp
      500                               505                               510
Pro Cys Thr Ser Asn Pro Thr Tyr Pro Val Arg Met Val Gly Ala Val
      515                               520                               525
Thr His Pro Lys Tyr Ala Gly Gln Gly Gly Ile Thr Thr Gln Ile Gly
      530                               535                               540
Asp Gln Gly Ile Thr Ala Ala Ser Ile Arg Ala Ile Ser Ala Ala Pro
      545                               550                               555                               560
Pro Asp Thr Tyr Thr Gln Ser Ala Phe Leu Lys Ala Pro Glu Thr Glu
      565                               570                               575
Lys Glu Glu Glu Arg Glu Ser Glu Thr Ser Phe Thr Ser Ala Glu Ser
      580                               585                               590
Ser Ser Glu Gly Asp Gly Ser Ser Asp Asp Gln Ala Glu Arg Arg Ala
      595                               600                               605
Ala Arg Lys Arg Val Ile Lys Leu Leu Leu Lys Arg Leu Ala Asp Arg
      610                               615                               620
Pro Val Asp Asn Lys Arg Arg Arg Phe Ser Glu
      625                               630                               635

```

```

<210> SEQ ID NO 14
<211> LENGTH: 639
<212> TYPE: PRT
<213> ORGANISM: Torque teno virus

```

```

<400> SEQUENCE: 14

```

```

Met Ala Pro Thr Arg Arg Trp Arg Arg Arg Phe Gly Arg Arg Arg Arg
 1      5      10      15
Arg Tyr Arg Lys Arg Arg Tyr Gly Trp Arg Arg Arg Tyr Tyr Arg Tyr
 20     25     30
Arg Pro Arg Tyr Tyr Arg Arg Arg Trp Leu Val Arg Arg Arg Arg
 35     40     45
Ser Val Tyr Arg Arg Gly Gly Arg Arg Ala Arg Pro Tyr Arg Ile Ser
 50     55     60
Ala Phe Asn Pro Lys Val Met Arg Arg Val Val Ile Arg Gly Trp Trp
 65     70     75     80
Pro Ile Leu Gln Cys Leu Lys Gly Gln Glu Ser Leu Arg Tyr Arg Pro
 85     90     95
Leu Gln Trp Asp Thr Glu Lys Gln Trp Arg Val Lys Lys Asp Tyr Glu
100    105    110
Asp Asn Tyr Gly Tyr Leu Val Gln Tyr Gly Gly Gly Trp Gly Ser Gly
115    120    125
Glu Val Thr Leu Glu Gly Leu Tyr Gln Glu His Leu Leu Trp Arg Asn
130    135    140
Ser Trp Ser Lys Gly Asn Asp Gly Met Asp Leu Val Arg Tyr Phe Gly
145    150    155    160
Cys Ile Val Tyr Leu Tyr Pro Leu Gln Asp Gln Asp Tyr Trp Phe Trp
165    170    175
Trp Asp Thr Asp Phe Lys Glu Leu Tyr Ala Glu Ser Ile Lys Glu Tyr
180    185    190
Ser Gln Pro Ser Val Met Met Met Ala Lys Arg Thr Arg Leu Val Ile
195    200    205
Ala Arg Asp Arg Ala Pro His Arg Arg Arg Val Arg Lys Ile Phe Ile
210    215    220

```

-continued

---

Pro Pro Pro Ser Arg Asp Thr Thr Gln Trp Gln Phe Gln Thr Asp Phe  
 225 230 235 240  
 Cys Lys Arg Pro Leu Phe Thr Trp Ala Ala Gly Leu Ile Asp Met Gln  
 245 250 255  
 Lys Pro Phe Asp Ala Asn Gly Ala Phe Arg Asn Ala Trp Trp Leu Glu  
 260 265 270  
 Thr Arg Asn Asp Gln Gly Glu Met Lys Tyr Ile Glu Leu Trp Gly Arg  
 275 280 285  
 Val Pro Pro Gln Gly Asp Thr Glu Leu Pro Lys Gln Ser Glu Phe Lys  
 290 295 300  
 Lys Gly Asp Asn Asn Pro Asn Tyr Asn Ile Thr Glu Gly His Glu Lys  
 305 310 315 320  
 Asn Ile Tyr Pro Ile Ile Ile Tyr Val Asp Gln Lys Asp Gln Lys Thr  
 325 330 335  
 Arg Lys Lys Tyr Cys Val Cys Tyr Asn Lys Thr Leu Asn Arg Trp Arg  
 340 345 350  
 Lys Ala Gln Ala Ser Thr Leu Ala Ile Gly Asp Leu Gln Gly Leu Val  
 355 360 365  
 Leu Arg Gln Leu Met Asn Gln Glu Met Thr Tyr Tyr Trp Lys Ser Gly  
 370 375 380  
 Glu Phe Ser Ser Pro Phe Leu Gln Arg Trp Lys Gly Thr Arg Leu Ile  
 385 390 395 400  
 Thr Ile Asp Ala Arg Lys Ala Asp Thr Glu Asn Pro Lys Val Ser Ser  
 405 410 415  
 Trp Glu Trp Gly Gln Asn Trp Asn Thr Ser Gly Thr Val Leu Gln Glu  
 420 425 430  
 Val Phe Asn Ile Ser Leu Asn Asn Thr Gln Ile Arg Gln Asp Asp Phe  
 435 440 445  
 Ala Lys Leu Thr Leu Pro Lys Ser Pro His Asp Ile Asp Phe Gly His  
 450 455 460  
 His Ser Arg Phe Gly Pro Phe Cys Val Lys Asn Glu Pro Leu Glu Phe  
 465 470 475 480  
 Gln Leu Leu Pro Pro Thr Pro Thr Asn Leu Trp Phe Gln Tyr Lys Phe  
 485 490 495  
 Leu Phe Gln Phe Gly Gly Glu Tyr Gln Pro Pro Thr Gly Ile Arg Asp  
 500 505 510  
 Pro Cys Ile Asp Thr Pro Ala Tyr Pro Val Pro Gln Ser Gly Ser Val  
 515 520 525  
 Thr His Pro Lys Phe Ala Gly Lys Gly Gly Met Leu Thr Glu Thr Asp  
 530 535 540  
 Arg Trp Gly Ile Thr Ala Ala Ser Ser Arg Thr Leu Ser Ala Asp Thr  
 545 550 555 560  
 Pro Thr Glu Ala Ala Gln Ser Ala Leu Leu Arg Gly Asp Ala Glu Lys  
 565 570 575  
 Lys Gly Glu Glu Thr Glu Glu Thr Ala Ser Ser Ser Ser Ile Thr Ser  
 580 585 590  
 Ala Glu Ser Ser Thr Glu Gly Asp Gly Ser Ser Asp Asp Glu Glu Thr  
 595 600 605  
 Ile Arg Arg Arg Arg Arg Thr Trp Lys Arg Leu Arg Arg Met Val Arg  
 610 615 620  
 Gln Gln Leu Asp Arg Arg Met Asp His Lys Arg Gln Arg Leu His  
 625 630 635

-continued

---

```

<210> SEQ ID NO 15
<211> LENGTH: 625
<212> TYPE: PRT
<213> ORGANISM: Torque teno virus

<400> SEQUENCE: 15

Met Pro Tyr Arg Arg Tyr Arg Arg Arg Arg Arg Arg Pro Thr Arg Arg
1          5          10          15

Trp Arg His Arg Arg Trp Arg Arg Tyr Phe Arg Tyr Arg Tyr Arg Arg
20          25          30

Ala Pro Arg Arg Arg Arg Pro Lys Val Arg Arg Arg Arg Arg Lys Ala
35          40          45

Pro Val Ile Gln Trp Phe Pro Pro Ser Arg Arg Thr Cys Leu Ile Glu
50          55          60

Gly Phe Trp Pro Leu Ser Tyr Gly His Trp Phe Arg Thr Cys Leu Pro
65          70          75          80

Met Arg Arg Leu Asn Gly Leu Ile Phe Thr Gly Gly Gly Cys Asp Trp
85          90          95

Thr Gln Trp Ser Leu Gln Asn Leu Phe His Glu Lys Leu Asn Trp Arg
100         105         110

Asn Ile Trp Thr Ala Ser Asn Val Gly Met Glu Phe Ala Arg Phe Leu
115         120         125

Arg Gly Lys Phe Tyr Phe Phe Arg His Pro Trp Arg Ser Tyr Ile Val
130         135         140

Thr Trp Asp Gln Asp Ile Pro Cys Lys Pro Leu Pro Tyr Gln Asn Leu
145         150         155         160

Gln Pro Leu Leu Met Leu Leu Lys Lys Gln His Lys Leu Val Leu Ser
165         170         175

Gln Lys Asp Cys Asn Pro Asn Arg Lys Gln Lys Pro Val Thr Leu Lys
180         185         190

Phe Arg Pro Pro Pro Lys Leu Thr Ser Gln Trp Arg Leu Ser Arg Glu
195         200         205

Leu Ser Lys Ile Pro Leu Ile Arg Leu Gly Ile Ser Leu Ile Asp Leu
210         215         220

Ser Glu Pro Trp Leu Glu Gly Trp Gly Asn Ala Phe Tyr Ser Val Leu
225         230         235         240

Gly Tyr Glu Ala Ser Lys His Ser Gly Arg Trp Ser Asn Trp Thr Gln
245         250         255

Met Lys Tyr Phe Trp Ile Tyr Asp Thr Gly Val Gly Asn Ala Val Tyr
260         265         270

Val Ile Leu Leu Lys Lys Asp Val Ser Asp Asn Pro Gly Asp Met Ala
275         280         285

Thr Gln Phe Val Thr Gly Ser Gly Gln His Pro Asp Ala Ile Asp His
290         295         300

Ile Glu Met Val Asn Glu Gly Trp Pro Tyr Trp Leu Phe Phe Tyr Gly
305         310         315         320

Gln Ser Glu Gln Asp Ile Lys Lys Leu Ala His Asp Gln Asp Ile Val
325         330         335

Arg Glu Tyr Ala Arg Asp Pro Lys Ser Lys Lys Leu Lys Ile Gly Val
340         345         350

Ile Gly Trp Ala Ser Ser Asn Tyr Thr Thr Ala Gly Ser Asn Gln Asn
355         360         365

Ser Val Leu Gln Thr Pro Glu Ala Ile Gln Gly Gly Tyr Val Ala Tyr
370         375         380

```

-continued

Ala Gly Ser Arg Ile Pro Gly Ala Gly Ser Ile Thr Asn Leu Phe Gln  
 385 390 395 400

Met Gly Trp Pro Gly Asp Gln Asn Trp Pro Pro Thr Asn Gln Asp Gln  
 405 410 415

Thr Asn Phe Asn Trp Gly Leu Arg Gly Leu Cys Val Leu Arg Asp Asn  
 420 425 430

Met Lys Leu Gly Ala Gln Glu Leu Asp Asp Glu Cys Thr Met Leu Ser  
 435 440 445

Leu Phe Gly Pro Phe Val Glu Lys Ala Asn Thr Ala Phe Ala Thr Asn  
 450 455 460

Asp Pro Lys Tyr Phe Arg Pro Glu Leu Lys Asp Tyr Asn Val Val Met  
 465 470 475 480

Lys Tyr Ala Phe Lys Phe Gln Trp Gly Gly His Gly Thr Glu Arg Phe  
 485 490 495

Lys Thr Thr Ile Gly Asp Pro Ser Thr Ile Pro Cys Pro Phe Glu Pro  
 500 505 510

Gly Glu Arg Tyr His His Gly Val Gln Asp Pro Ala Lys Val Gln Asn  
 515 520 525

Thr Val Leu Asn Pro Trp Asp Tyr Asp Cys Asp Gly Ile Val Arg Thr  
 530 535 540

Asp Thr Leu Lys Arg Leu Leu Glu Leu Pro Thr Glu Thr Glu Glu Thr  
 545 550 555 560

Glu Lys Ala Tyr Pro Leu Leu Gly Gln Lys Thr Glu Lys Glu Pro Leu  
 565 570 575

Ser Asp Ser Asp Glu Glu Ser Val Ile Ser Ser Thr Ser Ser Gly Ser  
 580 585 590

Ser Gln Glu Glu Glu Thr Gln Arg Arg Lys His His Lys Pro Ser Lys  
 595 600 605

Arg Arg Leu Leu Lys His Leu Gln Arg Val Val Lys Arg Met Lys Thr  
 610 615 620

Leu  
 625

&lt;210&gt; SEQ ID NO 16

&lt;211&gt; LENGTH: 625

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Torque teno virus

&lt;400&gt; SEQUENCE: 16

Met Pro Tyr Arg Arg Tyr Arg Arg Arg Arg Arg Pro Thr Arg Arg  
 1 5 10 15

Trp Arg His Arg Arg Trp Arg Arg Phe Phe Arg Tyr Arg Tyr Arg Arg  
 20 25 30

Ala Pro Arg Arg Arg Arg Thr Lys Val Arg Arg Arg Arg Lys Ala  
 35 40 45

Pro Val Ile Gln Trp Phe Pro Pro Ser Arg Arg Thr Cys Leu Ile Glu  
 50 55 60

Gly Phe Trp Pro Leu Ser Tyr Gly His Trp Phe Arg Thr Cys Leu Pro  
 65 70 75 80

Met Arg Arg Leu Asn Gly Leu Ile Phe Thr Gly Gly Gly Cys Asp Trp  
 85 90 95

Thr Gln Trp Ser Leu Gln Asn Leu Tyr His Glu Lys Leu Asn Trp Arg  
 100 105 110

Asn Ile Trp Thr Ala Ser Asn Val Gly Met Glu Phe Ala Arg Phe Leu

-continued

115				120				125							
Arg	Gly	Lys	Phe	Tyr	Phe	Phe	Arg	His	Pro	Trp	Arg	Ser	Tyr	Ile	Ile
	130					135					140				
Thr	Trp	Asp	Gln	Asp	Ile	Pro	Cys	Lys	Pro	Leu	Pro	Tyr	Gln	Asn	Leu
	145				150					155					160
His	Pro	Leu	Leu	Met	Leu	Leu	Lys	Lys	Gln	His	Lys	Leu	Val	Leu	Ser
				165					170						175
Gln	Lys	Asp	Cys	Asn	Pro	Asn	Arg	Arg	Gln	Lys	Pro	Val	Thr	Leu	Lys
		180						185					190		
Ile	Arg	Pro	Pro	Pro	Lys	Leu	Thr	Ser	Gln	Trp	Arg	Leu	Ser	Arg	Glu
		195					200					205			
Leu	Ala	Lys	Met	Pro	Leu	Val	Arg	Leu	Gly	Val	Ser	Leu	Ile	Asp	Leu
	210				215						220				
Ser	Glu	Pro	Trp	Leu	Glu	Gly	Trp	Gly	Asn	Ala	Phe	Tyr	Ser	Val	Leu
	225				230					235					240
Gly	Tyr	Glu	Ala	Ser	Lys	His	Ser	Gly	Arg	Trp	Ser	Asn	Trp	Thr	Gln
				245					250					255	
Ile	Lys	Tyr	Phe	Trp	Ile	Tyr	Asp	Thr	Gly	Val	Gly	Asn	Ala	Val	Tyr
			260					265						270	
Val	Ile	Leu	Leu	Lys	Gln	Glu	Val	Asp	Asp	Asn	Pro	Gly	Ala	Met	Ala
		275					280					285			
Thr	Lys	Phe	Val	Thr	Gly	Pro	Gly	Gln	His	Pro	Asp	Ala	Ile	Asp	Arg
	290					295					300				
Ile	Glu	Gln	Ile	Asn	Glu	Gly	Trp	Pro	Tyr	Trp	Leu	Phe	Phe	Tyr	Gly
	305				310					315					320
Gln	Ser	Glu	Gln	Asp	Ile	Lys	Lys	Leu	Ala	His	Asp	Gln	Glu	Ile	Ala
				325					330					335	
Arg	Glu	Tyr	Ala	Asn	Asn	Pro	Lys	Ser	Lys	Lys	Leu	Lys	Ile	Gly	Val
			340					345					350		
Ile	Gly	Trp	Ala	Ser	Ser	Asn	Phe	Thr	Thr	Ala	Gly	Ser	Ser	Gln	Asn
		355					360					365			
Gln	Thr	Pro	Gln	Thr	Pro	Glu	Ala	Ile	Gln	Gly	Gly	Tyr	Val	Ala	Tyr
	370					375					380				
Ala	Gly	Ser	Lys	Ile	Gln	Gly	Ala	Gly	Ala	Ile	Thr	Asn	Leu	Tyr	Thr
	385				390					395					400
Asp	Ala	Trp	Pro	Gly	Asp	Gln	Asn	Trp	Pro	Pro	Leu	Asn	Arg	Glu	Gln
				405					410					415	
Thr	Asn	Phe	Asn	Trp	Gly	Leu	Arg	Gly	Leu	Cys	Ile	Met	Arg	Asp	Asn
			420					425					430		
Met	Lys	Leu	Gly	Ala	Gln	Glu	Leu	Asp	Asp	Glu	Cys	Thr	Met	Leu	Thr
		435					440					445			
Leu	Phe	Gly	Pro	Phe	Val	Glu	Lys	Ala	Asn	Thr	Ala	Phe	Ala	Thr	Asn
	450					455					460				
Asp	Pro	Lys	Tyr	Phe	Arg	Pro	Glu	Leu	Lys	Asp	Tyr	Asn	Ile	Val	Met
	465				470					475					480
Lys	Tyr	Ala	Phe	Lys	Phe	Gln	Trp	Gly	Gly	His	Gly	Thr	Glu	Arg	Phe
				485					490					495	
Lys	Thr	Thr	Ile	Gly	Asp	Pro	Ser	Thr	Ile	Pro	Cys	Pro	Phe	Glu	Pro
			500						505				510		
Gly	Glu	Arg	Tyr	His	His	Gly	Val	Gln	Asp	Pro	Ala	Lys	Val	Gln	Asn
		515					520					525			
Thr	Val	Leu	Asn	Pro	Trp	Asp	Tyr	Asp	Cys	Asp	Gly	Ile	Val	Arg	Thr
	530					535					540				

-continued

Asp Thr Leu Lys Arg Leu Leu Glu Leu Pro Thr Glu Thr Glu Glu Thr  
 545 550 555 560  
 Glu Lys Ala Tyr Pro Leu Leu Gly Gln Lys Thr Glu Lys Glu Pro Leu  
 565 570 575  
 Ser Asp Ser Asp Glu Glu Ser Val Ile Ser Ser Thr Ser Ser Gly Ser  
 580 585 590  
 Ser Gln Glu Glu Glu Thr Gln Arg Arg Arg Gln His Lys Pro Ser Lys  
 595 600 605  
 Arg Arg Leu Leu Lys His Leu Gln Arg Val Val Lys Arg Met Lys Thr  
 610 615 620  
 Leu  
 625

<210> SEQ ID NO 17  
 <211> LENGTH: 73  
 <212> TYPE: PRT  
 <213> ORGANISM: Torque teno virus

<400> SEQUENCE: 17

Met Lys Glu Lys Asp Tyr Trp Glu Glu Ala Trp Leu Thr Ser Cys Thr  
 1 5 10 15  
 Ser Ile His Asp His His Cys Asn Cys Gly Ser Trp Arg Asp His Leu  
 20 25 30  
 Trp Thr Leu Cys Ala Leu Asp Asp Ala Asp Leu Ala Ala Ala Ala Asp  
 35 40 45  
 Ile Ile Glu Arg Glu Glu Ala Asp Gly Gly Glu Asp Phe Gly Phe Val  
 50 55 60  
 Asp Gly Asp Pro Gly Asp Ala Gly Gly  
 65 70

<210> SEQ ID NO 18  
 <211> LENGTH: 72  
 <212> TYPE: PRT  
 <213> ORGANISM: Torque teno virus

<400> SEQUENCE: 18

Met Pro Glu His Trp Glu Glu Ala Trp Leu Glu Ala Thr Lys Gly Trp  
 1 5 10 15  
 His Asp Leu Asp Cys Arg Cys Gly Asn Trp Gln Asp His Leu Trp Leu  
 20 25 30  
 Leu Leu Ala Asp Gly Asp Ala Ala Leu Ala Ala Val Asp Ala Ile  
 35 40 45  
 Glu Arg Asp Ala Met Gly Gly Glu Asp Val Thr Thr Ala Thr Asp Arg  
 50 55 60  
 Val Thr Ile Gly Asp Asp Gly Trp  
 65 70

<210> SEQ ID NO 19  
 <211> LENGTH: 68  
 <212> TYPE: PRT  
 <213> ORGANISM: Torque teno virus

<400> SEQUENCE: 19

Met Glu Glu Arg Trp Leu Thr Val Ala Tyr Cys Ala His Gly Leu Phe  
 1 5 10 15  
 Cys Asp Cys Lys Asn Pro Lys Lys His Leu Glu Lys Cys Leu Thr Asp  
 20 25 30

-continued

---

Ala Ile Ala Asp Ala Glu Gly Asp Arg His Gly Asp Gly Gly Thr Gly  
                   35                  40                  45  
 Gly Gly Asp Ala Thr Phe Asp Ile Gly Ile Asp Ala Leu Leu Ala Ala  
           50                  55                  60  
 Ala Ala Gln Arg  
 65

<210> SEQ ID NO 20  
 <211> LENGTH: 68  
 <212> TYPE: PRT  
 <213> ORGANISM: Torque teno virus

<400> SEQUENCE: 20

Met Glu Glu Arg Trp Leu Thr Val Ala Tyr Cys Ala His Gly Leu Phe  
 1                  5                  10                  15  
 Cys Gly Cys Lys Asp Pro Lys Lys His Leu Glu Lys Cys Leu Thr Asp  
           20                  25                  30  
 Ala Ile Ala Asp Ala Glu Gly Asp Arg His Gly Asp Gly Gly Thr Gly  
           35                  40                  45  
 Gly Gly Asp Ala Ser Phe Asp Ile Gly Ile Asp Ala Leu Leu Ala Ala  
           50                  55                  60  
 Ala Ala Gln Arg  
 65

<210> SEQ ID NO 21  
 <211> LENGTH: 174  
 <212> TYPE: PRT  
 <213> ORGANISM: Torque teno virus

<400> SEQUENCE: 21

Met Arg Phe Arg Arg Arg Arg Phe Gly Arg Arg Arg Arg Tyr Tyr Arg  
 1                  5                  10                  15  
 Lys Arg Arg Gly Gly Trp Arg Arg Arg Phe Arg Ile Arg Arg Arg Arg  
           20                  25                  30  
 Pro Trp Arg Arg Trp Arg Phe Gly Gly Met Tyr Gln Pro Pro Thr Gly  
           35                  40                  45  
 Ile Gln Asp Pro Cys Thr Ser Asn Pro Thr Tyr Pro Val Arg Met Val  
           50                  55                  60  
 Gly Ala Val Thr His Pro Lys Tyr Ala Gly Gln Gly Gly Ile Thr Thr  
           65                  70                  75                  80  
 Gln Ile Gly Asp Gln Gly Ile Thr Ala Ala Ser Ile Arg Ala Ile Ser  
           85                  90                  95  
 Ala Ala Pro Pro Asp Thr Tyr Thr Gln Ser Ala Phe Leu Lys Ala Pro  
           100                  105                  110  
 Glu Thr Glu Lys Glu Glu Glu Arg Glu Ser Glu Thr Ser Phe Thr Ser  
           115                  120                  125  
 Ala Glu Ser Ser Ser Glu Gly Asp Gly Ser Ser Asp Asp Gln Ala Glu  
           130                  135                  140  
 Arg Arg Ala Ala Arg Lys Arg Val Ile Lys Leu Leu Leu Lys Arg Leu  
           145                  150                  155                  160  
 Ala Asp Arg Pro Val Asp Asn Lys Arg Arg Arg Phe Ser Glu  
                   165                  170

<210> SEQ ID NO 22  
 <211> LENGTH: 182  
 <212> TYPE: PRT  
 <213> ORGANISM: Torque teno virus

-continued

&lt;400&gt; SEQUENCE: 22

```

Met Ala Pro Thr Arg Arg Trp Arg Arg Arg Phe Gly Arg Arg Arg Arg
1          5          10          15
Arg Tyr Arg Lys Arg Arg Tyr Gly Trp Arg Arg Arg Tyr Tyr Arg Tyr
20          25          30
Arg Pro Arg Tyr Tyr Arg Arg Arg Trp Leu Phe Gly Gly Glu Tyr Gln
35          40          45
Pro Pro Thr Gly Ile Arg Asp Pro Cys Ile Asp Thr Pro Ala Tyr Pro
50          55          60
Val Pro Gln Ser Gly Ser Val Thr His Pro Lys Phe Ala Gly Lys Gly
65          70          75          80
Gly Met Leu Thr Glu Thr Asp Arg Trp Gly Ile Thr Ala Ala Ser Ser
85          90          95
Arg Thr Leu Ser Ala Asp Thr Pro Thr Glu Ala Ala Gln Ser Ala Leu
100         105         110
Leu Arg Gly Asp Ala Glu Lys Lys Gly Glu Glu Thr Glu Glu Thr Ala
115         120         125
Ser Ser Ser Ser Ile Thr Ser Ala Glu Ser Ser Thr Glu Gly Asp Gly
130         135         140
Ser Ser Asp Asp Glu Glu Thr Ile Arg Arg Arg Arg Arg Thr Trp Lys
145         150         155         160
Arg Leu Arg Arg Met Val Arg Gln Gln Leu Asp Arg Arg Met Asp His
165         170         175
Lys Arg Gln Arg Leu His
180

```

&lt;210&gt; SEQ ID NO 23

&lt;211&gt; LENGTH: 178

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Torque teno virus

&lt;400&gt; SEQUENCE: 23

```

Met Pro Tyr Arg Arg Tyr Arg Arg Arg Arg Arg Arg Pro Thr Arg Arg
1          5          10          15
Trp Arg His Arg Arg Trp Arg Arg Tyr Phe Arg Tyr Arg Tyr Arg Arg
20          25          30
Ala Pro Arg Arg Arg Arg Pro Lys Trp Gly Gly His Gly Thr Glu Arg
35          40          45
Phe Lys Thr Thr Ile Gly Asp Pro Ser Thr Ile Pro Cys Pro Phe Glu
50          55          60
Pro Gly Glu Arg Tyr His His Gly Val Gln Asp Pro Ala Lys Val Gln
65          70          75          80
Asn Thr Val Leu Asn Pro Trp Asp Tyr Asp Cys Asp Gly Ile Val Arg
85          90          95
Thr Asp Thr Leu Lys Arg Leu Leu Glu Leu Pro Thr Glu Thr Glu Glu
100         105         110
Thr Glu Lys Ala Tyr Pro Leu Leu Gly Gln Lys Thr Glu Lys Glu Pro
115         120         125
Leu Ser Asp Ser Asp Glu Glu Ser Val Ile Ser Ser Thr Ser Ser Gly
130         135         140
Ser Ser Gln Glu Glu Glu Thr Gln Arg Arg Lys His His Lys Pro Ser
145         150         155         160
Lys Arg Arg Leu Leu Lys His Leu Gln Arg Val Val Lys Arg Met Lys
165         170         175

```

-continued

Thr Leu

<210> SEQ ID NO 24  
 <211> LENGTH: 178  
 <212> TYPE: PRT  
 <213> ORGANISM: Torque teno virus

&lt;400&gt; SEQUENCE: 24

```

Met Pro Tyr Arg Arg Tyr Arg Arg Arg Arg Arg Arg Pro Thr Arg Arg
1           5           10           15
Trp Arg His Arg Arg Trp Arg Arg Phe Phe Arg Tyr Arg Tyr Arg Arg
                20           25           30
Ala Pro Arg Arg Arg Arg Thr Lys Trp Gly Gly His Gly Thr Glu Arg
                35           40           45
Phe Lys Thr Thr Ile Gly Asp Pro Ser Thr Ile Pro Cys Pro Phe Glu
                50           55           60
Pro Gly Glu Arg Tyr His His Gly Val Gln Asp Pro Ala Lys Val Gln
        65           70           75           80
Asn Thr Val Leu Asn Pro Trp Asp Tyr Asp Cys Asp Gly Ile Val Arg
                85           90           95
Thr Asp Thr Leu Lys Arg Leu Leu Glu Leu Pro Thr Glu Thr Glu Glu
                100          105          110
Thr Glu Lys Ala Tyr Pro Leu Leu Gly Gln Lys Thr Glu Lys Glu Pro
                115          120          125
Leu Ser Asp Ser Asp Glu Glu Ser Val Ile Ser Ser Thr Ser Ser Gly
                130          135          140
Ser Ser Gln Glu Glu Glu Thr Gln Arg Arg Arg Gln His Lys Pro Ser
        145          150          155          160
Lys Arg Arg Leu Leu Lys His Leu Gln Arg Val Val Lys Arg Met Lys
                165          170          175

```

Thr Leu

<210> SEQ ID NO 25  
 <211> LENGTH: 224  
 <212> TYPE: PRT  
 <213> ORGANISM: Torque teno virus

&lt;400&gt; SEQUENCE: 25

```

Met Lys Glu Lys Asp Tyr Trp Glu Glu Ala Trp Leu Thr Ser Cys Thr
1           5           10           15
Ser Ile His Asp His His Cys Asn Cys Gly Ser Trp Arg Asp His Leu
                20           25           30
Trp Thr Leu Cys Ala Leu Asp Asp Ala Asp Leu Ala Ala Ala Ala Asp
                35           40           45
Ile Ile Glu Arg Glu Glu Ala Asp Gly Gly Glu Asp Phe Gly Phe Val
        50           55           60
Asp Gly Asp Pro Gly Asp Ala Gly Gly Ser Ala Ala Cys Thr Ser Leu
        65           70           75           80
Pro Pro Glu Ser Lys Ile Pro Ala Leu Leu Thr Arg Pro Ile Leu Ser
                85           90           95
Glu Trp Ser Glu Gln Leu His Thr Pro Asn Thr Pro Gly Lys Ala Glu
                100          105          110
Ser Arg Pro Lys Leu Glu Ile Lys Val Ser Pro Leu Pro Leu Ser Val
                115          120          125
Pro Ser Val Gln Leu His Gln Ile Pro Thr Arg Ser Arg Arg Ser Ser
        130          135          140

```

-continued

Lys Pro Arg Lys Pro Arg Lys Lys Arg Lys Glu Arg Val Arg Pro Val  
 145 150 155 160

Ser Arg Val Pro Lys Ala Leu Leu Arg Glu Met Asp Arg Leu Met Thr  
 165 170 175

Lys Gln Arg Asp Ala Leu Pro Glu Ser Glu Ser Ser Ser Tyr Phe Ser  
 180 185 190

Ser Asp Ser Leu Thr Asp Pro Trp Thr Thr Ser Asp Asp Asp Phe Gln  
 195 200 205

Ser Asp Pro Asp Pro Leu Thr Asn Lys Arg Lys Lys Arg Leu Gln Phe  
 210 215 220

<210> SEQ ID NO 26  
 <211> LENGTH: 228  
 <212> TYPE: PRT  
 <213> ORGANISM: Torque teno virus

<400> SEQUENCE: 26

Met Pro Glu His Trp Glu Glu Ala Trp Leu Glu Ala Thr Lys Gly Trp  
 1 5 10 15

His Asp Leu Asp Cys Arg Cys Gly Asn Trp Gln Asp His Leu Trp Leu  
 20 25 30

Leu Leu Ala Asp Gly Asp Ala Ala Leu Ala Ala Val Asp Ala Ile  
 35 40 45

Glu Arg Asp Ala Met Gly Gly Glu Asp Val Thr Thr Ala Thr Asp Arg  
 50 55 60

Val Thr Ile Gly Asp Asp Gly Cys Leu Ala Val Asn Thr Ser His Gln  
 65 70 75 80

Gln Val Ser Ala Ile Pro Ala Leu Ile His Gln Pro Ile Leu Cys Arg  
 85 90 95

Ser Gln Glu Val Leu His Thr Pro Asn Ser Pro Glu Arg Ala Glu Cys  
 100 105 110

Ser Arg Lys Gln Thr Val Gly Val Ser Leu Leu Pro Leu Pro Glu Pro  
 115 120 125

Ser Val Gln Ile His Pro Pro Lys Gln Arg Lys Val His Phe Ser Glu  
 130 135 140

Gly Thr Arg Lys Arg Lys Glu Arg Lys Pro Arg Lys Pro Arg His Arg  
 145 150 155 160

Pro Val Ser Arg Val Pro Lys Ala Leu Leu Arg Glu Met Asp Arg Leu  
 165 170 175

Met Met Lys Arg Gln Ser Asp Ala Glu Gly Gly Pro Gly Ser Asp Ser  
 180 185 190

Asp Gly Trp Ser Asp Ser Ser Leu Thr Asp Glu Trp Thr Thr Ser Asp  
 195 200 205

Ser Asp Phe Ile Asp Thr Pro Ile Arg Glu Arg Cys Leu Asn Lys Lys  
 210 215 220

Gln Lys Lys Arg  
 225

<210> SEQ ID NO 27  
 <211> LENGTH: 199  
 <212> TYPE: PRT  
 <213> ORGANISM: Torque teno virus

<400> SEQUENCE: 27

Met Glu Glu Arg Trp Leu Thr Val Ala Tyr Cys Ala His Gly Leu Phe  
 1 5 10 15



-continued

180	185	190	
Ser Ser Thr Ser Ser Gly Trp 195			
<210> SEQ ID NO 29 <211> LENGTH: 21 <212> TYPE: DNA <213> ORGANISM: Torque teno virus <400> SEQUENCE: 29			
tccgaatggc tgagtttatg c			21
<210> SEQ ID NO 30 <211> LENGTH: 17 <212> TYPE: DNA <213> ORGANISM: Torque teno virus <400> SEQUENCE: 30			
tccgctcagc tgctcct			17
<210> SEQ ID NO 31 <211> LENGTH: 18 <212> TYPE: DNA <213> ORGANISM: Torque teno virus <400> SEQUENCE: 31			
ggtggtaaag aggatgaa			18
<210> SEQ ID NO 32 <211> LENGTH: 19 <212> TYPE: DNA <213> ORGANISM: Torque teno virus <400> SEQUENCE: 32			
aatagattgg acacaggag			19
<210> SEQ ID NO 33 <211> LENGTH: 18 <212> TYPE: DNA <213> ORGANISM: Torque teno virus <400> SEQUENCE: 33			
tatcgggcag gacagct			18
<210> SEQ ID NO 34 <211> LENGTH: 18 <212> TYPE: DNA <213> ORGANISM: Torque teno virus <400> SEQUENCE: 34			
taggggcgcg ctctacgt			18
<210> SEQ ID NO 35 <211> LENGTH: 21 <212> TYPE: DNA <213> ORGANISM: Torque teno virus <400> SEQUENCE: 35			
cctacatgaa ggagaaagac t			--
<210> SEQ ID NO 36 <211> LENGTH: 17 <212> TYPE: DNA			

-continued

---

<213> ORGANISM: Torque teno virus  
 <400> SEQUENCE: 36  
 ccagcgtctc cagggtc 17

<210> SEQ ID NO 37  
 <211> LENGTH: 17  
 <212> TYPE: DNA  
 <213> ORGANISM: Torque teno virus  
 <400> SEQUENCE: 37  
 aagctaccaa gggctgg 17

<210> SEQ ID NO 38  
 <211> LENGTH: 19  
 <212> TYPE: DNA  
 <213> ORGANISM: Torque teno virus  
 <400> SEQUENCE: 38  
 gcggtctggt agcggtagt 19

<210> SEQ ID NO 39  
 <211> LENGTH: 22  
 <212> TYPE: DNA  
 <213> ORGANISM: Torque teno virus  
 <400> SEQUENCE: 39  
 cgaatggctg agtttatgcc gc 22

<210> SEQ ID NO 40  
 <211> LENGTH: 11  
 <212> TYPE: DNA  
 <213> ORGANISM: Torque teno virus  
 <400> SEQUENCE: 40  
 agtcctcatt t 11

<210> SEQ ID NO 41  
 <211> LENGTH: 11  
 <212> TYPE: DNA  
 <213> ORGANISM: Torque teno virus  
 <400> SEQUENCE: 41  
 aaccaatcag a 11

<210> SEQ ID NO 42  
 <211> LENGTH: 17  
 <212> TYPE: DNA  
 <213> ORGANISM: Torque teno virus  
 <400> SEQUENCE: 42  
 ctgggcccgggt gccggag 17

<210> SEQ ID NO 43  
 <211> LENGTH: 14  
 <212> TYPE: DNA  
 <213> ORGANISM: Torque teno virus  
 <400> SEQUENCE: 43  
 cggagtcaag gggc 14

<210> SEQ ID NO 44  
 <211> LENGTH: 11

-continued

---

<212> TYPE: DNA  
 <213> ORGANISM: Torque teno virus  
  
 <400> SEQUENCE: 44  
  
 tatcgggcag g 11

<210> SEQ ID NO 45  
 <211> LENGTH: 23  
 <212> TYPE: PRT  
 <213> ORGANISM: Torque teno virus  
  
 <400> SEQUENCE: 45  
  
 Thr Ala Cys Ala Cys Thr Thr Cys Cys Gly Gly Gly Thr Thr Cys Ala  
 1 5 10 15  
  
 Gly Gly Ala Gly Gly Cys Thr  
 20

<210> SEQ ID NO 46  
 <211> LENGTH: 22  
 <212> TYPE: DNA  
 <213> ORGANISM: Torque teno virus  
  
 <400> SEQUENCE: 46  
  
 actcagccat tcggaacctc ac 22

<210> SEQ ID NO 47  
 <211> LENGTH: 22  
 <212> TYPE: DNA  
 <213> ORGANISM: Torque teno virus  
  
 <400> SEQUENCE: 47  
  
 caatttggt cgcttcgctc gc 22

<210> SEQ ID NO 48  
 <211> LENGTH: 25  
 <212> TYPE: DNA  
 <213> ORGANISM: Torque teno virus  
  
 <400> SEQUENCE: 48  
  
 tacttatatt cgctttcgtg ggaac 25

<210> SEQ ID NO 49  
 <211> LENGTH: 22  
 <212> TYPE: DNA  
 <213> ORGANISM: Torque teno virus  
  
 <400> SEQUENCE: 49  
  
 agttacacat aaccacaaa cc 22

<210> SEQ ID NO 50  
 <211> LENGTH: 21  
 <212> TYPE: DNA  
 <213> ORGANISM: Torque teno virus  
  
 <400> SEQUENCE: 50  
  
 attaccgctt gcccgatagg c 21

<210> SEQ ID NO 51  
 <211> LENGTH: 21  
 <212> TYPE: DNA  
 <213> ORGANISM: Torque teno virus  
  
 <400> SEQUENCE: 51

-continued

---

 ccaaaccaca ggaaactgtg c 21

<210> SEQ ID NO 52  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Torque teno virus

<400> SEQUENCE: 52

cttgactccg ctctcaggag 20

<210> SEQ ID NO 53  
 <211> LENGTH: 2878  
 <212> TYPE: DNA  
 <213> ORGANISM: Torque teno virus

<400> SEQUENCE: 53

tacacttccg gggtcaggag gctcaatttg gctcgtctcg ctgcaccac gtttgetgcc 60

aggcggacct gattgaagac tgaaaaccgt taaattcaaa tttgaaaatg gcgggcaaaa 120

tggcggacag gggcggggga ttatgcaaat taatttatgc aaagtaggag gagctcgatt 180

ttaatttatg caaagtagga ggagtcattt ctgattggtc gggagctcaa gtcctcattt 240

gcataggggtg taaccaatca gatttaagge gttcccacta aagtgaatat aagtgagtgc 300

agttccgaat ggctgagttt atgccgccag cggtagacag aactgtctag cgactgggcg 360

ggtgcccggag gatcccagat ccggagtcaa ggggcctatc gggcaggagc agctgagcgg 420

agggcctaca tgaaggagaa agactactgg gaagaagcct ggctgaccag ctgtacatcc 480

atacacgacc accactgcga ctgcggtagc tggagagacc acctgtggac gctatgcgct 540

ttagacgacg cagatttggc cgccgcgca gatattatag aaagagaaga ggcggatgga 600

ggagaagatt tcggattcgt agacggcgac cctggagacg ctggcgggta aggagatggc 660

ggcgttccgt cttccgtaga aggggacgta gagcgcgccc ctaccgcatt agcgcgtgga 720

accctaaggt tctaagaaac tgccgcatca caggatggtg gccagtaata cagtgtatgg 780

acgggatgga gtggataaaa tacaagccga tggacttaag agtcgaggca aaccggatat 840

tcgataaaca gggcagtaag atagagacag aacagatggg atacttaatg cagtacggag 900

gaggatggtc aagcggagta atcagcttag agggactttt caatgaaaac agactgtgga 960

gaaacatatg gtctaaaagc aatgacggga tggacttggc cagatacttc ggggtgcagaa 1020

ttagactata tccaacagag aatcagggct acttgttctg gtatgacaca gaatttgacg 1080

aacagcagag aagaatgta gacgaatata cacaacctag tgtaatgctg caggctaaaa 1140

actcccgttt aatagtatgt aaacaaaaga tgccaattag acggagagta aagagcattt 1200

tcataccgcc accggcacag ttaacaacac agtggaaagt tcagcaggaa ctgtgtcaat 1260

ttccattatt taactgggcc tgtatctgta tagacatgga cacgccgttc gactacaacg 1320

gcgcatggcg aaatgcctgg tggctaataa gaaggcttca aaacggaaac atggagtaca 1380

tagaaagatg gggcagaata ccgatgacag gagacacaga actgccacca gcagacgact 1440

tcaaggcagg aggggtgaac aaaaacttca aaccgacagg tattcagaga atataccta 1500

tagtagcagt atgcctagtg gagggaaaca agagagtagt gaaatgggcc acagtacaca 1560

atgggccaat agacagatgg agaaaaaac agacaggaac gttaaaacta tctgcactga 1620

gaagactagt gcttagagta tgctcagaaa gtgagacata ctataagtgg acagcatcag 1680

aatttacagg agcatttcag caggactggc ggccagttag cggaacagaa taccggttat 1740

gtacaattaa aatggagcca gaattcagaa acccgacagt agaggtgtgg tcatggaaa 1800

-continued

---

```

caactatacc gacagcagga acaactgaaag actatattcgg gctcagttca gggcaacagt 1860
ggaaggacac tgactttggc aggctgcaat taccagaag cagccacaat gttgactttg 1920
gacataaagc tagatttggc ccattttgtg tgaaaaagcc tccagtagaa ttcagagact 1980
cagcccccaa cccactaaat atctgggtga aatacacatt ctattttcag ttcggcgcca 2040
tgtaccagcc tcccaccgga atccaagatc cctgcacttc taaccggacc tatectgtca 2100
gaatggtcgg agcagttaca cccccaaat acgcccggca aggcggaatc ggcacccaaa 2160
ttggagatca aggtatcacc gctgcctctc tccgtgccat cagtgcagct ccaccaaata 2220
cctacacgca gtcggcgctc ctcaaagccc cggaaaccga gaaagaagag gaaagagaga 2280
gtgagaccag tttcacgagt gccgaaagct cttctgaggg agatggatcg tctgatgacc 2340
aagcagagag acgctgtgcc agaaagcgag tcatcaagct acttotcaag cgactcgctg 2400
acagaccctg ggacaacaag cgacgacgat tttcagagtg accctgacct cctcaccaat 2460
aaacgcaaaa agccttgca attctaattc gctgtccgtg tattcattgg gggggtcgg 2520
gggggcttgc cccccgta gttgggttct cgcactcccg cctgccaagt gaaagtggg 2580
gacgagtgag tgcgggacat cccgtgtaat ggctacataa ctaccgggt ttgcttgac 2640
agtggccgtg gctcgacct cacacaacaa tgcagatagg gggcgcaatt gggatcgta 2700
gaaaactatg gccgagcatg gggggggctc cgccccccc aaccccccg gtgggggggc 2760
caaggcccc cctacacccc cccatggggg gctgctgccc cccaaacccc ccgctgga 2820
tggggggggc tgcgcccc ccaaaccccc cttgcccggg gctgtgcccc ggaccccc 2878

```

&lt;210&gt; SEQ ID NO 54

&lt;211&gt; LENGTH: 635

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Torque teno virus

&lt;400&gt; SEQUENCE: 54

```

Met Arg Phe Arg Arg Arg Arg Phe Gly Arg Arg Arg Arg Tyr Tyr Arg
1           5           10          15
Lys Arg Arg Gly Gly Trp Arg Arg Arg Phe Arg Ile Arg Arg Arg Arg
20          25          30
Pro Trp Arg Arg Trp Arg Val Arg Arg Trp Arg Arg Ser Val Phe Arg
35          40          45
Arg Arg Gly Arg Arg Ala Arg Pro Tyr Arg Ile Ser Ala Trp Asn Pro
50          55          60
Lys Val Leu Arg Asn Cys Arg Ile Thr Gly Trp Trp Pro Val Ile Gln
65          70          75          80
Cys Met Asp Gly Met Glu Trp Ile Lys Tyr Lys Pro Met Asp Leu Arg
85          90          95
Val Glu Ala Asn Arg Ile Phe Asp Lys Gln Gly Ser Lys Ile Glu Thr
100         105         110
Glu Gln Met Gly Tyr Leu Met Gln Tyr Gly Gly Gly Trp Ser Ser Gly
115         120         125
Val Ile Ser Leu Glu Gly Leu Phe Asn Glu Asn Arg Leu Trp Arg Asn
130         135         140
Ile Trp Ser Lys Ser Asn Asp Gly Met Asp Leu Val Arg Tyr Phe Gly
145         150         155         160
Cys Arg Ile Arg Leu Tyr Pro Thr Glu Asn Gln Gly Tyr Leu Phe Trp
165         170         175
Tyr Asp Thr Glu Phe Asp Glu Gln Gln Arg Arg Met Leu Asp Glu Tyr

```

-continued

180					185					190					
Thr	Gln	Pro	Ser	Val	Met	Leu	Gln	Ala	Lys	Asn	Ser	Arg	Leu	Ile	Val
		195						200					205		
Cys	Lys	Gln	Lys	Met	Pro	Ile	Arg	Arg	Arg	Val	Lys	Ser	Ile	Phe	Ile
	210					215					220				
Pro	Pro	Pro	Ala	Gln	Leu	Thr	Thr	Gln	Trp	Lys	Phe	Gln	Gln	Glu	Leu
	225				230					235					240
Cys	Gln	Phe	Pro	Leu	Phe	Asn	Trp	Ala	Cys	Ile	Cys	Ile	Asp	Met	Asp
				245					250					255	
Thr	Pro	Phe	Asp	Tyr	Asn	Gly	Ala	Trp	Arg	Asn	Ala	Trp	Trp	Leu	Met
			260					265						270	
Arg	Arg	Leu	Gln	Asn	Gly	Asn	Met	Glu	Tyr	Ile	Glu	Arg	Trp	Gly	Arg
		275					280					285			
Ile	Pro	Met	Thr	Gly	Asp	Thr	Glu	Leu	Pro	Pro	Ala	Asp	Asp	Phe	Lys
	290					295					300				
Ala	Gly	Gly	Val	Asn	Lys	Asn	Phe	Lys	Pro	Thr	Gly	Ile	Gln	Arg	Ile
	305				310					315					320
Tyr	Pro	Ile	Val	Ala	Val	Cys	Leu	Val	Glu	Gly	Asn	Lys	Arg	Val	Val
				325					330					335	
Lys	Trp	Ala	Thr	Val	His	Asn	Gly	Pro	Ile	Asp	Arg	Trp	Arg	Lys	Lys
			340					345					350		
Gln	Thr	Gly	Thr	Leu	Lys	Leu	Ser	Ala	Leu	Arg	Arg	Leu	Val	Leu	Arg
		355					360					365			
Val	Cys	Ser	Glu	Ser	Glu	Thr	Tyr	Tyr	Lys	Trp	Thr	Ala	Ser	Glu	Phe
	370					375					380				
Thr	Gly	Ala	Phe	Gln	Gln	Asp	Trp	Trp	Pro	Val	Ser	Gly	Thr	Glu	Tyr
	385				390					395					400
Pro	Leu	Cys	Thr	Ile	Lys	Met	Glu	Pro	Glu	Phe	Glu	Asn	Pro	Thr	Val
				405					410					415	
Glu	Val	Trp	Ser	Trp	Lys	Ala	Thr	Ile	Pro	Thr	Ala	Gly	Thr	Leu	Lys
			420					425					430		
Asp	Tyr	Phe	Gly	Leu	Ser	Ser	Gly	Gln	Gln	Trp	Lys	Asp	Thr	Asp	Phe
		435					440					445			
Gly	Arg	Leu	Gln	Leu	Pro	Arg	Ser	Ser	His	Asn	Val	Asp	Phe	Gly	His
	450					455					460				
Lys	Ala	Arg	Phe	Gly	Pro	Phe	Cys	Val	Lys	Lys	Pro	Pro	Val	Glu	Phe
	465				470					475					480
Arg	Asp	Ser	Ala	Pro	Asn	Pro	Leu	Asn	Ile	Trp	Val	Lys	Tyr	Thr	Phe
				485					490					495	
Tyr	Phe	Gln	Phe	Gly	Gly	Met	Tyr	Gln	Pro	Pro	Thr	Gly	Ile	Gln	Asp
			500					505					510		
Pro	Cys	Thr	Ser	Asn	Pro	Thr	Tyr	Pro	Val	Arg	Met	Val	Gly	Ala	Val
		515					520					525			
Thr	His	Pro	Lys	Tyr	Ala	Gly	Gln	Gly	Gly	Ile	Ala	Thr	Gln	Ile	Gly
	530					535					540				
Asp	Gln	Gly	Ile	Thr	Ala	Ala	Ser	Leu	Arg	Ala	Ile	Ser	Ala	Ala	Pro
	545					550				555					560
Pro	Asn	Thr	Tyr	Thr	Gln	Ser	Ala	Phe	Leu	Lys	Ala	Pro	Glu	Thr	Glu
				565					570					575	
Lys	Glu	Glu	Glu	Arg	Glu	Ser	Glu	Thr	Ser	Phe	Thr	Ser	Ala	Glu	Ser
				580					585				590		
Ser	Ser	Glu	Gly	Asp	Gly	Ser	Ser	Asp	Asp	Gln	Ala	Glu	Arg	Arg	Ala
		595						600					605		

-continued

Ala Arg Lys Arg Val Ile Lys Leu Leu Leu Lys Arg Leu Ala Asp Arg  
610 615 620

Pro Val Asp Asn Lys Arg Arg Arg Phe Ser Glu  
625 630 635

<210> SEQ ID NO 55  
<211> LENGTH: 73  
<212> TYPE: PRT  
<213> ORGANISM: Torque teno virus

<400> SEQUENCE: 55

Met Lys Glu Lys Asp Tyr Trp Glu Glu Ala Trp Leu Thr Ser Cys Thr  
1 5 10 15

Ser Ile His Asp His His Cys Asp Cys Gly Ser Trp Arg Asp His Leu  
20 25 30

Trp Thr Leu Cys Ala Leu Asp Asp Ala Asp Leu Ala Ala Ala Asp  
35 40 45

Ile Ile Glu Arg Glu Glu Ala Asp Gly Gly Glu Asp Phe Gly Phe Val  
50 55 60

Asp Gly Asp Pro Gly Asp Ala Gly Gly  
65 70

<210> SEQ ID NO 56  
<211> LENGTH: 2872  
<212> TYPE: DNA  
<213> ORGANISM: Torque teno virus  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (2719)..(2732)  
<223> OTHER INFORMATION: n i s a , c , g , o r t

<400> SEQUENCE: 56

tacactttgg ggttcaggag gctcaatttg gctcgttcg ctcgcaccac gtttgcgtcc	60
aggcggacct gattgaagac tgaaaaccgt taaattcaaa attgaaaagg gcgggcacaaa	120
tgccggacag ggggaggagt ttatgcaaat taatttatgc aaagtaggag gagctcgatt	180
ttaatttatg caaagtagga ggagtcaaat ctgattggtc gggagctcaa gtcctcattt	240
gcataggggtg taaccaatca gaattaaggc gttcccacga aagcgaatat aagtaggtga	300
ggttccgaat ggctgagttt atgccgccag cggtagacag aactgtctag cgactgggcg	360
ggtgccggag gatccctgat ccggagtcaa ggggcctatc gggcaggagc agctaggcgg	420
agggcctatg ccggaacct gggaggaagc ctggttgaa gctaccaagg gctggcacga	480
tctcgactgc cgctgcggta actggcagga ccacctatgg ctctactcg ccgatggaga	540
cgccgctttg gccgccgcc tagacgctat agaaagagac gctatggctg gagacgacgc	600
tactaccgct acaggccgcg tgactatcgg cgacgatggc tggttaaggag aaggcggcgt	660
tccgtctacc gtagaggtgg acgtagagcg cgcccctacc gactgtttaa tccaaaagta	720
atgccggagag tagtaattag ggggtggtgg cctattttac aatgcttaa aggacaggag	780
gcactaagat atagacctct acagtgggac acagagagac agtggagagt gagatcagac	840
ttcgaagacc agtacggata cctcgtacaa tacgggggag gttggggaag tggatgatgtg	900
acacttgaag gtctctacca agagcactta ttgtggagaa actcttggtc taaaggaaac	960
gatggaatgg acctagtaag atactttgga tgtgtagtat acctatatcc actaaaggac	1020
caggactatt ggttctggtg ggacacggac ttcaaagaat tatatgcaga aaacataaag	1080
gaatacagcc aaccatcagt aatgatgatg gcaaaaagaa caagaatagt aatagccaga	1140

-continued

---

```

gaaagggcac cacatagaag aaaagtaaga aaaatattta ttcggccacc ttcgagagac 1200
acaacacagt ggcagtttca gacagatttc tgcaatagaa agttatttac gtgggcagct 1260
ggtctaatac acatgcaaaa accgttcgat gctaattggag cctttagaaa tgcttggtgg 1320
ctggaacaga gaaatgatca gggagaaatg aaatacatag aactgtgggg aagagtaccc 1380
ccacaaggag attcagagct gccccaaaaa aaagaattct ccacaggaac agataacca 1440
aactacaatg ttcaggacaa tgaggagaaa aacatatacc ccattataat atacgtagac 1500
caaaaagatc aaaaaccaag aaaaaagtac tgcgtatggt ataataagac cctcaacaga 1560
tggagactag gacaggcaag tactctaaag ataggaaacc tgaaaggact agtactaaga 1620
cagctgatga atcaagaat gacgtatata tggaaagaag gagaatacag tgcccccttt 1680
gtacaaaggc ggaaggcag cagattcgct gtgatagacg caagaaaggc agaccaagaa 1740
aaccgaaaag tatcaacatg gccaatgag ggaacgtgga acacacagga cacagtactg 1800
aaggatgatc tcggtattaa cttgcaaaaat caacaattta gggcggcgga ctttggtaaa 1860
ctcacactac caaaatcacc gcatgactta gacttcggtc accacagcag atttgggcca 1920
ttttgtgtga aaaatgaacc actggagttt caggtatacc ctccagaacc aactaactg 1980
tggtttcagt acagattttt ctttcagttt ggaggtgaat accaaccccc cacaggaatc 2040
cgggatccat gcggtgatac accagcctat cctgtgccgc agtcagggaag tattacacac 2100
cccaaattcg ccggaaaaag aggaatgctc acggaacag accgttgggg tatcactgct 2160
gcctcttcca gagccctcag tgcagataca cccacagagg cagcgcaaaag tgcacttctc 2220
cgaggggact cggaaagcga aggagaggaa accgaggaaa ccgctgcatc gtccagtatc 2280
acgagtgcgc aaagctctac tgaggagat ggatcgtctg atgatgaaga gacaatcaga 2340
cgcagaagga ggacctgga ggcactcaga cgaatggta gagagcagct tgaccgacga 2400
atggaccaca agcagacagc acttcattga ccccccata agagaaagat gcctcaataa 2460
aaaacaaaag aaacgctaaa cagtgtccga ttactaatgg ggggggggtcc ggggggggct 2520
tgcccccccg caagctgggt taccgcaacta actcctgccc aagtgaaact cggggacgag 2580
tgagtgcggg acatcccgtg taatggctac ataactaccc ggctttgctt cgacagtggc 2640
cgtggctcga ccctcacaca aactgcagg tagggggcgc aattgggac gttagaaaac 2700
tatggcgcag catggggggn nnnnnnnnnn nccaacccc cccgggtggg gggccaaggc 2760
ccccctaca ccccccatg gggggctgcc gcccccaaa cccccgcgt cggatggggg 2820
gggctgcgcc cccccaaac ccccttgcc cgggctgtg ccccgaccc cc 2872

```

&lt;210&gt; SEQ ID NO 57

&lt;211&gt; LENGTH: 637

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Torque teno virus

&lt;400&gt; SEQUENCE: 57

```

Met Ala Pro Thr Arg Arg Trp Arg Arg Arg Phe Gly Arg Arg Arg Arg
1           5           10           15

```

```

Arg Tyr Arg Lys Arg Arg Tyr Gly Trp Arg Arg Arg Tyr Tyr Arg Tyr
20           25           30

```

```

Arg Pro Arg Asp Tyr Arg Arg Arg Trp Leu Val Arg Arg Arg Arg Arg
35           40           45

```

```

Ser Val Tyr Arg Arg Gly Gly Arg Arg Ala Arg Pro Tyr Arg Leu Phe
50           55           60

```

-continued

---

Asn Pro Lys Val Met Arg Arg Val Val Ile Arg Gly Trp Trp Pro Ile  
 65 70 75 80  
 Leu Gln Cys Leu Lys Gly Gln Glu Ala Leu Arg Tyr Arg Pro Leu Gln  
 85 90 95  
 Trp Asp Thr Glu Arg Gln Trp Arg Val Arg Ser Asp Phe Glu Asp Gln  
 100 105 110  
 Tyr Gly Tyr Leu Val Gln Tyr Gly Gly Gly Trp Gly Ser Gly Asp Val  
 115 120 125  
 Thr Leu Glu Gly Leu Tyr Gln Glu His Leu Leu Trp Arg Asn Ser Trp  
 130 135 140  
 Ser Lys Gly Asn Asp Gly Met Asp Leu Val Arg Tyr Phe Gly Cys Val  
 145 150 155 160  
 Val Tyr Leu Tyr Pro Leu Lys Asp Gln Asp Tyr Trp Phe Trp Trp Asp  
 165 170 175  
 Thr Asp Phe Lys Glu Leu Tyr Ala Glu Asn Ile Lys Glu Tyr Ser Gln  
 180 185 190  
 Pro Ser Val Met Met Met Ala Lys Arg Thr Arg Ile Val Ile Ala Arg  
 195 200 205  
 Glu Arg Ala Pro His Arg Arg Lys Val Arg Lys Ile Phe Ile Pro Pro  
 210 215 220  
 Pro Ser Arg Asp Thr Thr Gln Trp Gln Phe Gln Thr Asp Phe Cys Asn  
 225 230 235 240  
 Arg Lys Leu Phe Thr Trp Ala Ala Gly Leu Ile Asp Met Gln Lys Pro  
 245 250 255  
 Phe Asp Ala Asn Gly Ala Phe Arg Asn Ala Trp Trp Leu Glu Gln Arg  
 260 265 270  
 Asn Asp Gln Gly Glu Met Lys Tyr Ile Glu Leu Trp Gly Arg Val Pro  
 275 280 285  
 Pro Gln Gly Asp Ser Glu Leu Pro Lys Lys Lys Glu Phe Ser Thr Gly  
 290 295 300  
 Thr Asp Asn Pro Asn Tyr Asn Val Gln Asp Asn Glu Glu Lys Asn Ile  
 305 310 315 320  
 Tyr Pro Ile Ile Ile Tyr Val Asp Gln Lys Asp Gln Lys Pro Arg Lys  
 325 330 335  
 Lys Tyr Cys Val Cys Tyr Asn Lys Thr Leu Asn Arg Trp Arg Leu Gly  
 340 345 350  
 Gln Ala Ser Thr Leu Lys Ile Gly Asn Leu Lys Gly Leu Val Leu Arg  
 355 360 365  
 Gln Leu Met Asn Gln Glu Met Thr Tyr Ile Trp Lys Glu Gly Glu Tyr  
 370 375 380  
 Ser Ala Pro Phe Val Gln Arg Trp Lys Gly Ser Arg Phe Ala Val Ile  
 385 390 395 400  
 Asp Ala Arg Lys Ala Asp Gln Glu Asn Pro Lys Val Ser Thr Trp Pro  
 405 410 415  
 Ile Glu Gly Thr Trp Asn Thr Gln Asp Thr Val Leu Lys Asp Val Phe  
 420 425 430  
 Gly Ile Asn Leu Gln Asn Gln Gln Phe Arg Ala Ala Asp Phe Gly Lys  
 435 440 445  
 Leu Thr Leu Pro Lys Ser Pro His Asp Leu Asp Phe Gly His His Ser  
 450 455 460  
 Arg Phe Gly Pro Phe Cys Val Lys Asn Glu Pro Leu Glu Phe Gln Val  
 465 470 475 480  
 Tyr Pro Pro Glu Pro Thr Asn Leu Trp Phe Gln Tyr Arg Phe Phe Phe



-continued

---

ttacagacgc tatcgacgac gccgaagaag accgacacgg agatggaggc accggaggtg	540
gagacgctac tttc gatatc ggtatcgacg cgctcctcgc cgccgccgca caaaggtaag	600
gagacggagg aaaaaagctc cggtcataca atggttccct cctagccgga gaacctgcct	660
catagaggga ttttggcgtg tgagctacgg acactggttc cgtacctgtc tccccttag	720
gcggttaaat ggactagtat tcccgggtgg aggttgtagc tggagccagt ggagtttaca	780
aaacctttac aatgaaaaac ttaactggag aaatatatgg acagctagta atgttggaat	840
ggaattcgct agatttttaa aaggaaagt ttacttttcc agacatccat ggagaaatta	900
tataataact tgggatcaag atataccatg caggccacta ccttatcaaa acctgcaccc	960
actcctaagt ctactaaaaa aacagcacia aattgtactt tcacagcaaa actgtaaccc	1020
aaacagaaaa caaaaacctg tcacattaaa attcaaacct ccgcaaaaa taacatcaca	1080
atggagacta agtagagaat tagcaaatg gccactaata agacttggag taagctttat	1140
agacctaaca gaacctggg tagaagggg gggaaatgca ttttattccg tgctaggata	1200
tgaagcagta aaagaccaag gacactggtc aaactggaca caaataaaat actattggat	1260
ctatgacacg ggagtaggaa atgcagtata tgttatacta ttaaaaaag acgttactga	1320
taatccagga aacatggcaa caacctttaa agcatcagga ggacagcatc cagatgcaat	1380
agatcacatt gaattgataa accaaggatg gccttactgg ttatactttt atggtaaaaag	1440
tgaacaagac attaaaaaag aggcacacag cgcagaaata tcaagagaat atactagaga	1500
cccaaatct aaaaaactaa aaataggaat agtaggatgg gcatcttcaa actacacaac	1560
aacaggcagt gatcaaaaca gtgggtggatc aacatcagct atacaagggtg gatatgtagc	1620
atatgcaggg tccgggggtca taggagcagg gtcaatagga aatttatatc aacaaggatg	1680
gccatcctaat caaaactggc ctaatacaaa cagagacaaa acaaaacttg actggggaat	1740
acgaggacta tgtatactca gagataacat gcacttagga agccaagaat tagatgatga	1800
atgcacaatg ctcacattgt tcggaccctt tgtagaaaaa gcaaatccaa tatttgcaac	1860
aacagacctt aattctttaa aacctgaact caaagactat aatataatca tgaatatgct	1920
ctttaaattt cagtggggag gacatggcac agaaagattt aaaaccaaca tcgggagacc	1980
cagcaccata ccctgccctc tcgaaccctg ggaccgctc cacagcggga tacaagacct	2040
ctccaaggta caaaacaccg tcctcaacct ctgggactat gactgtgatg ggattgttag	2100
aaaagatact ctcaaaagac ttctcgaact cccacagag acagaggagg aggagaaggc	2160
gtaccactc cttggacaaa aaacagagaa agagccatta tcagactccg acgaagagag	2220
cgttatctca agcagcagca gtggatcctc tcaagaagaa gaaacgcaga gacgaagaca	2280
ccacaagcca agcaagcgc gactcctcaa gcacctccag cgggtggtaa agaggatgaa	2340
aacactgtga tagataaata tagaaacctc gcagaccctc cactcaatgt cacaggacac	2400
atggaaaaat tcattgcagtt acatattcaa aacgtacaag aaataagagc taaaaatgct	2460
aaaaaatccc tcaataaact ttacttttct gattaatagc ggctcctgt gtccaacctc	2520
tttttctcaa accccttcaa aatggcgggc gggacacaaa atggcggagg gactaagggg	2580
ggggcaagcc cccctnnnnn nnnnnnnnnn nnnnnnnnnn nngggggggc acccccccg	2640
ccccccctc gcgggggctc cgccccctgc acccccgga ggggggaaa cccccctca	2700
cccccccgcg gggggcaagc cccctgcac cccc	2735

-continued

&lt;211&gt; LENGTH: 624

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Torque teno virus

&lt;400&gt; SEQUENCE: 60

Met Pro Tyr Arg Arg Tyr Arg Arg Arg Arg Arg Arg Pro Thr Arg Arg  
 1 5 10 15  
 Trp Arg His Arg Arg Trp Arg Arg Tyr Phe Arg Tyr Arg Tyr Arg Arg  
 20 25 30  
 Ala Pro Arg Arg Arg Arg Thr Lys Val Arg Arg Arg Arg Lys Lys Ala  
 35 40 45  
 Pro Val Ile Gln Trp Phe Pro Pro Ser Arg Arg Thr Cys Leu Ile Glu  
 50 55 60  
 Gly Phe Trp Pro Leu Ser Tyr Gly His Trp Phe Arg Thr Cys Leu Pro  
 65 70 75 80  
 Phe Arg Arg Leu Asn Gly Leu Val Phe Pro Gly Gly Gly Cys Asp Trp  
 85 90 95  
 Ser Gln Trp Ser Leu Gln Asn Leu Tyr Asn Glu Lys Leu Asn Trp Arg  
 100 105 110  
 Asn Ile Trp Thr Ala Ser Asn Val Gly Met Glu Phe Ala Arg Phe Leu  
 115 120 125  
 Lys Gly Lys Phe Tyr Phe Phe Arg His Pro Trp Arg Asn Tyr Ile Ile  
 130 135 140  
 Thr Trp Asp Gln Asp Ile Pro Cys Arg Pro Leu Pro Tyr Gln Asn Leu  
 145 150 155 160  
 His Pro Leu Leu Met Leu Leu Lys Lys Gln His Lys Ile Val Leu Ser  
 165 170 175  
 Gln Gln Asn Cys Asn Pro Asn Arg Lys Gln Lys Pro Val Thr Leu Lys  
 180 185 190  
 Phe Lys Pro Pro Pro Lys Leu Thr Ser Gln Trp Arg Leu Ser Arg Glu  
 195 200 205  
 Leu Ala Lys Met Pro Leu Ile Arg Leu Gly Val Ser Phe Ile Asp Leu  
 210 215 220  
 Thr Glu Pro Trp Val Glu Gly Trp Gly Asn Ala Phe Tyr Ser Val Leu  
 225 230 235 240  
 Gly Tyr Glu Ala Val Lys Asp Gln Gly His Trp Ser Asn Trp Thr Gln  
 245 250 255  
 Ile Lys Tyr Tyr Trp Ile Tyr Asp Thr Gly Val Gly Asn Ala Val Tyr  
 260 265 270  
 Val Ile Leu Leu Lys Lys Asp Val Thr Asp Asn Pro Gly Asn Met Ala  
 275 280 285  
 Thr Thr Phe Lys Ala Ser Gly Gly Gln His Pro Asp Ala Ile Asp His  
 290 295 300  
 Ile Glu Leu Ile Asn Gln Gly Trp Pro Tyr Trp Leu Tyr Phe Tyr Gly  
 305 310 315 320  
 Lys Ser Glu Gln Asp Ile Lys Lys Glu Ala His Ser Ala Glu Ile Ser  
 325 330 335  
 Arg Glu Tyr Thr Arg Asp Pro Lys Ser Lys Lys Leu Lys Ile Gly Ile  
 340 345 350  
 Val Gly Trp Ala Ser Ser Asn Tyr Thr Thr Thr Gly Ser Asp Gln Asn  
 355 360 365  
 Ser Gly Gly Ser Thr Ser Ala Ile Gln Gly Gly Tyr Val Ala Tyr Ala  
 370 375 380  
 Gly Ser Gly Val Ile Gly Ala Gly Ser Ile Gly Asn Leu Tyr Gln Gln



-continued

20			25			30									
Ala	Pro	Arg	Arg	Arg	Arg	Ala	Lys	Val	Arg	Arg	Arg	Arg	Arg	Lys	Ala
	35						40						45		
Pro	Val	Ile	Gln	Trp	Asn	Pro	Pro	Ser	Arg	Arg	Thr	Cys	Leu	Ile	Glu
	50						55					60			
Gly	Phe	Trp	Pro	Leu	Ser	Tyr	Gly	His	Trp	Phe	Arg	Thr	Cys	Leu	Pro
	65				70						75				80
Met	Arg	Arg	Leu	Asn	Gly	Leu	Ile	Phe	Thr	Gly	Gly	Gly	Cys	Asp	Trp
				85					90					95	
Thr	Gln	Trp	Ser	Leu	Gln	Asn	Leu	Phe	His	Glu	Lys	Leu	Asn	Trp	Arg
			100						105					110	
Asn	Ile	Trp	Thr	Ala	Ser	Asn	Val	Gly	Met	Glu	Phe	Ala	Arg	Phe	Leu
			115				120						125		
Arg	Gly	Lys	Phe	Tyr	Phe	Phe	Arg	His	Pro	Trp	Arg	Ser	Tyr	Ile	Val
	130						135					140			
Thr	Trp	Asp	Gln	Asp	Ile	Pro	Cys	Lys	Pro	Leu	Pro	Tyr	Gln	Asn	Leu
	145				150						155				160
Gln	Pro	Leu	Leu	Met	Leu	Leu	Lys	Lys	Gln	His	Lys	Leu	Val	Leu	Ser
				165					170						175
Gln	Lys	Asp	Cys	Asn	Pro	Ser	Arg	Lys	Gln	Lys	Pro	Val	Thr	Leu	Lys
			180					185						190	
Phe	Arg	Pro	Pro	Pro	Lys	Leu	Thr	Ser	Gln	Trp	Arg	Leu	Ser	Arg	Glu
		195					200						205		
Leu	Ser	Lys	Ile	Pro	Leu	Ile	Arg	Leu	Gly	Ile	Ser	Leu	Ile	Asp	Leu
	210					215					220				
Ser	Glu	Pro	Trp	Leu	Glu	Gly	Trp	Gly	Asn	Ala	Phe	Tyr	Ser	Val	Leu
	225				230					235					240
Gly	Tyr	Glu	Ala	Ser	Lys	His	Ser	Gly	Arg	Trp	Ser	Asn	Trp	Thr	Gln
				245					250						255
Met	Lys	Tyr	Phe	Trp	Ile	Tyr	Asp	Thr	Gly	Val	Gly	Asn	Ala	Val	Tyr
			260					265						270	
Val	Ile	Leu	Leu	Lys	Lys	Asp	Val	Asp	Asp	Asn	Pro	Gly	Asp	Met	Ala
		275					280						285		
Thr	Lys	Phe	Val	Thr	Gly	Gln	Gly	Gln	His	Pro	Asp	Ala	Ile	Asp	His
	290					295					300				
Ile	Glu	Met	Val	Asn	Glu	Gly	Trp	Pro	Tyr	Trp	Leu	Phe	Phe	Tyr	Gly
	305				310					315					320
Gln	Ser	Glu	Gln	Asp	Ile	Lys	Lys	Leu	Ala	His	Asp	Gln	Asp	Ile	Ala
				325					330						335
Arg	Glu	Tyr	Ala	Arg	Asp	Pro	Lys	Ser	Lys	Lys	Leu	Lys	Ile	Gly	Val
			340					345						350	
Ile	Gly	Trp	Ala	Ser	Ser	Asn	Tyr	Thr	Thr	Ala	Gly	Ser	Asn	Gln	Asn
		355					360						365		
Thr	Thr	Ala	Gln	Thr	Pro	Glu	Ala	Ile	Gln	Gly	Gly	Tyr	Val	Ala	Tyr
	370					375						380			
Ala	Gly	Ser	Arg	Ile	Pro	Gly	Ala	Gly	Ser	Ile	Thr	Asn	Leu	Phe	Gln
	385				390					395					400
Met	Gly	Trp	Pro	Gly	Asp	Gln	Asn	Trp	Pro	Pro	Thr	Asn	Gln	Glu	Gln
				405					410						415
Thr	Asn	Phe	Asn	Trp	Gly	Leu	Arg	Gly	Leu	Cys	Val	Leu	Arg	Asp	Asn
			420					425						430	
Met	Lys	Leu	Gly	Ala	Gln	Glu	Leu	Asp	Asp	Glu	Cys	Thr	Met	Leu	Ser
		435						440					445		

-continued

---

Leu Phe Gly Pro Phe Val Glu Lys Ala Asn Thr Ala Phe Ala Thr Asn  
 450 455 460

Asp Pro Lys Tyr Phe Arg Pro Glu Leu Lys Asp Tyr Asn Val Val Met  
 465 470 475 480

Lys Tyr Ala Phe Lys Phe Gln Trp Gly Gly His Gly Thr Glu Arg Phe  
 485 490 495

Lys Thr Thr Ile Gly Asp Pro Ser Thr Ile Pro Cys Pro Phe Glu Pro  
 500 505 510

Gly Glu Arg Tyr His His Gly Val Gln Asp Pro Ala Lys Val Gln Asn  
 515 520 525

Thr Val Leu Asn Pro Trp Asp Tyr Asp Cys Asp Gly Ile Val Arg Thr  
 530 535 540

Asp Thr Leu Lys Arg Leu Leu Glu Leu Pro Thr Glu Thr Glu Glu Thr  
 545 550 555 560

Glu Lys Ala Tyr Pro Leu Leu Gly Gln Lys Thr Glu Lys Glu Pro Leu  
 565 570 575

Ser Asp Ser Asp Glu Glu Ser Val Ile Ser Ser Thr Ser Ser Gly Ser  
 580 585 590

Ser Gln Glu Glu Glu Thr Gln Arg Arg Arg Gln His Lys Pro Ser Lys  
 595 600 605

Arg Arg Leu Leu Lys His Leu Gln Arg Val Val Lys Arg Met Lys Thr  
 610 615 620

Leu  
 625

---

What is claimed is:

1. An immunogenic composition comprising a biologically functional plasmid or baculovirus expression vector comprising an open reading frame (ORF); wherein said ORF encodes a protein comprising amino acids 310-625 of SEQ ID NO:16.
2. The vaccine composition according to claim 1, further comprising an adjuvant.
3. method of eliciting an immune response in a pig against porcine Torque teno virus (PTTV), comprising administering to a pig an immunologically effective amount of the composition according to claim 1.

35 4. The method according to claim 3, which comprises administering the composition parenterally, intranasally, intradermally, or transdermally to the pig.

40 5. The method according to claim 3, which comprises administering the composition intralymphoidly or intramuscularly to the pig.

\* \* \* \* \*