

Transplacental Transmission of a North American Isolate of *Leishmania infantum* in an Experimentally Infected Beagle

Author(s): Alexa C. Rosypal , Gregory C. Troy , Anne M. Zajac , Glenn Frank , and David S. Lindsay

Source: Journal of Parasitology, 91(4):970-972. 2005.

Published By: American Society of Parasitologists

DOI: <http://dx.doi.org/10.1645/GE-483R.1>

URL: <http://www.bioone.org/doi/full/10.1645/GE-483R.1>

BioOne (www.bioone.org) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/page/terms_of_use.

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

summer (June) showed no clinical signs of disease, while histologic examination of the intestines showed only a minimal coccidian infection. This indicates that as fish grow older and, presumably, increasingly more immunocompetent, or as water temperatures increase and stabilize, the fish may be able to eradicate the parasites or become more resistant to infection, reinfection, or pathologic effects (Steinhagen et al., 1998; Paperna, 1999). This is fortunate, because no chemotherapeutics are approved by the U.S. Food and Drug Administration for treatment of piscine coccidiosis (Center for Veterinary Medicine, 2000). Piscine coccidiosis is also not considered a notifiable or significant disease by the World Organization for Animal Health (Office International des Epizooties, 2003), and thus, development of treatments for this disease has received limited attention. Therefore, prevention of clinical coccidiosis currently depends on maintaining a proper environment, providing optimal nutrition, and reducing stress among fish populations.

The authors wish to thank Sheila M. Mitchell, Kathy Lowe, and Virginia Viers for technical assistance with transmission electron microscopy. This project was supported in part by USDA/CSREES grant 2003-34395-13918 and the VT Commercial Fish and Shellfish Technology (CFAST) Program.

LITERATURE CITED

CENTER FOR VETERINARY MEDICINE. 2000. Drugs approved for use in aquaculture (poikilothermic food species). Food and Drug Administration, Blacksburg, Virginia. Web site: www.fda.gov/cvm/index/aquaculture/appendixa6.htm.

- HAYAT, M. A. 1989. Principles of electron microscopy, 3rd ed. CRC Press, Boca Raton, Florida, 325 p.
- LOM, J., S. S. DESSER, AND I. DYKOVA. 1989. Some little-known and new protozoan parasites of fish from Lake Sasajewun, Algonquin Park, Ontario. *Canadian Journal of Zoology* **67**: 1372-1379.
- , AND I. DYKOVA. 1992. Apicomplexans (Phylum Apicomplexa Levine, 1970). In *Protozoan parasites of fishes*, J. Lom and I. Dykova (eds.). Elsevier, New York, New York, p. 87-123.
- MOLNÁR, K. 1995. Phylum Apicomplexa. In *Fish diseases and disorders, Volume 1: Protozoan and metazoan infections*, P. T. K. Woo (ed.). CAB International, Wallingford, U.K., p. 263-287.
- , AND G. HANEK. 1974. Seven new *Eimeria* spp. (Protozoa, Coccidia) from freshwater fishes of Canada. *Journal of Protozoology* **21**: 489-493.
- OFFICE INTERNATIONAL DES EPIZOOTIES. 2003. Diseases and pathogens encompassed within the OIE surveillance system. World Organization for Animal Health. Web site: www.oie.int/fdc/eng/brochure/en_pathogens.htm.
- PAPERNA, I. 1999. Parasites in warmwater aquaculture. In *Fish diseases and disorders*, P. T. K. Woo (ed.). CAB International, Wallingford, U.K., p. 162-165.
- STEINHAGEN, D., K. HESPE, B. ELLMER, AND W. KORTING. 1998. *Goussia carpelli* (Protozoa: Coccidia) infection in stressed and immunosuppressed common carp *Cyprinus carpio*. *Diseases of Aquatic Organisms* **34**: 199-204.
- WOOTEN, R. 1989. The parasitology of teleosts. In *Fish pathology*, R. J. Roberts (ed.). W. B. Saunders, London, U.K., p. 242-287.

J. Parasitol., 91(4), 2005, pp. 970-972
© American Society of Parasitologists 2005

Transplacental Transmission of a North American Isolate of *Leishmania infantum* in an Experimentally Infected Beagle

Alexa C. Rosypal, Gregory C. Troy*, Anne M. Zajac, Glenn Frank†, and David S. Lindsay, Department of Biomedical Sciences and Pathobiology, Virginia-Maryland College of Veterinary Medicine, Virginia Tech, Blacksburg, Virginia 24061; *Department of Small Animal Clinical Sciences, Virginia-Maryland College of Veterinary Medicine, Virginia Tech, Blacksburg, Virginia 24061; †Heska Corporation, 1825 Sharp Point Drive, Fort Collins, Colorado 80525. e-mail: arosypal@vt.edu

ABSTRACT: *Leishmania infantum*, an etiologic agent of zoonotic visceral leishmaniasis, is widespread among foxhounds in the United States. Although sand flies are widely distributed throughout the United States, epidemiological data do not support a major role for sand flies in the transmission of *L. infantum* in foxhounds in this country. Congenital transmission of human visceral leishmaniasis is reported in humans and might also occur in dogs. We have previously isolated *L. infantum* from Virginia foxhounds and used this isolate (LIVT-1) to experimentally infect beagles. Four female beagles, chronically infected with LIVT-1, were bred to a male beagle chronically infected with *L. infantum chagasi*. One beagle was able to maintain her pregnancy, and 4 puppies were delivered by cesarean section. One puppy was malformed and autolytic at delivery, and tissues were not collected or analyzed. The remaining puppies were killed at the time of cesarean section, and selected tissues were collected for parasite culture and PCR. Promastigotes were not cultured from tissues in any of the puppies. *Leishmania* sp. DNA was detectable by PCR in liver, bone marrow, and heart from all 3 puppies and in the spleen, lymph node, kidney, and placenta in 2 puppies. Placental tissue from the dam was PCR negative. This is the first report of maternal transmission of a North American isolate of *L. infantum* from an experimentally infected dog.

Leishmaniasis is a vectorborne disease caused by infection with *Leishmania* spp.; these parasites are usually transmitted by phlebotomine sand flies. Studies have suggested that transmission might also occur infrequently by sexual contact (Catone et al., 2003), blood transfusions (Owens et al., 2001; Giger et al., 2002), and direct contact (Lainson and Bray, 1964; Nuwayri-Salti and Khansa, 1985). Rare cases of con-

genital transmission have been documented in humans (Low and Cooke, 1926; Nyakundi et al., 1988; Yadav et al., 1989; Eltoun et al., 1992; Meinecke et al., 1999). Similarly, it has been suggested that maternal transmission can occasionally occur in dogs (Mancianti and Sozzi, 1995; Masucci et al., 2003).

Leishmania infantum, an etiologic agent of human visceral leishmaniasis, is widespread among foxhounds in the United States (Rosypal et al., 2003). Sand flies are present throughout much of the country, particularly in the southeast (Young and Perkins, 1984). However, epidemiological data do not support a role for sand flies in the transmission of *L. infantum*. This suggests that an alternative mode of transmission could maintain infections in U.S. foxhounds. The risk of zoonotic spread from infected dogs to humans is currently unknown for the United States. In this study, we determined whether vertical transmission can occur in dogs infected with a North American isolate of *L. infantum*.

Intact female beagles (n = 4) were experimentally infected with promastigotes of the LIVT-1 strain of *L. infantum* originally isolated from a naturally infected foxhound from Virginia (Rosypal et al., 2003). All of the dogs had clinical signs compatible with canine leishmaniasis. The parasite had been reisolated from the beagles by culture of lymph node and bone marrow aspirates, and it was detectable by polymerase chain reaction (PCR) conducted on bone marrow (A. Rosypal, unpubl. obs.). The female dogs were bred to a male beagle chronically infected with *L. infantum chagasi* both by natural service and by artificial insemination. Before breeding, semen was collected from the male dog and placed in *Leishmania* sp. culture media (30% v/v fetal bovine serum, 1% penicillin/streptomycin, 2% human urine, in Grace's Insect Media) at 25 C.

TABLE I. PCR results from selected tissues taken at necropsy from puppies delivered from a beagle experimentally infected with a North American isolate of *Leishmania infantum*.

Tissue	PCR results*		
	Puppy 1	Puppy 2	Puppy 3
Liver	+	+	+
Bone marrow	+	+	+
Spleen	+	+	-
Heart	+	+	+
Lymph node	+	-	+
Kidney	-	+	+
Placenta	+	+	-

* +, positive; -, negative.

One of the 4 dogs became pregnant and was able to sustain the pregnancy, but the other 3 dogs either did not conceive or did not maintain their pregnancies. For this reason, the following data were collected from a single female dog that sustained her pregnancy. Four puppies were delivered by cesarean section on day 60 of gestation (normal gestation period = 63 days) to preclude the possibility of transvaginal transmission during natural birth. Puppies were either unable to breathe at birth or were killed immediately following delivery. One puppy was nonviable and deformed at the time of delivery and tissues were not collected. Portions of liver, femoral bone marrow, spleen, heart, lymph node, kidney, and placental tissues were collected from the remaining 3 puppies for parasite culture and PCR.

DNA was extracted from tissue samples with a commercial kit (DNA Mini Kit, Quiagen®, Valencia, California). For each 50- μ l reaction, 1 μ l of DNA was added to 45 μ l of Platinum® PCR Supermix (Invitrogen® Life Technologies, Carlsbad, California) in a 0.5-ml thin-walled microcentrifuge tube. To the reaction tube, 2 μ l of primers 13A and 13B were added; these primers amplify a conserved minicircle region of kinetoplast DNA from all species of *Leishmania* (Rodgers et al., 1990).

Optimal PCR amplification conditions consisted of initial denaturation at 95 C for 2 min; 38 cycles of denaturation at 94 C for 30 sec, annealing at 62 C for 30 sec, and extension at 68 C for 30 sec; and a final extension at 72 C for 10 min. PCR product (1 μ l) was used as template DNA in a second 50- μ l reaction and was subjected to the same cycling conditions described. PCR products were electrophoresed on a 2% agarose gel and with size markers to detect the 116-bp PCR product. DNA extracted from LIVT-1 promastigotes was used a positive control, and a negative control without DNA was included. To determine the sensitivity of the PCR assay, DNA was extracted from a known number of LIVT-1 promastigotes. Tenfold serial dilutions of DNA were made and subjected to PCR. DNA detection limit was <1 organism.

Promastigotes were not cultured from any puppy tissues. Parasites were not isolated from the sire's semen culture. The expected 116-bp product was amplified by PCR in liver, bone marrow, and heart tissue from all 3 puppies (Table I). *Leishmania* sp. DNA was also detectable in spleen, lymph node, kidney, and placental tissues from 2 puppies. No PCR product was detectable in testes collected at necropsy from the sire (data not shown). *Leishmania* sp. DNA was demonstrable by PCR in uterine tissue collected at necropsy from the dam (data not shown). Placental tissue from the dam was PCR negative.

This is the first report of maternal transmission of a strain of *L. infantum* from North America. The most likely route of transmission was across the placenta. Puppies were delivered by cesarean section, which eliminated the possibility of transmission through microscopic lesions in the birth canal, and *Leishmania* sp. was detected by PCR in placenta and uterine tissue from the dam.

Leishmania sp. parasites circulate in blood. The placental blood supply is in close proximity to the maternal circulation, and parasites might pass into fetal circulation. This mode of transmission has been previously reported in humans (Low and Cooke, 1926) and in experimentally infected mice (Nuwayri-Salti and Khansa, 1985). Paternal transmission was excluded because no parasites were cultured from semen and there was no detectable *Leishmania* sp. DNA in testes tissue tested by PCR.

Pregnancy has a systemic effect that biases the immune system toward a Th2 immune response that protects the fetoplacental unit while simultaneously increasing susceptibility to intracellular pathogens (Wegmann et al., 1993). It has been previously shown that Th1 cytokines can be harmful to the placenta and might compromise fetal survival (Wegmann et al., 1993; Raghupathay, 1997). Resistance to leishmaniasis is associated with a strong Th1 type response marked by increased secretion of interferon- γ , tumor necrosis factor, and interleukin 2 compared with symptomatic dogs (Pinelli et al., 1994; Santos-Gomes et al., 2002). Moreover, dogs with clinical disease develop a strong humoral response (Nieto et al., 1999) and a simultaneous lack of an appropriate cell-mediated immune response, resulting in disease progression (Pinelli et al., 1994). Humoral immunity is driven by a Th2 immune response and the production of cytokines by type 2 helper T cells.

The dogs used in this study were clinically symptomatic with high anti-*Leishmania* sp. antibody responses (data not shown), so they were already biased toward a Th2 immune response. In addition to generalized immunosuppression required during pregnancy to prevent immune reactions directed against foreign antigens on the fetus, a shift to a Th2-from a Th1-type response during clinical disease could increase both the severity of disease and the chance of congenital transmission of canine leishmaniasis. Previous work in experimentally infected mice demonstrated increased susceptibility to leishmaniasis during pregnancy (Krishnan, Guilbert, Russell et al., 1996).

In this study, 3 of 4 beagles that were bred were unable to conceive or maintain their pregnancies. Similarly, in a murine model, Krishnan, Guilbert, Wegmann et al. (1996) found that *Leishmania major* infection was associated with increased frequencies of fetal resorption and implantation failure in pregnant mice. Although mice and dogs have different types of chorioallantoic placentas (Loke, 1982), the various numbers of placental layers and types of placental attachments found between species might play a role in the probability of congenital transmission of leishmaniasis.

Zoonotic visceral leishmaniasis is caused by both *L. infantum* and *Leishmania chagasi* (Mauricio et al., 1999). The finding that a North American isolate of *L. infantum* is transmitted vertically is in contrast to previous work that inferred maternal transmission of *L. chagasi* does not occur in dogs, even though some placental tissue was PCR positive (Andrade et al., 2002). Rare cases of vertical transmission of *L. infantum* from the Mediterranean Basin have been previously described in humans and in dogs (Mancianti and Sozzi, 1995; Meinecke et al., 1999; Masucci et al., 2003). *Leishmania infantum* MON-1 is the most common zymodeme causing canine leishmaniasis in southern Europe and is the same type identified in infected foxhounds in the United States (Gaskin et al., 2002). It is possible that the strain of *L. infantum* infecting U.S. foxhounds has the unique ability to routinely cross the placenta. This mode of transmission could be responsible for maintaining the infections in this group of dogs. Additional research with larger numbers of dogs experimentally or naturally infected with the North American strain of *L. infantum* need to be conducted to determine the efficiency of maternal transmission and parasite survival in transplacentally infected puppies.

LITERATURE CITED

- ANDRADE, H. M., V. DE P. C. P. DE TOLEDO, M. J. MARQUES, J. C. F. SILVA, W. L. TAFURI, W. MAYRINK, AND O. GENARO. 2002. *Leishmania (Leishmania) chagasi* is not vertically transmitted in dogs. *Veterinary Parasitology* **103**: 71-81.
- CATONE, G., G. MARINO, G. POGLAYEN, M. GRAMICCIA, A. LUDOVISI, AND A. ZANGHI. 2003. Canine transmissible venereal tumour parasitized by *Leishmania infantum*. *Veterinary Research Communications* **27**: 549-553.
- ELTOUM, I. A., E. E. ZILJSTRA, M. S. ALI, H. W. GHALIB, M. M. H. SATTI, B. ELTOUM, AND A. M. EL-HASSAN. 1992. Congenital kala-azar and leishmaniasis in the placenta. *American Journal of Tropical Medicine and Hygiene* **46**: 57-62.
- GASKIN, A. A., P. SCHANTZ, J. JACKSON, A. BIRKENHEUER, L. TOMLINSON, M. GRAMICCIA, M. LEVY, F. STEURER, E. KOLLMAR, B. C. HEGARTY, A. AHN, AND E. B. BREITSCHWERDT. 2002. Visceral leishmaniasis in a New York foxhound kennel. *Journal of Veterinary Internal Medicine* **16**: 34-44.
- GIGER, U., D. A. OAKLEY, S. D. OWENS, AND P. SCHANTZ. 2002. *Leish-*

- mania donovani* transmission by packed RBC transfusion to anemic dogs in the United States. *Transfusion* **42**: 381–383.
- KRISHNAN, L., L. J. GUILBERT, A. S. RUSSELL, T. G. WEGMANN, T. R. MOSMANN, AND M. BELOSEVIC. 1996. Pregnancy impairs resistance of C57BL/6 mice to *Leishmania major* infection and causes decreased antigen-specific responses and increased production of T helper 2 cytokines. *Journal of Immunology* **156**: 644–652.
- , T. G. WEGMANN, M. BELOSEVIC, AND T. R. MOSMANN. 1996. T helper 1 response against *Leishmania major* in pregnant C57BL/6 mice increases implantation failure and fetal resorptions. *Journal of Immunology* **156**: 653–662.
- LAINSON, R., AND R. S. BRAY. 1964. Transmission of *Leishmania mexicana* among laboratory hamsters in the absence of an insect vector. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **58**: 287.
- LOKE, Y. W. 1982. Transmission of parasites across the placenta. *Advances in Parasitology* **21**: 155–228.
- LOW, G. C., AND W. E. COOKE. 1926. A congenital case of kala-azar. *The Lancet* **11**(ii): 1209–1211.
- MANCIANTI, F., AND S. SOZZI. 1995. Isolation of *Leishmania* from a newborn puppy. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **89**: 402.
- MASUCCI, M., M. DE MAJO, R. B. CONTARINO, G. BORRUTO, F. VITALE, AND M. G. PENNISI. 2003. Canine leishmaniasis in the newborn puppy. *Veterinary Research Communications* **27**(suppl. 1): 771–774.
- MAURICIO, I. L., M. K. HOWARD, AND M. A. MILES. 1999. Genomic diversity in the *Leishmania donovani* complex. *Parasitology* **119**: 237–246.
- MEINECKE, C. K., J. SCHOTTELIUS, L. OKSAM, AND B. FLEISCHER. 1999. Congenital transmission of visceral leishmaniasis (kala azar) from an asymptomatic mother to her child. *Pediatrics* **104**: e65.
- NIETO, C. G., M. GARCIA-ALONSO, J. M. REQUENA, C. MIRON, M. SOTO, C. ALONSO, AND I. NAVARRETE. 1999. Analysis of the humoral immune response against total and recombinant antigens of *Leishmania infantum*: Correlation with disease progression in canine experimental leishmaniasis. *Veterinary Immunology and Immunopathology* **67**: 117–130.
- NUWAYRI-SALTI, N., AND H. F. KHANSA. 1985. Direct non-insect-vector transmission of *Leishmania* parasites in mice. *International Journal of Parasitology* **15**: 497–500.
- NYAKUNDI, P. M., R. MUIGAI, J. B. O. WERE, C. N. OSTER, G. S. CACHIHI, AND G. KIRIGI. 1988. Congenital visceral leishmaniasis: Case report. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **82**: 564.
- OWENS, S. D., D. A. OAKLEY, K. MARRYOTT, W. HATCHETT, R. WALTON, T. J. NOLAN, A. NEWTON, F. STEURER, P. SCHANTZ, AND U. GIGER. 2001. Transmission of visceral leishmaniasis through blood transfusions from infected English foxhounds to anemic dogs. *Journal of the American Veterinary Medical Association* **219**: 1076–1083.
- PINELLI, E., R. KILLICK-KENDRICK, J. WAGENAAR, W. BERNADINA, G. DEL REAL, AND J. RUITENBERG. 1994. Cellular and humoral immune responses in dogs experimentally and naturally infected with *Leishmania infantum*. *Infection and Immunity* **62**: 229–235.
- RAGHUPATHY, R. 1997. Th1-type immunity is incompatible with successful pregnancy. *Immunology Today* **18**: 478–482.
- RODGERS, M. R., S. J. POPPER, AND D. F. WIRTH. 1990. Amplification of kinetoplast DNA as a tool in the diagnosis of *Leishmania*. *Experimental Parasitology* **71**: 267–275.
- ROSYPAL, A. C., G. C. TROY, A. M. ZAJAC, R. B. DUNCAN, JR., K. WAKI, K.-P. CHANG, AND D. S. LINDSAY. 2003. Emergence of zoonotic canine leishmaniasis in the United States: Isolation and immunohistochemical detection of *Leishmania infantum* from foxhounds from Virginia. *Journal of Eukaryotic Microbiology* **50**: S691–S693.
- SANTOS-GOMES, G. M., R. ROSA, C. LEANDRO, S. CORTES, P. ROMAO, AND H. SILVEIRA. 2002. Cytokine expression during the outcome of canine experimental infection by *Leishmania infantum*. *Veterinary Immunology and Immunopathology* **88**: 21–30.
- WEGMANN, T. G., H. LIN, L. GUILBERT, AND T. R. MOSMANN. 1993. Bidirectional cytokine interactions in the maternal–fetal relationship: Is successful pregnancy a Th2 phenomenon? *Immunology Today* **14**: 353–356.
- YADAV, T. P., H. GUPTA, U. SATTEYA, R. KUMAR, AND V. MITTAL. 1989. Congenital kala-azar. *Annals of Tropical Medicine and Parasitology* **83**: 535–537.
- YOUNG, D. G., AND P. V. PERKINS. 1984. *Phlebotomine* sand flies of North America (Diptera: Psychodidae). *Mosquito News* **44**: 264–304.

The Finding of *Echinostoma* (Trematoda: Digenea) and Hookworm Eggs in Coprolites Collected From a Brazilian Mummified Body Dated 600–1,200 Years Before Present

L. Sianto, K. J. Reinhard*, M. Chame, S. Chaves, S. Mendonça, M. L. C. Gonçalves, A. Fernandes, L. F. Ferreira, and A. Araújo, Escola Nacional de Saude Publica—Fundacao Oswaldo Cruz, Rio de Janeiro, Brazil; *School of Natural Resources Sciences, University of Nebraska—Lincoln, Lincoln, Nebraska 68588-0340. e-mail: adauto@ensp.fiocruz.br

ABSTRACT: The identification of parasites from ancient cultures expands our list of parasites infective to extant humans. A partially mummified human body from the archeological site of Lapa do Boquete, Minas Gerais State, Brazil, was recently discovered. It was interred between 600 and 1,200 yr ago. Dietary analysis showed that the mummified body was from a society that had a mixed subsistence of agriculture and gathering of wild foods. Coprolites from the body contained numerous helminth eggs. The eggs were identified as those of *Echinostoma* sp. and hookworm. Hookworm infection in pre-Columbian populations is already established, but this is the first evidence of *Echinostoma* sp. eggs found in human coprolites. The diagnosis of a true infection, as opposed to false parasitism, is discussed. The possibility of *Echinostoma ilocanum* infection is discussed, as this is a common species found in humans in the Asiatic region, which could have been introduced in South America in the pre-Columbian period. Alternative possibilities are also considered, including indigenous Brazilian *Echinostoma* species.

One of the most significant contributions of archeology to parasitology is the documentation of parasite species infective to ancient humans that are not known from the present clinical literature. In some cases, false parasitism is implicated, such as the find of *Cryptocotyle lingua* eggs in an Alaskan Yupik mummy (Zimmerman, 1998). False parasitism occurs when parasite eggs are passed in the feces of a subject who is not infected with the parasite. In other cases, real infection is implicated, such as the discovery of acanthocephalan eggs in archeological sites of the Great Basin of North America (Fry, 1970). Diagnosing infection from the archeological record is only possible when the physical remains analyzed are of human origin and when the dietary practices of the human population are known (Reinhard et al., 1987; Reinhard, 1988). If these 2 criteria are met, then the possibility of confusing false parasitism with true infection can be reduced.

Archeologists recently excavated the cave, Lapa do Boquete. This site is situated in the Peruagu River Valley of northern Minas Gerais State, Brazil. The region is characterized by cerrado vegetation com-