

Table of Contents		Page #
Abstract		ii
Acknowledgments		iv
List of Figures		vii
List of Tables		viii
Chapter I	Introduction and Review of Literature	1
	• Cotton	1
	• Weed Control in Cotton	2
	• Objectives	13
	• Literature Cited	14
Chapter II	Influence of Adjuvants on Cotton Response to Postemergence Applications of CGA 362622	21
	• Abstract	21
	• Introduction	22
	• Materials and Methods	23
	• Results and Discussion	26
	• Literature Cited	29
Chapter III	CGA 362622 Plus Pyriithiobac Combinations for Broadleaf Weed Control in Cotton (<i>Gossypium hirsutum</i>)	37
	• Abstract	37
	• Introduction	38
	• Materials and Methods	40
	• Results and Discussion	43
	• Literature Cited	48
Chapter IV	Combinations of CGA 362622 and Glyphosate for Weed Control in Glyphosate-Tolerant Cotton	60
	• Abstract	60
	• Introduction	61
	• Materials and Methods	63
	• Results and Discussion	65

	• Literature Cited	71
Chapter V	Combinations of CGA 362622 and Bromoxynil for Broadleaf Weed Control in Bromoxynil-Resistant Cotton	82
	• Abstract	82
	• Introduction	83
	• Materials and Methods	85
	• Results and Discussion	86
	• Literature Cited	92
Chapter VI	Absorption, Translocation, and Metabolism of CGA 362622 in Cotton, Spurred Anoda (<i>Anoda cristata</i>), and Smooth Pigweed (<i>Amaranthus hybridus</i>)	100
	• Abstract	100
	• Introduction	101
	• Materials and Methods	103
	• Results and Discussion	106
	• Literature Cited	110
Chapter VII	Summary	119
Vita		121

List of Figures

Chapter	Figure	Title	Page #
I	1.1.A	Chemical structure of pyriithiobac.	20
	1.1.B	Structure and functional groups of a general sulfonylurea Molecule.	20
	1.1.C	Chemical structure of CGA 362622.	20
VI	6.1	Chemical structures of A) CGA 362622, B) nicosulfuron, and C) rimsulfuron.	113
	6.2	Negative image of autoradiograph of A) cotyledon- to one-leaf cotton, B) two- to three-leaf cotton, C) spurred anoda, and D) smooth pigweed harvested 72-h after treatment.	117
	6.3	Metabolism of parent molecule (Rf = 0.6) and formation of metabolites in the treated leaves of cotyledon cotton (<i>Gossypium hirsutum</i>), 2-lf cotton, spurred anoda (<i>Anoda cristata</i>), and smooth pigweed (<i>Amaranthus hybridus</i>). The LSD value compares means across species and time (P = 0.05).	118

List of Tables

Chapter	Table	Title	Page #
II	2.1	Rainfall by month at Eastern Shore AREC near Painter, VA, in 1999, 2000, and 2001.	31
	2.2	Effect of CGA 362622 rate and adjuvant applied postemergence on cotton response from 1999 through 2001 near Painter, VA.	32
	2.3	Analysis of main and interaction effects of CGA 362622 rate and adjuvant on cotton height and significance of time interval on height means from 1999 through 2001 near Painter, VA.	33
	2.4	Effect of CGA 362622 rate and adjuvant applied postemergence on cotton height from 1999 through 2001 near Painter, VA.	34
	2.5	Effect of CGA 362622 rate and adjuvant applied postemergence on cotton lint yield and fiber quality from 1999 through 2001 near Painter, VA.	35
	2.6	Effect of CGA 362622 rate and adjuvant applied postemergence on cotton response, cotton leaf area, and cotton shoot dry weight at 3 WAT in the greenhouse.	36
III	3.1	Height of weed species from 1999, 2000, and 2001 measured at herbicide application near Painter, VA.	51
	3.2	Effect of CGA 362622 rate and pyrithiobac rate applied postemergence on cotton response pooled over 1999, 2000, and 2001 near Painter, VA.	52
	3.3	Analysis of main and interaction effects of CGA 362622 rate and pyrithiobac rate on control of velvetleaf, jimsonweed, ivy-leaf morningglory, and tall morningglory at 56 DAT from 1999, 2000, and 2001 near Painter, VA.	53

	3.4	Analysis of main and interaction effects of CGA 362622 rate and pyrithiobac rate on control of common ragweed, spurred anoda, common lambsquarters, annual morningglory species, and common cocklebur at 56 DAT from 1999, 2000, and 2001 near Painter, VA.	54
	3.5	Effect of CGA 362622 rate and pyrithiobac rate applied postemergence on weed control 56 DAT from 1999, 2000, and 2001 near Painter, VA.	55
	3.6	Effect of CGA 362622 rate and pyrithiobac rate applied postemergence on cotton lint yield and fiber quality from 2000 and 2001 near Painter, VA.	56
	3.7	Control of jimsonweed in the greenhouse with postemergence applications of CGA 362622 and pyrithiobac.	57
	3.8	Control of velvetleaf in the greenhouse with postemergence applications of CGA 362622 and pyrithiobac.	58
	3.9	Control of spurred anoda in the greenhouse with postemergence applications of CGA 362622 and pyrithiobac.	59
IV	4.1	Height of weed species present at herbicide application in 1999, 2000, and 2001.	74
	4.2	Analysis of main and interaction effects of CGA 362622 rate and glyphosate rate on percent cotton response from 1999, 2000, and 2001 near Painter, VA.	75
	4.3	Effect of CGA 362622 rate and glyphosate rate applied postemergence on cotton response from 1999, 2000, and 2001 near Painter, VA.	76
	4.4	Analysis of main and interaction effects of CGA 362622 rate and glyphosate rate on visual weed control 56 DAT from 1999, 2000, and 2001 near Painter, VA.	77
	4.5	Analysis of main and interaction effects of CGA 362622 rate and glyphosate rate on visual weed control 56 DAT in 1999, 2000, and 2001 near Painter, VA.	78

	4.6	Effect of glyphosate and CGA 362622 applied postemergence on weed control at 56 DAT in 1999, 2000, and 2001 near Painter, VA.	79
	4.7	Effect of CGA 362622 rate and glyphosate rate applied postemergence on weed control 56 DAT from 1999, 2000, and 2001 near Painter, VA.	80
	4.8	Effect of glyphosate and CGA 362622 applied postemergence on cotton lint yield and fiber quality from 1999 through 2001 near Painter, VA.	81
V	5.1	Height of weed species from 1999, 2000, and 2001 measured at herbicide application near Painter, VA.	94
	5.2	Effect of CGA 362622 rate and bromoxynil rate applied postemergence on cotton response from 1999, 2000, and 2001 near Painter, VA.	95
	5.3	Analysis of main and interaction effects of CGA 362622 rate and bromoxynil rate on visual weed control 28 DAT in 1999, 2000, and 2001 near Painter, VA.	96
	5.4	Effect of CGA 362622 rate and bromoxynil rate applied postemergence on weed control 28 DAT from 1999, 2000, and 2001 near Painter, VA.	97
	5.5	Effects of bromoxynil and CGA 362622 applied postemergence on weed control at 28 DAT in 1999, 2000, and 2001 near Painter, VA.	98
	5.6	Effects of bromoxynil and CGA 362622 applied postemergence on cotton lint yield and fiber quality from 1999 through 2001 near Painter, VA.	99
VI	6.1	Absorption and translocation of foliar applied ¹⁴ C-CGA 362622 in cotton (<i>Gossypium hirsutum</i>), spurred anoda (<i>Anoda cristata</i>), and smooth pigweed (<i>Amaranthus hybridus</i>).	114

6.2	Metabolism of foliar-applied ^{14}C -CGA 362622 in treated leaf extracts of cotton (<i>Gossypium hirsutum</i>), spurred anoda (<i>Anoda cristata</i>), and smooth pigweed (<i>Amaranthus hybridus</i>) at 6, 24, 48, and 72-h after treatment.	116
-----	---	-----