

## List of Tables:

<b>Table 2.1.</b> Summary of preparation conditions and the surface properties of the oxynitrides synthesized by TPR.	13
<b>Table 2.2.</b> Crystal structure and lattice parameter for the synthesized niobium oxynitride. Comparison with some others niobium compound.	16
<b>Table 3.1.</b> Characteristics of catalysts.	47
<b>Table 4.1.</b> Reactivities of different amines.	72
<b>Table 4.2.</b> List of retention times and relative response factors of all reagents and products.	82
<b>Table 4.3.</b> Characteristics of catalysts.	84
<b>Table 5.1.</b> Pyridine HDN results for Mo <sub>2</sub> C, NbMo <sub>2</sub> C, and MoS <sub>2</sub> /SiO <sub>2</sub> .	131

## List of Schemes:

<b>Scheme 4.1.</b> $\beta$ -Elimination.	70
<b>Scheme 4.2.</b> Nucleophilic substitution.	70
<b>Scheme 4.3.</b> E1 and S <sub>N</sub> 1.	71
<b>Scheme 4.4.</b> Mechanism proposed for the HDN of a pentylamine.	100
<b>Scheme 4.5.</b> Mechanism proposed for the isomerization of olefins.	103
<b>Scheme 4.6.</b> Mechanism for initial HDN of cyclic amines.	103
<b>Scheme 5.1.</b> Reaction scheme for HDN of pyridine.	113
<b>Scheme 5.2.</b> Reaction scheme for HDN of piperidine.	124

<b>Scheme 5.3.</b> Reaction pathway for formation of sulfur products.	124
<b>Scheme 5.4.</b> Rake mechanism for the reaction network.	134

**List of Figures:**

<b>Figure 2.1.</b> Synthesis traces of niobium oxynitride.	14
<b>Figure 2.2.</b> Activation energy analysis using the Redhead equation. Plot $\ln(T_f^2/\beta)$ vs. $1/T_f$ .	17
<b>Figure 2.3.</b> X-ray diffraction patterns of reference samples and the oxynitrides synthesized using a molar space velocity of $1640 \text{ h}^{-1}$ and the indicated heating rates.	18
<b>Figure 2.4.</b> Points A (1020 K), B (1065 K) C (1085 K) D (1100 K) and FP (final product at 1110 K) in the water formation profile where the nitridation was stopped ( $\beta = 0.0500 \text{ K s}^{-1}$ ).	19
<b>Figure 2.5.</b> Development of the surface area during synthesis ( $\beta = 0.0500 \text{ K s}^{-1}$ ).	20
<b>Figure 2.6.</b> X-ray diffraction of the intermediates of reaction obtained by interruption of the nitridation process ( $\beta = 0.0500 \text{ K s}^{-1}$ ) in different points: (A) 1020 K; (B) 1065 K; (C) 1085 K; (D) 1100 K. Comparison with $\text{Nb}_2\text{O}_5$ , $\text{NbO}_2$ , $\text{NbO}$ and the final product (FP).	21
<b>Figure 2.7.</b> SEM pictures of the starting material, $\text{Nb}_2\text{O}_5$ , final product ( $\text{NbN}_x\text{O}_y$ ) and intermediates obtained by interruption of the nitridation process ( $\beta = 0.0500 \text{ K s}^{-1}$ ).	23
<b>Figure 2.8.</b> Results for the elemental analysis of nitrogen of the intermediates (A, B, C, D) and final product (FP).	24
<b>Figure 2.9.</b> Temperature programmed reduction experiment to $T = 1373 \text{ K}$ .	25
<b>Figure 2.10.</b> X-ray diffraction of the fresh catalyst and the spent catalyst.	26

<b>Figure 2.11.</b> Comparison of HDN activity of the catalyst at 3.1 MPa and 643 K [26].	27
<b>Figure 3.1.</b> Mass Spectrometer signal during carburization.	48
<b>Figure 3.2.</b> XRD patterns of the supported bimetallic oxide and oxycarbide.	49
<b>Figure 3.3.</b> XPS results the supported bimetallic oxycarbide.	51
<b>Figure 3.4.</b> C K-edge NEXAFS features of the bimetallic and monometallic oxycarbides.	54
<b>Figure 3.5.</b> C K-edge NEXAFS features of the Nb-Mo supported oxycarbides.	55
<b>Figure 3.6.</b> O K-edge NEXAFS features of the bimetallic and monometallic oxycarbides.	56
<b>Figure 3.7.</b> O K-edge NEXAFS features of the Nb-Mo supported oxycarbides.	59
<b>Figure 3.8.</b> Comparison of HDN and HDS activity at 3.1 MPa and 643 K.	60
<b>Figure 3.9.</b> Effect of Mo/Nb ratio over the activity.	61
<b>Figure 3.10.</b> C K-edge NEXAFS features of the Nb-Mo supported oxycarbides before and after reaction.	62
<b>Figure 4.1.</b> Typical chromatogram of a liquid-sample from <i>n</i> -pentylamine reaction.	81
<b>Figure 4.2.</b> X-ray diffraction of Nb-Mo oxycarbide and its parent monometallic carbides.	83
<b>Figure 4.3.</b> Conversion of <i>tert</i> -pentylamine, <i>n</i> -pentylamine, and <i>neo</i> -pentylamine over: a) Mo <sub>2</sub> C; b) NbMo <sub>2</sub> C; c) MoS <sub>2</sub> /SiO <sub>2</sub> .	88
<b>Figure 4.4.</b> Conversion and product distribution for the <i>n</i> -pentylamine reaction over: a) Mo <sub>2</sub> C; b) NbMo <sub>2</sub> C; c) MoS <sub>2</sub> /SiO <sub>2</sub> .	89
<b>Figure 4.5.</b> Conversion and product distribution for the <i>tert</i> -pentylamine reaction over: a) Mo <sub>2</sub> C; b) NbMo <sub>2</sub> C; c) MoS <sub>2</sub> /SiO <sub>2</sub> .	90
<b>Figure 4.6.</b> Conversion and product distribution for the <i>neo</i> -pentylamine reaction over: a) Mo <sub>2</sub> C; b) NbMo <sub>2</sub> C; c) MoS <sub>2</sub> /SiO <sub>2</sub> .	91

<b>Figure 4.7.</b> TPD curves for ethylamine in Mo <sub>2</sub> C, NbMo <sub>2</sub> C, and MoS <sub>2</sub> /SiO <sub>2</sub> : a) ethylamine (m=30), b) ethylene (m=28), and c) ammonia (m=17) were observed during the desorption.	93
<b>Figure 4.8.</b> Comparison of the DRIFT spectra of ethylamine in the gas-phase and adsorbed on Mo <sub>2</sub> C, NbMo <sub>2</sub> C, and MoS <sub>2</sub> /SiO <sub>2</sub> .	94
<b>Figure 4.9.</b> Relationship between the number of moles of ethylamine converted during TPD and the specific rate of reaction of <i>ter</i> -pentylamine on Mo <sub>2</sub> C, NbMo <sub>2</sub> C, and MoS <sub>2</sub> /SiO <sub>2</sub> .	102
<b>Figure 4.10.</b> Overall turnover rates at 500 K of <i>tert</i> -pentylamine, <i>n</i> -pentylamine, and <i>neo</i> -pentylamine on Mo <sub>2</sub> C, NbMo <sub>2</sub> C, and MoS <sub>2</sub> /SiO <sub>2</sub> based on the number of sites measured by: a) CO uptake for Mo <sub>2</sub> C, NbMo <sub>2</sub> C, and O <sub>2</sub> uptake for MoS <sub>2</sub> /SiO <sub>2</sub> from Table 2; b) moles of ethylamine converted from Figure 8.	106
<b>Figure 5.1.</b> Conversion of <i>n</i> -pentylamine, piperidine, and pyridine over: a) Mo <sub>2</sub> C; b) NbMo <sub>2</sub> C; c) MoS <sub>2</sub> /SiO <sub>2</sub> .	118
<b>Figure 5.2.</b> Conversion and product distribution for the <i>n</i> -pentylamine reaction over: a) Mo <sub>2</sub> C; b) NbMo <sub>2</sub> C; c) MoS <sub>2</sub> /SiO <sub>2</sub> .	120
<b>Figure 5.3.</b> Conversion and product distribution for the piperidine reaction over: a) Mo <sub>2</sub> C; b) NbMo <sub>2</sub> C; c) MoS <sub>2</sub> /SiO <sub>2</sub> .	121
<b>Figure 5.4.</b> Conversion and product distribution for the pyridine reaction over: a) Mo <sub>2</sub> C; b) NbMo <sub>2</sub> C; c) MoS <sub>2</sub> /SiO <sub>2</sub> .	122
<b>Figure 5.5.</b> Piperidine product distribution at constant conversion levels: a) 10% conversion; b) 30% conversion.	126

<b>Figure 5.6.</b> Pyridine product distribution at constant conversion levels: a) 20% conversion; b) 40% conversion.	127
<b>Figure 5.7.</b> $\ln (r)$ vs $1/T$ plots for: a) $\text{Mo}_2\text{C}$ ; b) $\text{NbMo}_2\text{C}$ ; c) $\text{MoS}_2/\text{SiO}_2$ .	133
<b>Figure 5.8.</b> $\ln (r')$ vs $1/T$ plots for: a) $\text{Mo}_2\text{C}$ ; b) $\text{NbMo}_2\text{C}$ ; c) $\text{MoS}_2/\text{SiO}_2$ .	137
<b>Figure 5.9.</b> Energy diagram sketch.	138