



Tips for Profitable Variety Selection: How to Use Data From Different Types of Variety Trials

Wade E. Thomason, Professor and Extension Grain Crops Specialist, Crop and Soil Environmental Sciences

David Holshouser, Associate Professor, Tidewater Agricultural Research and Extension Center

Keith Balderson, Extension Agent, Agriculture and Natural Resources, Crop and Soil Sciences, VCE, Essex County Office

David Moore, Extension Agent, Agriculture and Natural Resources, Crop and Soil Sciences, VCE Middlesex County Office

Eric Bowen, Extension Agent, Agriculture and Natural Resources, Family and Consumer Sciences, VCE Prince Edward County Office

Glenn Chappell II, Assistant Professor, VSU

Bobby Clark, Extension Agent, Agriculture and Natural Resources, Crop and Soil Sciences, VCE Shenandoah County Office

Paul Davis, Extension Agent, Agriculture and Natural Resources, Crop and Soil Sciences, VCE New Kent County Office, Retired

Watson Lawrence, Extension Agent, Agriculture and Natural Resources, Crop and Soil Sciences, VCE Chesapeake Office

Mike Parrish, Agriculture and Natural Resources, Crop and Soil Sciences, VCE Dinwiddie County Office

Scott Reiter, Extension Agent, Agriculture and Natural Resources, Crop and Soil Sciences, VCE Prince George County Office

Sam Johnson, Extension Agent, Agriculture and Natural Resources, Crop and Soil Sciences, VCE Westmoreland County Office, Retired

Glenn Slade, Extension Agent, Agriculture and Natural Resources, Crop and Soil Sciences, VCE Surry County Office

Kelvin Wells, Extension Agent, Agriculture and Natural Resources, Crop and Soil Sciences, VCE Sussex County Office

Matthew Yancey, Extension Agent, Agriculture and Natural Resources, Forestry, VCE Rockingham County Office

Keith Burgess, Monacan SWCD

Introduction

Selecting an appropriate, high-yielding variety is one of the most important management decisions that producers make. Yield potential is clearly important, but the decision is complicated by such factors as the cropping system, the need for disease resistance, end-use quality goals, year-to-year climatic variation, and the need to select multiple varieties in order to reduce risk by spreading out flowering and maturity dates.

Choosing varieties based on these many variables dictates that a significant amount of information be available about each and every one. This information is available in summaries of variety performance trials. Proper use of variety test performance data is the first step in making these important variety decisions. But in order to make proper use of the data, an understanding of the types of variety trials that are most commonly conducted and the pros and cons of each is needed.

Types of Variety Tests

“This variety yielded better on my farm than this other one.” or “This variety performed well for me last year.”

There are many, many factors that could cause either of these statements to be true: year-to-year weather variation, yield

potential differences in and between fields, planting date differences, etc.

Yearly or farm-specific evidence involves a single observation with no direct comparison to other varieties. This information is not without value, however. Most scientifically sound variety trials are designed and sited to avoid major same-soil variability along with pest or previous crop management problems that can and usually will be encountered on an individual’s farm. So, these types of observations about variety performance should be used in conjunction with other variety tests when making a final decision.

Split-Field Comparisons



Split-field comparisons are planned so that two (or sometimes more) varieties are planted side by side in the same or adjacent fields. These tests allow for a “real world” assessment using farm equipment and large areas that should represent the range of soils and crop management scenarios that would normally be seen. The variability in this type of test can also lead to either inaccurate comparisons or difficulty in making comparisons.

For example, in the fields shown in figure 1A, it is apparent that there are soybean growth differences between the two sides of this field, which are separated by a line. Those differences could be due to soils or they might be due to historical

management practices. There's a farmstead within the field planted to variety A. If, for example, this is an animal operation, guess which field might have received the bulk of the manure.

In figure 1B, where varieties A and B are planted side by side in the same field, there is similar variation in crop growth. If the entire area for each variety is harvested and compared on the basis of yield, is the difference due to variety or to field variation? The danger of this type of test is that the comparison may favor field variability rather than variety differences.

It is true that a yield monitor that logs in yield data as the crop is harvested can give some good information about how a variety changes with soil type, but there is still only one side-by-side comparison of varieties.




Figures 1A-B. Two examples of how split-field comparisons could be conducted.

On-Farm Tests With Multiple Varieties

Unreplicated Strip Plots

An example of unreplicated strip-plot trials is shown in figures 2A-B. Varieties are planted in an adjacent pattern, usually the length of the field. The width of each strip for a variety may be a whole planter-/drill-width or a portion of the total. The term “unreplicated” indicates that each variety is planted once at each site. Studies that are “replicated” have each variety planted multiple (generally three or more) times in each field. Unreplicated trials can be a good source of information because they combine the use of farmer equipment and relatively large plot size with the testing of a number of varieties in the same field. The major problem, again, is field variability. It is apparent from figure 2 that there are more problem areas with reduced soybean growth in the part of the field where varieties 1-15 are planted.

One way to avoid this complication is to use one variety as a repeating check (the “C” in figure 2). With these checks, we can partially measure the variability as we move across the



Figures 2A-B. Unreplicated strip plots.

Interpreting Data From Variety Trials

There are two important points to understand when using the data from any variety trial:

1. The highest-yielding variety from a single location will rarely be the highest-yielding variety the following year at that location.
2. The highest-yielding variety selected from multiple location averages will usually be a variety that will yield among the best the following year at any location.

For example, table 1 shows relative yields of soybean varieties over three locations and three years of testing. The first two columns identify the location and the number of years used to choose the highest-yielding variety, respectively. The third column is the relative yield of this highest-yielding variety at a particular location based on one, two, or three years of data. The variety chosen with only one year of data was not always (and not likely) the highest-yielding variety chosen when using two or three years of data. The next columns (4-7) show relative yield of that variety at each of the three testing locations and the average over locations in the following year.

Let's look at an example. If we chose a variety with only one year of data at the Suffolk location (relative yield = 114), the next year that variety had a relative yield in the Suffolk test of 100, meaning its performance was as good as the average for all entries in the test. Even with two or three years of data

for this variety at Suffolk, its performance the following year at Suffolk was not good. However, using the multi-location average data resulted in much greater predictability of performance the following year. In fact, using the multilocation data to choose a variety always gave at least 100 percent of the test average the following year. In fact, when all locations and years were considered, this variety had an average relative yield of 105 percent.

Varieties that have performed well under diverse conditions, as evidenced over locations and years, are likely to perform well again. Depending on a grower's location, it should be kept in mind that additional performance data about the same variety may be available from other state or private variety testing programs.

Conclusion

Growers and advisers should utilize multilocation and multiyear performance data to develop a list of the best-suited varieties. Then, once several varieties are selected, performance of those varieties in the closest regional test or from on-farm testing should be considered.

After identifying a group of varieties with high grain-yield potential and stable performance across sites and years, the variety selection process should continue with consideration given to secondary characteristics such as test weight, disease resistance, lodging, height, maturity, and end-use quality.

Table 1. Example of Virginia soybean variety relative yield performance (as a percentage) over multiple sites and years.

Data set location	Years in data set	Relative yield for highest-yielding variety in data set	Relative yield for highest-yielding variety the following year at each location			
			Orange	Suffolk	Warsaw	Average the following year
Orange	1	114	105	102	101	103
	2	114	99	108	100	104
	3	108	99	96	94	97
Suffolk	1	114	106	100	93	100
	2	113	85	84	98	90
	3	114	109	96	94	100
Warsaw	1	117	98	89	101	95
	2	116	101	107	108	108
	3	112	105	102	101	103
Multilocation	1	110	101	107	100	103
	2	109	103	108	108	106
	3	108	102	103	109	105